

SUMMARY

Evaluation of electromyography in patients with long-term low-back pain and in healthy subjects

Created by Britt Elfving

- Subject** : Evaluation of electromyography
in patients with long-term low-back pain
and in healthy subjects
- Subject Alt** : Evaluation of electromyography
in patients with long-term low-back pain
and in healthy subjects
- Keyword** : Disability, electromyography, exponential, force level, gender, healthy subjects,
interelectrode distance, low-back pain, erector spinae, maximal voluntary contraction, muscle
fatigue, recovery, reliability, validity.

Description :

Lumbar muscle function is considered to be an important component of longterm low-back pain. The change in the median frequency of the power spectrum of the electromyographic (EMG) signal is commonly used to estimate muscle fatigue.

Aim: The main purpose was to evaluate a test method to estimate lumbar muscle fatigue using frequency analysis of the electromyographic signal during isometric contraction.

Methods: In the different studies, 73 healthy subjects participated; 55 of these constituted a reference group which was compared to 57 patients with long-term low-back pain. The subjects performed a back extensor test in seated position: maximal voluntary contraction (MVC) torque, 45 s isometric fatigue contraction at 80% of MVC, 5 s contractions after 1, 2, 3 and 5 min in the recovery period. Surface EMG was recorded from the lumbar muscles bilaterally at spinal levels L1 and L5. To study reliability, this test was repeated five times by 11 of the healthy subjects and once by 20 of the patients. In a further study, 15 of the healthy subjects performed the fatigue test at 40% and 80% of MVC with both 2 cm and 4 cm interelectrode distances. EMG variables were initial median frequency, median frequency slope (obtained from linear regression of the fatigue contraction) and recovery half-time (obtained from non-linear regression of the recovery data). Activity, participation and other health-related factors were estimated by the patients in five questionnaires.

Results: The reliability was somewhat higher for patients ($ICC > 0.6$) than for healthy subjects. The initial median frequency and, to a greater extent, the slope were sensitive to the exerted torque. The patients differed from the healthy subjects by lower MVC torque, higher initial median frequency at L5, and by flatter slope and longer recovery half-time at both lumbar levels. Using logistic regression entering EMG initial median frequency, slope and recovery half-time in combination with MVC, about 80% of the patients and the healthy subjects could be correctly classified. By analyses of individual linear and non-linear regressions, it was found that patients with not significantly negative slopes and/or not exponential-like EMG recovery had lower self-efficacy and more activity limitations. Female patients had significantly flatter slopes, lower physical functioning and self-efficacy than male patients. The results indicate that the ability to fatigue the lumbar muscles in a 80% MVC contraction might be a healthy sign

Date Create : 16/12/2014
Type : Text
Format : pdf
Language : Indonesian
Identifier : UEU-Master-undergraduate_54
Collection : undergraduate_54
Call Number : 658.1 BEe
Source : magister these management of faculty
Relation Collection Universitas Esa Unggul
COverage : Civitas Akademika Universitas Esa Unggul
Right : copyright2014_Library@esaunggul

Full file - Member Only

If You want to view FullText...Please Register as MEMBER

Contact Person :

Astrid Chrisafi (mutiaraadinda@yahoo.com)

Thank You,

Astrid (astrid.chrisafi@esaunggul.ac.id)

Supervisor