

SUMMARY

PENGARUH TEPID SPONGE BATH DENGAN AIR HANGAT TERHADAP PENURUNAN SUHU TUBUH PADA ANAK USIA TODLER DAN PRA SEKOLAH YANG MENGALAMI DEMAM DI UGD RSIA FAMILY JAKARTA

TEPID EFFECT OF SPONGE BATH WITH WARM WATER TEMPERATURE ON THE DECREASE IN CHILDREN AND PRE SCHOOL toddler EXPERIENCE FEVER AT UGD RSIA FAMILY JAKARTA

Created by Puji Melasi

Subject : antipiretik, efektif, suhu tubuh

Subject Alt : air hangat, demam

Keyword : anak, demam, tepid sponge bath

Description :

Latar Belakan : Studi literature tentang pemberian antipiretik disertai tepid sponge bath dengan air hangat menunjukkan bahwa tindakan ini efektif menurunkan demam dibandingkan jika hanya dengan pemberian antipiretik saja.

Tujuan Penelitian : untuk mengetahui pengaruh pemberian antipiretik disertai tepid sponge bath dengan air hangat terhadap penurunan suhu tubuh anak demam di UGD RSIA Family Jakarta.

Desain Penelitian : yang digunakan adalah quasi experiment pre-post non equifalen control group dengan uji t tidak berpasangan. Sampel 32 responden dengan metode non probability sampling.

Hasil Penelitian : Hasil uji t tidak berpasangan menunjukkan penurunan suhu tubuh post test I nilai pada kelompok kontrol 0.21°C (+ 0.68), kelompok intervensi 0.39°C (+ 0.574), p value 0.086 ($p > 0.05$), post test II nilai pada kelompok kontrol 0.74°C (+ 0.718), kelompok intervensi 0.89°C (+ 0.574), p value 0.193 ($p > 0.05$). Penelitian ini menunjukkan tidak ada perbedaan yang bermakna dalam penurunan suhu tubuh antara kelompok intervensi maupun kelompok kontrol, tetapi ada kecenderungan bahwa pemberian antipiretik yang disertai tepid sponge bath mengalami penurunan suhu tubuh yang lebih besar dibandingkan dengan pemberian antipiretik saja.

Kesimpulan : disarankan anak yang mengalami demam agar mendapatkan tindakan tepid sponge bath karena dapat mengurangi resiko kejang demam.

Description Alt:

Background: The study of literature about giving antipyretics with tepid sponge bath with warm water indicates that this action effectively reduce fever than if only by giving an antipyretic alone.

Objective: To determine the effect of antipyretics with tepid sponge bath with warm water to decrease body temperature in febrile children RSIA Family Emergency Jakarta.

Research Design: quasi experiment used was a pre-post non equifalen control group by unpaired t test. Sample of 32 respondents with a non-probability sampling.

Results: The results of the unpaired t test showed a decrease in body temperature post

I test the value in the control group 12:21 ° C (+ 0.68), the intervention group 12:39 ° C (+ 0.574), p value 0.086 (P> 0.05), post test II values in the control group 0.74 ° C (+ 0.718), intervention group 0.89 ° C (+ 0.574), p value = 0.193 (p> 0.05). this study showed no significant difference in the decrease in body temperature between intervention group and the control group, but there is a tendency that the administration antipyretic with tepid sponge bath body temperature decreased more large compared to administration of antipyretics alone.
Conclusion: The recommended children who have a fever in order to get tepid action sponge bath because it can reduce the risk of febrile seizures.

Contributor : Widaningsih, S.Kp, M.Kep
Date Create : 10/07/2013
Type : Text
Format : pdf
Language : Indonesian
Identifier : UEU-Undergraduate-201033054
Collection : 201033054
Call Number : 610.73
Source : Undergraduate theses hygiene faculty
Relation Collection Universitas Esa Unggul
COverage : Sivitas Akademika Universitas Esa Unggul
Right : Copyright @2013 by UEU Library. This publication is protected by copyright and per obtained from the UEU Library prior to any prohibited reproduction, storage in a re transmission in any form or by any means, electronic, mechanical, photocopying, reco For information regarding permission(s), write to UEU Library

Full file - Member Only

If You want to view FullText...Please Register as MEMBER

Contact Person :

Astrid Chrisafi (mutiaraadinda@yahoo.com)

Thank You,

Astrid (astrid.chrisafi@esaunggul.ac.id)

Supervisor