

Effects of the Methanol extracts Phaleria macrocarpa (Scheff) Boerl fruits on Angiotensin Converting Enzyme (ACE) activity

by Aprilita Rina Yanti Eff

Submission date: 22-Oct-2019 05:47PM (UTC+0700)

Submission ID: 1197946103

File name: B17.pdf (208.82K)

Word count: 3991

Character count: 22293

Effects of the Methanol extracts *Phaleria macrocarpa* (Scheff) Boerl fruits on Angiotensin Converting Enzyme (ACE) activity

Aprilita Rina Yanti¹, Maksum Radji², Abdul Mun'im², FD Suyatna³.

¹ Faculty of Pharmacy University of 17 Agustus 1945 Jakarta, Jakarta Indonesia

² Faculty of Pharmacy, University of Indonesia, Depok, Indonesia.

³ Department of Pharmacology and Therapeutic, Faculty of Medicine, University of Indonesia, Jakarta, Indonesia.

Abstract

The present study was carried out to investigate the antihypertensive effect and Angiotensin Converting Enzyme activity of methanolic extract of *Phaleria macrocarpa* (Scheff) Boerl fruits in . Hypertension was induced experimentally by 10% w/v fructose diet ad libitum for 8 weeks. *P. macrocarpa* was given at the dose 0.5, 1 and 2 g/kg BW start at week 6 until week 8 and 1 g/kg BW for 8 weeks. Blood pressure was measured at baseline and at every weeks for 8 weeks using tail-cuff plethysmography. After completion of treatment schedule rats from each group were anesthetized with urethane (120 mg/100 gm, i. p.). Blood was collected and then centrifugated to obtain plasma and later store at -70°C for analysis activity ACE plasma. The results suggests that methanol extract of *Phaleria macrocarpa* could prevent the development high blood pressure and normalized blood pressure in rat induced by diet rich in fructose. The results tend to suggest that methanol extracts of possesses antihypertensive activity, through inhibition of angiotensin converting enzyme.

Keywords: fructose, hypertension, *P. macrocarpa*, angiotensin converting enzyme

14. INTRODUCTION

Angiotensin-converting enzyme (ACE) is a key enzyme in the renin-angiotensin aldosterone system (RAAS). ACE converts the inactive decapeptide angiotensin I, forming the active octapeptide angiotensin II. Angiotensin II is a main effector of the RAAS and seems to be involved in developing cardiovascular disease. Angiotensin II binds to the angiotensin receptors AT1 and AT2¹. Activation of AT1 receptors is associated with endothelial dysfunction, vasoconstriction, cell proliferation, platelet aggregation, inhibition of nitric oxide synthase, aldosterone release, and increases in reactive oxygen species².

High blood pressure is the leading preventable cause of cardiovascular diseases such as stroke, myocardial infarction, left ventricular hypertrophy, nephropathy, retinopathy and dementia³. In middle and old age

groups, every 20 mmHg of systolic or 10 mmHg diastolic increase in blood pressure above 115 mmHg/75 mmHg results doubling of mortality from ischemic heart disease and stroke⁴.

Fructose induced hypertension model gives clue about the role of dietary changes in hypertension. High fructose diets have been documented to increase blood pressure in experimental rats^{5,6}. Hypertension, a component of metabolic/insulin resistance, impaired glucose tolerance syndrome is an important risk factor for cardiovascular diseases⁷. Insulin resistance is often linked to the macronutrient content in the diet. Oxidative stress, over lipid production plays an important role in development of hypertension and other cardiovascular complications⁸. Feeding of a high fructose diet to rats results in hyperglycemia, hypertriglyceridemia and

hyperinsulinemia. The hypertension is accompanied by several metabolic abnormalities. Insulin resistance and hyperinsulinemia are frequently associated with both clinical and experimental hypertension⁹. Fructose-induced hypertension has been related to volume overload, sodium retention by kidneys and increased sympathetic activity among other things¹⁰. Alterations in renal function may also play an important role in the development of hypertension in fructose-fed rats¹¹.

Several classes of pharmacological agents have been used in the treatment of hypertension. One class of antihypertensive drugs known as angiotensin I converting enzyme (ACE) inhibitors are associated with a low rate of adverse side-effects and are the preferred class of anti-hypertensive agents for treating patients with concurrent secondary diseases¹². The potent ACE inhibitors were frequently derived from food proteins¹³.

Phaleria macrocarpa (Scheff.) Boerl is native plants from Indonesia, which has long been used as folk medicines for treatments various types of diseases such as cancer, liver disorders, heart disease, diabetes, arthritis, kidney disorders, stroke, and high blood pressure¹⁴. Qualitatively this plants showed the presence of alkaloids, saponins, polyphenols, and from fruit contained alkaloids, saponins and flavonoids¹⁵. *Phaleria macrocarpa* was reported to contain phenolic glycoside such as mahkotoside, mangiferin, kaempferol-3-O- β -dglucoside, dodecanoic acid, palmitic acid, ethyl stearate, and sucrose¹⁶. The content of lignans in *Phaleria macrocarpa* (Scheff.) Boerl are pinorensinol, lariciresinol and matairesinol¹⁷. *Phaleria macrocarpa* (Scheff.) Boerl has antidiabetic effects that inhibit alpha-glucosidase and anti-diabetic effect in mice induced streptozotocin.

Our previous study demonstrated that the fruits of *Phaleria macrocarpa* has ACE inhibitory activity with IC₅₀ values in the fruits of Petroleum ether extract, aethyl acetate extract and methanol extract were 162 μ g/²⁶, 139 μ g/mL and 122 μ g/mL respectively¹⁸. The aim of the present study was to evaluate the effect of metanolic extract on systolic and diastolic blood pressure and plasma angiotensin converting enzyme levels in the fructose induced hypertensive rat.

1.2. Material and Methods

1.1.2. Extraction

1000 gram of *P. macrocarpa* fruits powder was put in 2 litre methanol 80% solvent for 7 days at a room temperature with 4 times replications, and then the mixture was filtered and evaporated by rotary vacuum evaporator at 40°C until the concentrated methanol extract is obtained.

1.1.3. Animal and study design

Forty-two adult male Sprague-Dawley rats (aged 3 months), weighing 200 - 280 g were obtained from the Animal Source Unit, Indonesia University. The rats were randomly assigned into seven dietary groups (two control and five experimental groups) comprising of six animals each. Prior study approval was obtained from the Ethics Committee Medical Faculty of Indonesia University. All animal management and procedures were performed in accordance with the recommended guidelines. The rats were kept in stainless-steel cages and maintained at room temperature of 27°C \pm 2°C with a 12 h light-dark cycle. All rats had free access to food and water ad libitum during the study period. Hypertension was induced experimentally by fructose 10% W/V diet ad libitum for 8 weeks. Fructose solution was prepared every two days by dissolving the fructose in distilled water. After one week of acclimatization, each group of rats were fed on the following diets: group I (normal control/ NC), rats received no medication but were given distilled water for drinking for 8 weeks; group II (negative control/ NCG), rats received no medication but were given 10% fructose solution for drinking for 8 weeks; group III, rats received 10% fructose solution for drinking for 8 weeks and received captopril 10 mg/kg BW start at week 6 until week 8 (CG); group IV, rats received 10% fructose solution for drinking for 8 weeks and received methanol extract of *P. macrocarpa* 0.5 g/kg BW start at week 6 until week 8 (PM 0.5); Group V, rats received 10% fructose solution for drinking for 8 weeks and received methanol extract of PM 1 g/kg BW start at week 6 until week 8 (PM1); group VI, rats received 10% fructose solution for drinking for 8 weeks and received methanol extract of PM 2 g/kg BW start at week 6 until week 8 (PM2); group VII, rats received 10% fructose solution for drinking for 8 weeks with methanol extract of PM 1 g/kg BW (PM3). Blood pressure was measured at baseline and at every weeks for 8 weeks using tail-cuff plethysmography. After completion of treatment schedule rats from each group were anesthetized with urethane (120 mg/100 gm, i. p.). Blood was collected and then centrifugated to obtain plasma and later store at -70°C for analysis activity ACE plasma. The activity of ACE was measured using commercially available kit (USCNLife, West Lake, Wuhan, China) following manufacturer's instruction. The intensity of coloured product was measured in a microplate reader 450 nm.

1.3. RESULTS AND DISCUSSION

1.3.1. Blood pressure

Results of blood pressure measurement in all treatment groups showed that systolic blood increase

start at the second week of fructose administration, except in the normal controls and the group that received 10% fructose solution for drinking for 8 weeks with methanol extract of *P. marcocarpha* 1 g/kg BW (Figure 1). Statistical analysis showed a significant difference in systolic blood pressure at week 2 between the normal control group (NG) with the negative control group (NCG), captopril group (CG), PM group at dose 0.5 g / kg BW, PM group at dose 1 g / kg BW and PM group at dose 2 g / kg BW ($p < 0.05$). Systolic blood pressure of all treatment groups at week 0 was comparable, providing 10 % fructose in drinking water starting at week 0 to week 8 increased systolic blood pressure. Systolic blood pressure in group rats that given captopril, PM0,5, PM1 and PM 2 seen decreased and not significantly different with normal control. Systolic blood pressure in rats were given PM 3 was not increase, and showed that methanol extract *P. marcocarpha* can prevent increasing systolic blood pressure.

Measurement of diastolic blood pressure in all treatment groups showed that blood pressure diastolic increase start the second week of fructose administration, except in the normal controls and in group received 10% fructose solution for drinking for 8 weeks with methanol extract of PM 1 g/kg BW (Figure 2). Statistical analysis showed no difference in diastolic pressure in all treatment groups at week -0 and week 1. There was no significant difference in diastolic blood pressure between the NG with CG, PM 1, PM 2 and PM 3 ($p < 0.05$).

Fructose is a mediator of hypertension and hyperuricemia. Various studies have demonstrated administration of a high-fructose in rats leads to the onset of metabolic syndrome which includes hypertension, hyperglycemia, insulin resistance, obesity, and hyperuricemia¹⁹. The development of hypertension occurs in the second week after the administration of 10% fructose in drinks or 60% in the diet²⁰. Increasing production or activity of vasoactive mediators such as endothelin-1 (ET1)²¹, angiotensin II (ANGII)²² and thromboxane A2²³ is the mechanism underlying the relationship between resistance insulin and hypertension.

Studies in animals and humans suggest a link between the renin-angiotensin system to the pathogenesis of insulin resistance. Angiotensin II affects glucose metabolism through its effect on the insulin signaling pathway, decrease tissue blood flow, oxidative stress, increase sympathetic activity and adipogenesis. Increasing of sympathetic nerve activity causes increasing of catecholamine that leads to endothelial dysfunction and elevation in blood pressure²⁴. Angiotensin receptor antagonists prevent Ang II from binding to angiotensin receptors and

thereby inhibit AT1R-induced vasoconstriction, increasing blood flow through vasodilation. ACE inhibitors improve insulin sensitivity through two mechanisms. They reduce ACE-mediated generation of Ang II and they prevent the degradation of bradykinins, prolonging their half-life and allowing relaxation of blood vessels through endothelium-dependent mechanisms to occur²⁵. Insulin resistance also causes increasing in sympathetic nerve activity that causes sodium reabsorption in the kidney and contributes to hypertension²⁶. Fructose cause of hypertension by reducing renal sodium excretion, increasing of sodium absorption in the intestine and kidney, and this process is mediated by a transporter Slc26a6 and Slc2a5²⁷.

SLC26a6 in humans and in mice were also known as PAT1 (Putative Anion Transporter 1) or CFEX (Chloride / Formate Exchanger). There were major apical chloride/ base exchanger in the small intestine and proximal tubule, functioning in exchange $\text{Cl}^-/\text{HCO}_3^-$ and $\text{Cl}^-/\text{oxalate}$ and plays important role in the excretion of salt and bicarbonate in the small intestine. Fructose absorption in the small intestine and kidney proximal tubule is mediated primarily by Glut5 (Slc2a5)²⁷.

Another mechanism of fructose cause hypertension in rats through the formation of aldehyde conjugate levels is the result of the metabolism of fructose. Aldehyde binds to membrane protein sulfhydryl groups interfere calcium channels, thereby increasing the levels of free calcium, peripheral resistance and increased blood pressure²⁷. In vitro studies showed that the methanol extract of mahkota dewa (*P. marcocarpha*) has effect as ACE inhibitors with IC_{50} value of 139.11 ppm¹⁸. Mahkota dewa contain mahkotoside, mangiferin, kaempferol-3-glucoside, dodecanoat acid, palmitic acid, ethyl stearate, sucrose and lignan. The lignans were pinoresinol, lariciresinol and matairesinol¹⁶.

1.3.2. Activity ACE

Mean plasma ACE levels in each treatment group are summarized in Figure 3. Statistical analysis showed that the plasma ACE levels significant difference between the normal control group with the NCG, PM 0.5, PM 1 and PM2 ($p < 0.005$). There was no significant difference between the normal control group and the CG with Group PM 1 (8 weeks delivery) ($p > 0.005$). There was a significant difference between the NCG with NC group, CG, PM 0,5, PM1 and PM2 AND PM3 ($p > 0.05$). There was no significant difference between CG with PM3 ($p < 0.005$) and no significant difference between the group PM1 and PM2 ($p > 0.05$). Plasma ACE levels seen in the NCG 16 times larger than the CN.

The renin-angiotensin system (RAS) plays a pivotal role in the homeostatic control of arterial pressure, tissue perfusion, and extracellular volume leading to the development of hypertension. Dysregulation of the RAS can result in the pathogenesis of hypertension, cardiovascular and renal disorders. Owing to this, blockade of the RAS with angiotensin-converting enzyme inhibitors or angiotensin receptor blockers has become important therapeutic approach in cardiovascular and renal medicine¹. Pharmacological interruption of the RAAS is possible at five major site, renin release from the juxtaglomerular cells, renin catalyzed cleavage of angiotensinogen, ACE conversion of A-I to A-II, AII action at angiotensin receptors and aldosterone action at the mineralocorticoid receptors²⁸.

ACE converts inactive AI to potent vasoconstrictor AII and raising blood pressure. AII increases generation of superoxide free radicals via NADPH/NADH oxidase system. AII has dual role in elevating blood pressure, direct vasoconstrictor effects and increasing production of free radicals which reduce of bioavailability of NO and indirectly attenuating endothelium dependent relaxation responses²⁹.

The mechanisms underlying fructose-induced hypertension are not completely clarified, it has been proposed that elevation of blood pressure in fructose-fed rats is secondary to the development of insulin resistance and hyperinsulinemia. That is, compensatory hyperinsulinemia has been thought to be a cause of hypertension because insulin could cause sodium retention, sympathetic nerve activation, and vascular smooth muscle cell proliferation⁷.

Hypertension after chronic fructose administration is due to the interaction between the ET - 1 system and the renin angiotensin system. ET - 1 is a vasoactive peptide produced by endothelial cells, are working on various tissues endocrine, paracrine or autocrine. ET-1 acts as an endogenous vasoconstrictor and growth factor for vascular smooth muscle cells. On the condition of hyperinsulinemia and insulin resistance, insulin can induce the release of ET-1 and causes an increase in blood pressure and vascular dysfunction³⁰. Studies conducted by Tran et.al in rats fed fructose and given ET-1 receptor antagonists can normalize levels of chronic angiotensin II³¹. Renin-angiotensin system is a mediator in maintaining the blood pressure. Excessive activation of this system will lead to an increase in angiotensin II that causes hypertension and insulin resistance. Ang II may mediate the development of insulin resistance through its vasoconstrictor actions, thereby reducing blood flow and glucose uptake into insulin-sensitive tissues and the findings that Ang II infusion induced endothelial dysfunction, hepatic insulin resistance

and elevations in blood pressure^{19,31}. ACE inhibitors improve insulin sensitivities by reducing the formation of angiotensin II and prevent the degradation of bradykinin, prolonging the half-life and cause relaxation of blood vessels through endothelial-dependent mechanism³².

Phaleria macrocarpa is native to Indonesia, efficacious as a medicine. Fruits of *Phaleria macrocarpa* empirically been used to treat various types of diseases such as cancer, liver disorders, heart disease, diabetes, arthritis, kidney disorders, stroke, and high blood pressure¹⁴. Qualitatively this plants showed the presence of alkaloids, saponins, polyphenols, and from fruit contained alkaloids, saponins and flavonoids¹⁵. *Phaleria macrocarpa* was reported to contain phenolic glycoside such as mahkotaside, mangiferin, kaempferol-3-O- β -dglucoside, dodecanoic acid, palmitic acid, ethyl stearate, and sucrose¹⁶. *Phaleria macrocarpa* (Scheff.) Boerl also contain lignat, that are pinoresinol, lariciresinol and matairesinol¹⁷. The results showed that *Phaleria macrocarpa* (Scheff.) Boerl has antidiabetic effects that inhibit the enzyme α -glukosidase and have anti-diabetic effect in mice induced sterptozotisin. The fruits of *Phaleria macrocarpa* has ACE inhibitory activity with IC₅₀ values was 122 g/ml in methanol extracts.

Elevations in Ang II and increased oxidative stress or impaired nitric oxide (NO) levels may also contribute to the pathogenesis of fructose fed rats. Some of evidence suggests that vascular oxidative stress is involved in the pathogenesis of many cardiovascular disorders in the insulin-resistant states, including diabetes, hypertension, and atherosclerosis. Fructose-fed rats had increased vascular production of superoxide anions that was accompanied by increased nicotinamide adenine dinucleotide phosphate (NADPH) oxidase activity³. Pericarp and mesocarp *P. macrocarpa* fruit show good antioxidant and anti-inflammatory activities. These activities might be due to the presence of phenolic and flavonoid compounds³³.

The results suggests that methanol extract of *Phalleria macrocarpha* could prevent the development high blood pressure and normalized blood pressure in rat induced by diet rich in fructose. The results tend to suggest that methanol extracts of possesses antihypertensive activity, through inhibition I of angiotensin converting enzyme AII.

1.4. ACKNOWLEDGEMENT

This Research was supported by funds from the Riset Unggulan Perguruan Tinggi- Direktorat Perguruan Tinggi.




Figure 1
Effect of methanol extract *P. marcocarpa* in mean change systolic blood pressure in fructose induced hypertensive in rats




Figure 2
Effect of methanol extract *P. marcocarpa* in mean change diastolic blood pressure in fructose induced hypertensive in rats

1.5. REFERENCES

1. Kher V. Renin inhibition--benefit beyond hypertension control, The J. of the Assoc. of Physic of India. 2009; 57: 518–521.
2. Reid IA. The Renin-Angiotensin System: Physiology, Pathophysiology, and Pharmacology. Advanc. in Phys. Edu. 1998; 20: 236–245.
3. Shinozaki K, Ayajiki K, Nishio Y, Sugaya T, Kashiwagi A., and Okamura T. Evidence for a causal role of the renin-angiotensin system in vascular dysfunction associated with insulin resistance. Hypertension. 2004; 6: 255–62.
4. Stamler J, Stamler N and Neaton JD. Blood Pressure, Systolic and Diastolic, and Cardiovascular Risks US Population Data. Arch Intern Med. 1993; 153: 598–615.
5. Weir MR. Dietary Fructose and Elevated Levels of Blood Pressure. J. Am Soc Nephrol. 2010; 21: 1416–1418.
6. Madero M, Perez-pozo SE, Jalal D, Johnson RJ and Sánchez-lozada LG. Dietary Fructose and Hypertension. Curr Hypertens Rep. 2001; 13: 29–35.
7. Martinez FJ, Rizza RA and Romero JC. High-fructose feeding elicits insulin resistance, hyperinsulinism, and hypertension in normal mongrel dogs. Hypertension 1994; 23: 456–463.
8. Abdulla MH, Sattar MA, Johns EJ. The Relation between Fructose-Induced Metabolic Syndrome and Altered Renal Haemodynamic and Excretory Function in the Rat. Internat J of Neph. 2011; 934659.
9. Bhanot S, and Mcneil JH. Insulin and hypertension: a causal relationship?. Cardiovas Research. 1996; 31: 95.
10. Hwang Is, Ho H, Hoffman BB, Reaven GM. Fructose induced insulin resistance and hypertension in rats. Hypertension. 1987; 10: 512-516.
11. Johnson RJ, Sanchez-lozada LG and Nakagawa T. The Effect of Fructose on Renal Biology and Disease. J Am Soc Nephrol. 2010; 21: 2036–2039.
12. Voors AA, Van Veldhuisen DJ, Van Gilst WH, et al. The current role of ACE inhibitors for secondary prevention in cardiovascular disease; from pathogenesis to clinical practice. Cardiovasc Drugs Ther. 2006; 20: 69–73.
13. Iwaniak A and Dziuba J. Animal and Plant Proteins as Precursors of Peptides with ACE Inhibitory Activity – An in silico Strategy of Protein Evaluation. Food Technology Biotechnology. 2009; 47(4): 441–449.
14. Harmanto, Research against Leaves and Fruit Crown god. One Day Seminar Mahkota Dewa. Center for Research and Development of Pharmaceutical and Traditional Medicine, Ministry of Health of the Republic of Indonesia. 2003.
15. Sumastuti R.. Research against Leaves and Fruit Crown god. One Day Seminar Mahkota Dewa. Center for Research and Development of Pharmaceutical and Traditional Medicine, Ministry of Health of the Republic of Indonesia. 2003.
16. Oshimi S, Zaima K, Matsuno Y, Hirasawa Y, Iizuka T, Studiawan H, Morita H. Studies on the constituents from the fruits of *Phaleria macrocarpa*. J Nat Med. 2008; 62(2): 207–210.
17. Saufi A, Heimendahl CBV, Alfermann AW, Fuss. Stereochemistry of lignans in *Phaleria macrocarpa* (Scheff.) Boerl. Naturforsch. 2008; 63(1-2): 13–16.
18. Rinayanti A, Radji M, Mun'im A., and Suyatna FD. Screening Angiotensin Converting Enzyme (ACE) Inhibitor activity of antihypertensive medicinal plants from Indonesia. Internat. J Pharm Teach & Pract. 2011; 3; 4(1): 527–32.
19. Tran LT, Yuen GP, Mcneill HJ. The fructose-fed rat: a review on the mechanisms of fructose-induced insulin resistance and hypertension. Mol Cell Biochem. 2009; 332: 145–159.
20. Vasdev S, Prabhakaran VM, Whelan M, Ford CAL, and Parai S. Fructose-induced hypertension, hypertriglyceridemia and elevated cytosolic calcium in rats: prevention by Deuterium oxide. Artery. 1994; 21(3): 124–147.
21. Verma S, Bhanot S, and McNeill J. (1995). Effect of chronic endothelin blockade in hyperinsulinemic hypertensive rats. American Journal Physiology. 1995; 269: H2017–H2021.
22. Erlich Y and Rosenthal T. Effect of angiotensin- enzyme inhibitors on fructose induced hypertension and hyperinsulinaemia in rats. Clinical Experimental Pharmacology Physiology. 1995; suppl 22: S347–S349.
23. Galipeau D, Arikawa E, Sekirov I, and McNeil JJ.. Chronic thromboxane synthase inhibition prevents fructose-induced hypertension. Hypertension. 2001; 38: 872.

24. Kurtz TW and Pravenec M. Antidiabetic mechanisms of angiotensin-converting enzyme inhibitors and angiotensin II receptor antagonists: beyond the renin-angiotensin system. *J. Hypertens.* 2004; 22(12): 2253–2261.
25. Corry DB and Tuck M. Obesity, hypertension, and sympathetic nervous system activity. *Current Hypertens. Reports.* 1999; 1: 119–126.
26. Singh AK, Amlal H, Haas PJ, Dringenberg Fussell S, Barone S L, Soleimani M. Fructose-induced hypertension: essential role of chloride and fructose absorbing transporters PAT1 and Glut5. *Kidney International* 2008; 74(4): 438–447.
27. Sijani P. Konsumsi Fruktosa Berlebihan dapat Berdampak Buruk bagi Kesehatan nusia. *JKM.* 2011; 10 (2): 173–189.
28. Stanton A. Therapeutic potential of renin inhibitors in the management of cardiovascular disorders. *Am J cardiovasc Drug.* 2003; 3(6): 389–394.
29. Leong X, Mustafa MR, Das S and Jaarin K. Association of elevated blood pressure and impaired vasorelaxation in experimental Sprague-Dawley rats fed with heated vegetable oil. *Lipids in Health and Disease.* 2010; 9(6): 1–10.
30. Kang DG, Moon MK, Sohn EJ, Lee D H and Lee HS. Effects of morin on blood pressure and metabolic changes in fructose-induced hypertensive rats. *Biological and Pharmaceutical Bulletin.* 2004; 27(1): 1779–83.
31. n LT, KM Macleod and JH.McNeill. Endothelin-1 modulates angiotensin II in the development of hypertension in fructose-fed s. *Moll Cell Biochem.* 2009; 325: 89–97.
32. Wiemer G, Scholkens BA, Becker RH and Busse R. Ramiprilat enhances endothelial autacoid formation by inhibiting breakdown of endothelium-derived bradykinin. *Hypertension.* 1991; 18(4): 558–563.
33. Hendra R, Ahmad S, Oskoueian E, Sukari A and Shukor MY. Antioxidant, anti-inflammatory and cytotoxicity of *Phaleria macrocarpa* (Boerl.) Scheff Fruit. *BMC Complementary and Alternative Medicine.* 2011; 11(1): 110.

Effects of the Methanol extracts Phaleria macrocarpa (Scheff) Boerl fruits on Angiotensin Converting Enzyme (ACE) activity

ORIGINALITY REPORT

14%

SIMILARITY INDEX

9%

INTERNET SOURCES

12%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

- 1

Ji Young Park, Xia Han, Mei Jing Piao, Min Chang Oh et al. "Hyperoside Induces Endogenous Antioxidant System to Alleviate Oxidative Stress", Journal of Cancer Prevention, 2016
Publication

1%
- 2

H. R. Black. "Metabolic and Clinical Outcomes in Nondiabetic Individuals With the Metabolic Syndrome Assigned to Chlorthalidone, Amlodipine, or Lisinopril as Initial Treatment for Hypertension: A report from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT)", Diabetes Care, 11/16/2007
Publication

1%
- 3

www.ftb.com.hr
Internet Source

1%
- 4

Delbosc, S.. "Involvement of oxidative stress and NADPH oxidase activation in the development of cardiovascular complications in

1%

a model of insulin resistance, the fructose-fed rat", Atherosclerosis, 200503

Publication

5

johtn.com

Internet Source

1 %

6

jamanetwork.com

Internet Source

1 %

7

T. F. LÜSCHER. "The endothelium as a target and mediator of cardiovascular disease", European Journal of Clinical Investigation, 11/1993

Publication

1 %

8

Roberto Ferrari. "ACE-Inhibition for Secondary Prevention of Cardiovascular Events—Should We Change Our Recommendation After PEACE?", Cardiovascular Drugs and Therapy, 2006

Publication

1 %

9

www.jabonline.in

Internet Source

1 %

10

Justo Pedroche. "Utilisation of chickpea protein isolates for production of peptides with angiotensin I-converting enzyme (ACE)-inhibitory activity", Journal of the Science of Food and Agriculture, 07/2002

Publication

1 %

11

Zorica Zakula, Goran Koricanac, Snezana Tepavcevic, Mojca Stojiljkovic, Tijana Milosavljevic, Esma R. Isenovic. "Impairment of cardiac insulin signaling in fructose-fed ovariectomized female Wistar rats", European Journal of Nutrition, 2011

Publication

<1 %

12

P. R. Nagareddy. "GPCR agonist-induced transactivation of the EGFR upregulates MLC II expression and promotes hypertension in insulin-resistant rats", Cardiovascular Research, 07/01/2010

Publication

<1 %

13

Maria G. Pavlatou, George Mastorakos, Ioannis Lekakis, Stavros Liatis et al. "Chronic administration of an angiotensin II receptor antagonist resets the hypothalamic–pituitary–adrenal (HPA) axis and improves the affect of patients with diabetes mellitus type 2: Preliminary results", Stress, 2009

Publication

<1 %

14

ahro.austin.org.au

Internet Source

<1 %

15

Liu, Qing, Jinjin Liu, Haili Guo, Shengnan Sun, Shifeng Wang, Yanling Zhang, Shiyu Li, and Yanjiang Qiao. "[6]-Gingerol: A Novel AT1 Antagonist for the Treatment of Cardiovascular

<1 %

Disease", Planta Medica, 2013.

Publication

16

Keiko WADA. "A Positive Association between Leptin and Blood Pressure of Normal Range in Japanese Men", Hypertension Research, 2006

Publication

<1 %

17

www.visitechsystems.com

Internet Source

<1 %

18

hotbetterpop.com

Internet Source

<1 %

19

qjmed.oxfordjournals.org

Internet Source

<1 %

20

Teruo Shiba, Masahiro Inoue, Hisaya Tada, Yoichi Hayashi et al. "Delapril versus manidipine in hypertensive therapy to halt the type-2-diabetes-mellitus-associated nephropathy", Diabetes Research and Clinical Practice, 2000

Publication

<1 %

21

jama.ama-assn.org

Internet Source

<1 %

22

eprints.undip.ac.id

Internet Source

<1 %

23

Mattace Raso, Giuseppina, Claudio Pirozzi, Roberta d'Emmanuele di Villa Bianca, Raffaele Simeoli, Anna Santoro, Adriano Lama,

<1 %

Francesca Di Guida, Roberto Russo, Carmen De Caro, Raffaella Sorrentino, Antonio Calignano, and Rosaria Meli.

"Palmitoylethanolamide Treatment Reduces Blood Pressure in Spontaneously Hypertensive Rats: Involvement of Cytochrome P450-Derived Eicosanoids and Renin Angiotensin System", PLoS ONE, 2015.

Publication

24

www.jraas.com

Internet Source

<1 %

25

Submitted to Chung-Ang University

Student Paper

<1 %

26

Flávio Curbani, Fernanda de Oliveira Busato, Maynara Marcarini do Nascimento, David Nicholas Olivieri, Carlos Eduardo Tadokoro.

"Inhale, exhale: Why particulate matter exposure in animal models are so acute? Data and facts behind the history", Data in Brief, 2019

Publication

<1 %

27

Jonathan M. Whittamore, Marguerite Hatch.

"Chronic metabolic acidosis reduces urinary oxalate excretion and promotes intestinal oxalate secretion in the rat", Urolithiasis, 2015

Publication

<1 %

28

intjem.biomedcentral.com

Internet Source

<1 %

29

Gunawan, Alexander A S, Wisnu Jatmiko, Vektor Dewanto, F. Rachmadi, and F. Jovan. "Tracking efficiency measurement of dynamic models on geometric particle filter using KLD-resampling", 2014 International Conference on Advanced Computer Science and Information System, 2014.

Publication

<1 %

30

docplayer.net

Internet Source

<1 %

31

sundoc.bibliothek.uni-halle.de

Internet Source

<1 %

32

A. T. A. Nandhini, C. V. Anuradha. "Hoe 140 abolishes the blood pressure lowering effect of taurine in high fructose-fed rats", Amino Acids, 2003

Publication

<1 %

33

Submitted to iGroup

Student Paper

<1 %

34

www.ijrte.org

Internet Source

<1 %

35

Submitted to Udayana University

Student Paper

<1 %



Exclude quotes

On

Exclude matches

< 10 words

Exclude bibliography

Off