

## Lampiran 12

Nilai-nilai kritis bagi uji satu-contoh Kolmogorov-Smirnov

uji satu-arah, $\alpha =$	.10	.05	.025	.01	.005
uji satu-arah, $\alpha =$	.20	.10	.05	.02	.01
n = 1	.900	.950	.975	.990	.995
2	.684	.776	.842	.900	.929
3	.565	.636	.708	.785	.829
4	.493	.565	.624	.689	.734
5	.447	.509	.563	.627	.669
6	.410	.468	.519	.577	.617
7	.381	.436	.483	.538	.576
8	.358	.410	.454	.507	.542
9	.339	.387	.430	.480	.513
10	.323	.369	.409	.457	.489
11	.308	.352	.391	.437	.468
12	.296	.338	.375	.419	.449
13	.285	.325	.361	.404	.432
14	.275	.314	.349	.390	.418
15	.266	.304	.338	.377	.404
16	.258	.295	.327	.366	.392
17	.250	.286	.318	.355	.381
18	.244	.279	.309	.346	.371
19	.237	.271	.301	.337	.361
20	.232	.265	.294	.329	.352
21	.226	.259	.287	.321	.344
22	.221	.253	.281	.314	.337
23	.216	.247	.275	.307	.330
24	.212	.242	.269	.301	.323
25	.208	.238	.264	.295	.317
26	.204	.233	.259	.290	.311
27	.200	.229	.254	.284	.305
28	.197	.225	.250	.279	.300
29	.193	.221	.246	.275	.295
30	.190	.218	.242	.270	.285
31	.187	.214	.238	.266	.281
32	.184	.211	.234	.262	.277
33	.182	.208	.231	.258	.273
34	.179	.205	.227	.254	.269
35	.177	.202	.224	.251	.265
36	.174	.199	.221	.247	.262
37	.172	.196	.218	.244	.258
38	.170	.194	.215	.241	.255
39	.168	.191	.213	.238	.252
40	.165	.189	.210	.235	.249
Hampiran bagi n > 40 :	$\frac{1.0730}{\sqrt{n}}$	$\frac{1.2239}{\sqrt{n}}$	$\frac{1.3581}{\sqrt{n}}$	$\frac{1.5174}{\sqrt{n}}$	$\frac{1.6276}{\sqrt{n}}$