REFERENCES

Aagaard P, Simonsen EB, Trolle M, Bangsbo J and Klausen K. Moment and power generation during maximal knee extensions performed at low and high speeds. Eur J Appl Physiol Occup Physiol 69: 376-381, 1994.

Aagaard P and Andersen JL. Correlation between contractile strength and myosin heavy chain isoform composition in human skeletal muscle. Med Sci Sports Exerc 30: 1217-1222, 1998.

Aagaard P, Simonsen EB, Andersen JL, Magnusson P and Dyhre-Poulsen P. Increased rate of force development and neural drive of human skeletal muscle following resistance training. J Appl Physiol 93: 1318-1326, 2002.

Aagaard P. Making muscles "stronger": exercise, nutrition, drugs. J Musculoskelet Neuronal Interact 4: 165-174, 2004.

Adams GR, Hather BM, Baldwin KM and Dudley GA. Skeletal muscle myosin heavy chain composition and resistance training. J Appl Physiol 74: 911-915, 1993.

Adams GR, Hather BM and Dudley GA. Effect of short-term unweighting on human skeletal muscle strength and size. Aviat Space Environ Med 65: 1116-1121, 1994.

Adams GR, Caiozzo VJ and Baldwin KM. Skeletal muscle unweighting: spaceflight and ground-based models. J Appl Physiol 95: 2185-2201, 2003.

Agureev AN and Kalandarov S. [Preserved food nutrition in the confinement study "Simulation of flight of International Crew on Space Station"]. [Experiment with long-term confinement: Problems and achievements], Moscow, 2001.

Akima H, Kawakami Y, Kubo K, Sekiguchi C, Ohshima H, Miyamoto A and Fukunaga T. Effect of short-duration spaceflight on thigh and leg muscle volume. Med Sci Sports Exerc 32: 1743-1747, 2000a.

Akima H, Kubo K, Kanehisa H, Suzuki Y, Gunji A and Fukunaga T. Leg-press resistance training during 20 days of 6 degrees head-down- tilt bed rest prevents muscle deconditioning. Eur J Appl Physiol 82: 30-38, 2000b.

Akima H, Kubo K, Imai M, Kanehisa H, Suzuki Y, Gunji A and Fukunaga T. Inactivity and muscle: effect of resistance training during bed rest on muscle size in the lower limb. Acta Physiol Scand 172: 269-278, 2001.

Akima H, Ushiyama JI, Kubo J, Tonosaki SI, Itoh M, Kawakami Y, Fukuoka H, Kanehisa H and Fukunaga T. Resistance Training during Unweighting Maintains Muscle Size and Function in Human Calf. Med Sci Sports Exerc 35: 655-662, 2003.

Alkner BA, Tesch PA and Berg HE. Quadriceps EMG/force relationship in knee extension and leg press. Med Sci Sports Exerc 32: 459-463, 2000.

Andersen JL, Klitgaard H and Saltin B. Myosin heavy chain isoforms in single fibres from m. vastus lateralis of sprinters: influence of training. Acta Physiol Scand 151: 135-142, 1994.

Andersen JL, Gruschy-Knudsen T, Sandri C, Larsson L and Schiaffino S. Bed rest increases the amount of mismatched fibers in human skeletal muscle. J Appl Physiol 86: 455-460, 1999.

Andersen JL and Aagaard P. Myosin heavy chain IIX overshoot in human skeletal muscle. Muscle Nerve 23: 1095-1104, 2000.

Andersen LL, Andersen JL, Magnusson SP, Suetta C, Madsen JL, Christensen LR and Aagaard P. Changes in the human muscle force-velocity relationship in response to resistance training and subsequent detraining. J Appl Physiol 99: 87-94, 2005.

Antonutto G, Capelli C, Girardis M, Zamparo P and di Prampero PE. Effects of microgravity on maximal power of lower limbs during very short efforts in humans. J Appl Physiol 86: 85-92, 1999.

Asher RAJ. The dangers of going to bed. Br Med J 4: 967-968, 1947.

Askling C, Karlsson J and Thorstensson A. Hamstring injury occurrence in elite soccer players after preseason strength training with eccentric overload. Scand J Med Sci Sports 13: 244-250, 2003.

Augustsson J, Esko A, Thomee R and Svantesson U. Weight training of the thigh muscles using closed vs. open kinetic chain exercises: a comparison of performance enhancement. J Orthop Sports Phys Ther 27: 3-8, 1998.

Baldwin KM, White TP, Arnaud SB, Edgerton VR, Kraemer WJ, Kram R, Raab-Cullen D and Snow CM. NASA roundtable: Musculoskeletal adaptations to weightlessness and development of effective countermeasures. Med Sci Sports Exerc 28: 1247-1253, 1996.

Bamman MM, Hunter GR, Stevens BR, Guilliams ME and Greenisen MC. Resistance exercise prevents plantar flexor deconditioning during bed rest. Med Sci Sports Exerc 29: 1462-1468, 1997.

Bamman MM, Clarke MS, Feeback DL, Talmadge RJ, Stevens BR, Lieberman SA and Greenisen MC. Impact of resistance exercise during bed rest on skeletal muscle sarcopenia and myosin isoform distribution. J Appl Physiol 84: 157-163, 1998.

Bamman MM and Caruso JF. Resistance exercise countermeasures for space flight: implications of training specificity. J Strength Cond Res 14: 45-49, 2000.

Bassey EJ and Short AH. A new method for measuring power output in a single leg extension: feasibility, reliability and validity. Eur J Appl Physiol Occup Physiol 60: 385-390, 1990.

Berg HE, Dudley GA, Häggmark T, Ohlsén H and Tesch PA. Effects of lower limb unloading on skeletal muscle mass and function in humans. J Appl Physiol 70: 1882-1885, 1991.

Berg HE, Dudley GA, Hather B and Tesch PA. Work capacity and metabolic and morphologic characteristics of the human quadriceps muscle in response to unloading. Clin Physiol 13: 337-347, 1993a.

Berg HE, Tedner B and Tesch PA. Changes in lower limb muscle cross-sectional area and tissue fluid volume after transition from standing to supine. Acta Physiol Scand 148: 379-385, 1993b.

Berg HE and Tesch PA. A gravity-independent ergometer to be used for resistance training in space. Aviat Space Environ Med 65: 752-756, 1994.

Berg HE and Tesch PA. Changes in muscle function in response to 10 days of lower limb unloading in humans. Acta Physiol Scand 157: 63-70, 1996.

Berg HE, Larsson L and Tesch PA. Lower limb skeletal muscle function after 6 wk of bed rest. J Appl Physiol 82: 182-188, 1997.

Berg HE and Tesch PA. Force and power characteristics of a resistive exercise device for use in space. Acta Astronaut 42: 219-230, 1998.

Bergström J. Muscle electrolytes in man. Scandinavian Journal of Clinical Laboratory Investigation 14: 7-110, 1962.

Berry P, Berry I and Manelfe C. Magnetic resonance imaging evaluation of lower limb muscles during bed rest--a microgravity simulation model. Aviat Space Environ Med 64: 212-218, 1993.

Boda WL, Watenpaugh DE, Ballard RE and Hargens AR. Supine lower body negative pressure exercise simulates metabolic and kinetic features of upright exercise. J Appl Physiol 89: 649-654, 2000.

Bonting SL, Ed. European isolation and confinement study. Advances in space biology and medicine, 1993.

Bosco C, Cardinale M and Tsarpela O. Influence of vibration on mechanical power and electromyogram activity in human arm flexor muscles. Eur J Appl Physiol Occup Physiol 79: 306-311, 1999.

Bottinelli R, Canepari M, Pellegrino MA and Reggiani C. Force-velocity properties of human skeletal muscle fibres: myosin heavy chain isoform and temperature dependence. J Physiol 495 (Pt 2): 573-586, 1996.

Bottinelli R. Functional heterogeneity of mammalian single muscle fibres: do myosin isoforms tell the whole story? Pflugers Arch 443: 6-17, 2001.

Caiozzo VJ, Perrine JJ and Edgerton VR. Training-induced alterations of the in vivo force-velocity relationship of human muscle. J Appl Physiol 51: 750-754, 1981.

Caruso J, Hamill J, Yamauchi M, Mercado D, Cook T, Higginson B, O'Meara S, Elias J and Siconolfi S. Albuterol Aids Resistance Exercise in Reducing Unloading-Induced Ankle Extensor Strength Losses. J Appl Physiol 98: 1705-1711, 2005.

Caruso JF, Hamill JL, Yamauchi M, Mercado DR, Cook TD, Keller CP, Montgomery AG and Elias J. Albuterol helps resistance exercise attenuate unloading-induced knee extensor losses. Aviat Space Environ Med 75: 505-511, 2004.

Chopard A, Arrighi N, Carnino A and Marini JF. Changes in dysferlin, proteins from dystrophin glycoprotein complex, costameres, and cytoskeleton in human soleus and vastus lateralis muscles after a long-term bedrest with or without exercise. Faseb J, 2005.

Clément G and Pavy-Le Traon A. Centrifugation as a countermeasure during actual and simulated microgravity: a review. Eur J Appl Physiol 92: 235-248, 2004.

Cobb S and Forbes A. Electromyographic studies of muscular fatigue in man. Am J Physiol 65: 234-251, 1923.

Conley MS, Foley JM, Ploutz-Snyder LL, Meyer RA and Dudley GA. Effect of acute head-down tilt on skeletal muscle cross-sectional area and proton transverse relaxation time. J Appl Physiol 81: 1572-1577, 1996.

Convertino VA, Doerr DF, Mathes KL, Stein SL and Buchanan P. Changes in volume, muscle compartment, and compliance of the lower extremities in man following 30 days of exposure to simulated microgravity. Aviat Space Environ Med 60: 653-658, 1989.

Convertino VA. Planning strategies for development of effective exercise and nutrition countermeasures for long-duration space flight. Nutrition 18: 880-888, 2002.

De Lorme TL, Ferris B, G. and Gallagher JR. Effect of progressive resistance exercise on muscle contraction time. Arch Phys Med Rehabil 33: 86-92, 1952.

Deitrick JE, Whedon GD and Shorr E. Effects of immobilization upon various metabolic and physiologic functions of normal men. Am J Med 4: 3-35, 1948.

Deschenes MR, Giles JA, McCoy RW, Volek JS, Gomez AL and Kraemer WJ. Neural factors account for strength decrements observed after short- term muscle unloading. Am J Physiol Regul Integr Comp Physiol 282: R578-583, 2002.

Desmedt JE and Godaux E. Ballistic contractions in man: characteristic recruitment pattern of single motor units of the tibialis anterior muscle. J Physiol 264: 673-693, 1977.

Desmedt JE and Godaux E. Voluntary motor commands in human ballistic movements. Ann Neurol 5: 415-421, 1979.

di Prampero PE. The Twin Bikes System for artificial gravity in space. J Gravit Physiol 1: P12-14, 1994.

Duchateau J. Bed rest induces neural and contractile adaptations in triceps surae. Med Sci Sports Exerc 27: 1581-1589, 1995.

Dudley GA, Duvoisin MR, Convertino VA and Buchanan P. Alterations of the in vivo torque-velocity relationship of human skeletal muscle following 30 days exposure to simulated microgravity. Aviat Space Environ Med 60: 659-663, 1989.

Dudley GA, Tesch PA, Miller BJ and Buchanan P. Importance of eccentric actions in performance adaptations to resistance training. Aviat Space Environ Med 62: 543-550, 1991.

Dudley GA, Duvoisin MR, Adams GR, Meyer RA, Belew AH and Buchanan P. Adaptations to unilateral lower limb suspension in humans. Aviat Space Environ Med 63: 678-683, 1992.

Dupui P, Montoya R, Costes-Salon MC, Severac A and Guell A. Balance and gait analysis after 30 days -6 degrees bed rest: influence of lower-body negative-pressure sessions. Aviation Space & Environmental Medicine 63: 1004-1010, 1992.

Duvoisin MR, Convertino VA, Buchanan P, Gollnick PD and Dudley GA. Characteristics and preliminary observations of the influence of electromyostimulation on the size and function of human skeletal muscle during 30 days of simulated microgravity. Aviat Space Environ Med 60: 671-678, 1989.

Edgerton VR and Roy RR. Neuromuscular adaptions to actual and simulated spaceflight. Handbook of Physiology: Adaption to the environment, Section 4: Part 3: The gravitational environment. Ed. Fregly MJ and Blatteis CM. New York, Oxford University Press: 721-763, 1995.

Edgerton VR, Zhou MY, Ohira Y, Klitgaard H, Jiang B, Bell G, Harris B, Saltin B, Gollnick PD, Roy RR and et al. Human fiber size and enzymatic properties after 5 and 11 days of spaceflight. J Appl Physiol 78: 1733-1739, 1995.

Edström L. Selective atrophy of red muscle fibres in the quadriceps in long-standing knee-joint dysfunction. Injuries to the anterior cruciate ligament. J Neurol Sci 11: 551-558, 1970.

Elmann-Larsen B and Schmitt D. Staying in bed to benefit ESA's astronauts and Europe's citicens. ESA Bull 113: 34-39, 2003.

Enoka RM. Neural adaptations with chronic physical activity. J Biomech 30: 447-455, 1997.

Ericson MO, Nisell R and Ekholm J. Quantified electromyography of lower-limb muscles during level walking. Scand J Rehabil Med 18: 159-163, 1986.

ESA-CNES-NASDA. Final report. Simulation of a mission aboard the international space station by a long-term (90 days) anti-orthostatic bed rest at -6° on healthy subjects, 2003a.

ESA-CNES-NASDA. Clinical report. Simulation of a mission aboard the international space station by a long-term (90 days) anti-orthostatic bed rest at -6° on healthy subjects, 2003b.

Farina D, Cescon C and Merletti R. Influence of anatomical, physical, and detection-system parameters on surface EMG. Biol Cybern 86: 445-456, 2002.

Ferrando AA, Stuart CA, Brunder DG and Hillman GR. Magnetic resonance imaging quantitation of changes in muscle volume during 7 days of strict bed rest. Aviat Space Environ Med 66: 976-981, 1995.

Ferrando AA, Lane HW, Stuart CA, Davis-Street J and Wolfe RR. Prolonged bed rest decreases skeletal muscle and whole body protein synthesis. Am J Physiol 270: E627-633, 1996.

Ferretti G, Antonutto G, Denis C, Hoppeler H, Minetti AE, Narici MV and Desplanches D. The interplay of central and peripheral factors in limiting maximal O2 consumption in man after prolonged bed rest. J Physiol 501 (Pt 3): 677-686, 1997.

Ferri A, Scaglioni G, Pousson M, Capodaglio P, Van Hoecke J and Narici MV. Strength and power changes of the human plantar flexors and knee extensors in response to resistance training in old age. Acta Physiol Scand 177: 69-78, 2003.

Fitts RH, Riley DR and Widrick JJ. Physiology of a microgravity environment invited review: microgravity and skeletal muscle. J Appl Physiol 89: 823-839, 2000.

Fitts RH, Riley DR and Widrick JJ. Functional and structural adaptations of skeletal muscle to microgravity. J Exp Biol 204: 3201-3208, 2001.

Fluckey JD, Dupont-Versteegden EE, Montague DC, Knox M, Tesch P, Peterson CA and Gaddy-Kurten D. A rat resistance exercise regimen attenuates losses of musculoskeletal mass during hindlimb suspension. Acta Physiol Scand 176: 293-300, 2002.

Fortney SM, Schneider VS and Greenleaf JE. The physiology of bed rest. Handbook of Physiology: Adaption to the environment, Section 4: Part 3: The gravitational environment. Ed. Feregley MJ and Blatteis CM, Oxford Univ. Press. 2: 889-939, 1996.

Funato K, Matsuo A, Yata H, Akima H, Suzuki Y, Gunji A and Fukunaga T. Changes in forcevelocity and power output of upper and lower extremity musculature in young subjects following 20 days bed rest. J Gravit Physiol 4: S22-30, 1997.

Germain P, Guell A and Marini JF. Muscle strength during bedrest with and without muscle exercise as a countermeasure. Eur J Appl Physiol Occup Physiol 71: 342-348, 1995.

Gogia P, Schneider VS, LeBlanc AD, Krebs J, Kasson C and Pientok C. Bed rest effect on extremity muscle torque in healthy men. Arch Phys Med Rehabil 69: 1030-1032, 1988.

Gollnick PD, Karlsson J, Piehl K and Saltin B. Selective glycogen depletion in skeletal muscle fibres of man following sustained contractions. J Physiol 241: 59-67, 1974a.

Gollnick PD, Sjödin B, Karlsson J, Jansson E and Saltin B. Human soleus muscle: a comparison of fiber composition and enzyme activities with other leg muscles. Pflugers Arch 348: 247-255, 1974b.

Greenleaf JE, Bulbulian R, Bernauer EM, Haskell WL and Moore T. Exercise-training protocols for astronauts in microgravity. J Appl Physiol 67: 2191-2204, 1989.

Greenleaf JE. Intensive exercise training during bed rest attenuates deconditioning. Med Sci Sports Exerc 29: 207-215, 1997.

Greenleaf JE, Rehrer NJ, Mohler SR, Quach DT and Evans DG. Airline chair-rest deconditioning: induction of immobilisation thromboemboli? Sports Med 34: 705-725, 2004.

Grimby L and Hannerz J. Firing rate and recruitment order of toe extensor motor units in different modes of voluntary conraction. J Physiol 264: 865-879, 1977.

Gür H, Gransberg L, vanDyke D, Knutsson E and Larsson L. Relationship between in vivo muscle force at different speeds of isokinetic movements and myosin isoform expression in men and women. Eur J Appl Physiol 88: 487-496, 2003.

Hargens AR, Parazynski S, Aratow M and Fridén J. Muscle changes with eccentric exercise: implications on Earth and in space. Adv Myochem 2: 299-312, 1989.

Harridge SD, Bottinelli R, Canepari M, Pellegrino MA, Reggiani C, Esbjornsson M and Saltin B. Whole-muscle and single-fibre contractile properties and myosin heavy chain isoforms in humans. Pflugers Arch 432: 913-920, 1996.

Hather BM, Tesch PA, Buchanan P and Dudley GA. Influence of eccentric actions on skeletal muscle adaptations to resistance training. Acta Physiol Scand 143: 177-185, 1991.

Hather BM, Adams GR, Tesch PA and Dudley GA. Skeletal muscle responses to lower limb suspension in humans. J Appl Physiol 72: 1493-1498, 1992.

Hikida RS, Gollnick PD, Dudley GA, Convertino VA and Buchanan P. Structural and metabolic characteristics of human skeletal muscle following 30 days of simulated microgravity. Aviat Space Environ Med 60: 664-670, 1989.

Holmbäck AM, Porter MM, Downham D and Lexell J. Reliability of isokinetic ankle dorsiflexor strength measurements in healthy young men and women. Scand J Rehabil Med 31: 229-239, 1999.

Hortobágyi T, Lambert NJ and Hill JP. Greater cross education following training with muscle lengthening than shortening. Med Sci Sports Exerc 29: 107-112, 1997.

Hwang TI, Hill K, Schneider V and Pak CY. Effect of prolonged bedrest on the propensity for renal stone formation. J Clin Endocrinol Metab 66: 109-112, 1988.

Häggmark T and Eriksson E. Hypotrophy of the soleus muscle in man after achilles tendon rupture. Discussion of findings obtained by computed tomography and morphologic studies. Am J Sports Med 7: 121-126, 1979.

Häggmark T, Jansson E and Eriksson E. Fiber type area and metabolic potential of the thigh muscle in man after knee surgery and immobilization. Int J Sports Med 2: 12-17, 1981.

Häggmark T, Eriksson E and Jansson E. Muscle fiber type changes in human skeletal muscle after injuries and immobilization. Orthopedics 9: 181-185, 1986.

Häkkinen K and Komi PV. Training-induced changes in neuromuscular performance under voluntary and reflex conditions. Eur J Appl Physiol Occup Physiol 55: 147-155, 1986.

Häkkinen K. Neuromuscular and hormonal adaptations during strength and power training. A review. J Sports Med Phys Fitness 29: 9-26, 1989.

Häkkinen K, Kallinen M, Linnamo V, Pastinen UM, Newton RU and Kraemer WJ. Neuromuscular adaptations during bilateral versus unilateral strength training in middle-aged and elderly men and women. Acta Physiol Scand 158: 77-88, 1996.

Johnson MA, Polgar J, Weightman D and Appleton D. Data on the distribution of fibre types in thirtysix human muscles. An autopsy study. J Neurol Sci 18: 111-129, 1973.

Kanehisa H and Miyashita M. Specificity of velocity in strength training. Eur J Appl Physiol Occup Physiol 52: 104-106, 1983.

Komi PV. Stretch-shortening cycle. Strength and Power in Sports. Ed. Komi PV. Oxford, Blackwell Science: 184-202, 2003.

Koryak Y. Contractile properties of the human triceps surae muscle during simulated weightlessness. Eur J Appl Physiol Occup Physiol 70: 344-350, 1995a.

Koryak Y. Mechanical and electrical adaptation of skeletal muscle to gravitational unloading. J Gravit Physiol 2: P76-79, 1995b.

Koryak Y. Mechanical and electrical changes in human muscle after dry immersion. Eur J Appl Physiol Occup Physiol 74: 133-140, 1996.

Koryak Y. Effect of 120 days of bed-rest with and without countermeasures on the mechanical properties of the triceps surae muscle in young women. Eur J Appl Physiol Occup Physiol 78: 128-135, 1998a.

Koryak Y. Electromyographic study of the contractile and electrical properties of the human triceps surae muscle in a simulated microgravity environment. J Physiol 510: 287-295, 1998b.

Koryak Y. The effects of long-term simulated microgravity on neuromuscular performance in men and women. Eur J Appl Physiol Occup Physiol 79: 168-175, 1999a.

Koryak Y. Electrical and contractile parameters of muscle in man: effects of 7- day "dry" water immersion. Aviat Space Environ Med 70: 459-464, 1999b.

Koryak Y. "DRY" immersion induces neural and contractile adaptations in the human triceps surae muscle. Environ Med 46: 17-27, 2002.

Koryak YA. Influence of 120-days 6 degrees head-down tilt bed rest on the functional properties of the neuromuscular system in man. Aviat Space Environ Med 69: 766-770, 1998c.

Koryak YU. Electrically evoked and voluntary properties of the human triceps surae muscle: effects of long-term spaceflights. Acta Physiol Pharmacol Bulg 26: 21-27, 2001.

Kozlovskaya I. "Operational and physiological evaluation of the resistive exercise training using "Flywheel" technology, in the framework of the Russian "Simulation of Flight of International Crew on Space Station" Programme (SFINCSS-99)", State Research Center - Institute for Biomedical Problems, Moscow, Russia, 2000.

Kozlovskaya IB, Kreidich YV, Oganov VS and Koserenko OP. Pathophysiology of motor functions in prolonged manned space flights. Acta Astronaut 8: 1059-1072, 1981.

Kozlovskaya IB and Egorov AD. Some approaches to medical support for Martian expedition. Acta Astronaut 53: 269-275, 2003.

Kreitenberg A, Baldwin KM, Bagian JP, Cotten S, Witmer J and Caiozzo VJ. The "Space Cycle" Self Powered Human Centrifuge: a proposed countermeasure for prolonged human spaceflight. Aviat Space Environ Med 69: 66-72, 1998.

Kubo K, Akima H, Kouzaki M, Ito M, Kawakami Y, Kanehisa H and Fukunaga T. Changes in the elastic properties of tendon structures following 20 days bed-rest in humans. Eur J Appl Physiol 83: 463-468, 2000.

Kubo K, Akima H, Ushiyama J, Tabata I, Fukuoka H, Kanehisa H and Fukunaga T. Effects of 20 days of bed rest on the viscoelastic properties of tendon structures in lower limb muscles. Br J Sports Med 38: 324-330, 2004a.

Kubo K, Akima H, Ushiyama J, Tabata I, Fukuoka H, Kanehisa H and Fukunaga T. Effects of resistance training during bed rest on the viscoelastic properties of tendon structures in the lower limb. Scand J Med Sci Sports 14: 296-302, 2004b.

Lambertz D, Perot C, Kaspranski R and Goubel F. Effects of long-term spaceflight on mechanical properties of muscles in humans. J Appl Physiol 90: 179-188, 2001.

Lambertz D, Goubel F, Kaspranski R and Perot C. Influence of long-term spaceflight on neuromechanical properties of muscles in humans. J Appl Physiol 94: 490-498, 2003.

Larsson L and Moss RL. Maximum velocity of shortening in relation to myosin isoform composition in single fibres from human skeletal muscles. J Physiol 472: 595-614, 1993.

Larsson L, Li X, Berg HE and Frontera WR. Effects of removal of weight-bearing function on contractility and myosin isoform composition in single human skeletal muscle cells. Pflugers Arch 432: 320-328, 1996.

Layne CS, Mulavara AP, Pruett CJ, McDonald PV, Kozlovskaya IB and Bloomberg JJ. The use of in-flight foot pressure as a countermeasure to neuromuscular degradation. Acta Astronaut 42: 231-246, 1998.

LeBlanc A, Gogia P, Schneider V, Krebs J, Schonfeld E and Evans H. Calf muscle area and strength changes after five weeks of horizontal bed rest. Am J Sports Med 16: 624-629, 1988.

LeBlanc A, Rowe R, Schneider V, Evans H and Hedrick T. Regional muscle loss after short duration spaceflight. Aviat Space Environ Med 66: 1151-1154, 1995.

LeBlanc A, Rowe R, Evans H, West S, Shackelford L and Schneider V. Muscle atrophy during long duration bed rest. Int J Sports Med 18 Suppl 4: S283-285, 1997.

LeBlanc A, Lin C, Shackelford L, Sinitsyn V, Evans H, Belichenko O, Schenkman B, Kozlovskaya I, Oganov V, Bakulin A, Hedrick T and Feeback D. Muscle volume, MRI relaxation times (T2), and body composition after spaceflight. J Appl Physiol 89: 2158-2164, 2000a.

LeBlanc A, Schneider V, Shackelford L, West S, Oganov V, Bakulin A and Voronin L. Bone mineral and lean tissue loss after long duration space flight. J Musculoskel Neuron Interact 1: 157-160, 2000b.

LeBlanc AD, Schneider VS, Evans HJ, Pientok C, Rowe R and Spector E. Regional changes in muscle mass following 17 weeks of bed rest. J Appl Physiol 73: 2172-2178, 1992.

Lee SM, Bennett BS, Hargens AR, Watenpaugh DE, Ballard RE, Murthy G, Ford SR and Fortney SM. Upright exercise or supine lower body negative pressure exercise maintains exercise responses after bed rest. Med Sci Sports Exerc 29: 892-900, 1997.

Lee SM, Cobb K, Loehr JA, Nguyen D and Schneider SM. Foot-ground reaction force during resistive exercise in parabolic flight. Aviat Space Environ Med 75: 405-412, 2004.

Lindh M. Increase of muscle strength from isometric quadriceps exercises at different knee angles. Scand J Rehabil Med 11: 33-36, 1979.

Louisy F, Gaudin C, Oppert JM, Guell A and Guezennec CY. Haemodynamics of leg veins during a 30-days-6 degrees head-down bedrest with and without lower body negative pressure. European Journal of Applied Physiology & Occupational Physiology 61: 349-355, 1990.

MacDougall JD. Hypertrophy and hyperplasia. Strength and Power in Sports. Ed. Komi PV. Oxford, Blackwell Science: 252-264, 2003.

Maillet A and Pavy-Le Traon A. MEDES clinical research facility as a tool to prepare ISSA space flights. Acta Astronaut 43: 321-328, 1998.

Mayr W, Bijak M, Girsch W, Hofer C, Lanmuller H, Rafolt D, Rakos M, Sauermann S, Schmutterer C, Schnetz G, Unger E and Freilinger G. MYOSTIM-FES to prevent muscle atrophy in microgravity and bed rest: preliminary report. Artif Organs 23: 428-431, 1999.

Miller RG, Mirka A and Maxfield M. Rate of tension development in isometric contractions of a human hand muscle. Exp Neurol 73: 267-285, 1981.

Monod H. Contribution à l'étude du travail statique. Institut National de Securite pour la prevention des accidents du travail et des maladies professionelles. Paris, Institut National de Securite: 124, 1956.

Moritani T and deVries HA. Neural factors versus hypertrophy in the time course of muscle strength gain. Am J Phys Med 58: 115-130, 1979.

Morrissey M. Comparison of conventional multi-gym (MG) training and the fly-wheel ergometer (FWE) for strengthening the quadriceps muscle group in patients with unilateral knee injury. Proc. of Modern methods in the rehabilitation of sportsmen, Ljubljana, 2004.

Morrissey MC, Harman EA and Johnson MJ. Resistance training modes: specificity and effectiveness. Med Sci Sports Exerc 27: 648-660, 1995.

Nardone A and Schieppati M. Shift of activity from slow to fast muscle during voluntary lengthening contractions of the triceps surae muscles in humans. J Physiol 395: 363-381, 1988.

Nardone A, Romano C and Schieppati M. Selective recruitment of high-threshold human motor units during voluntary isotonic lengthening of active muscles. J Physiol 409: 451-471, 1989.

Narici MV, Roi GS, Landoni L, Minetti AE and Cerretelli P. Changes in force, cross-sectional area and neural activation during strength training and detraining of the human quadriceps. Eur J Appl Physiol Occup Physiol 59: 310-319, 1989.

Narici MV, De-Boer M, Goldstraw C and Seynnes O. Timecourse of Muscular and Tendinous Adaptations with Strength Training. Proc. of Isokinetic congress, Bologna, 2005.

Nicogossian AE, Sawin AI and Grigoriev AI. Countermeasures to space deconditioning. Space physiology and medicine. Ed. Nicogossian, Huntoon and Pool. Malvern, Lea & Febiger: 447-467, 1994a.

Nicogossian AE, Sawin AI and Leach Huntoon C. Overall physiologic response to space flight. Space physiology and medicine. Ed. Nicogossian, Huntoon and Pool. Malvern, Lea & Febiger: 213-227, 1994b.

Nordander C, Willner J, Hansson GA, Larsson B, Unge J, Granquist L and Skerfving S. Influence of the subcutaneous fat layer, as measured by ultrasound, skinfold calipers and BMI, on the EMG amplitude. Eur J Appl Physiol 89: 514-519, 2003.

Ohira Y, Jiang B, Roy RR, Oganov V, Ilyina-Kakueva E, Marini JF and Edgerton VR. Rat soleus muscle fiber responses to 14 days of spaceflight and hindlimb suspension. J Appl Physiol 73: 51S-57S, 1992.

Ohira Y, Yoshinaga T, Ohara M, Nonaka I, Yoshioka T, Yamashita-Goto K, Shenkman BS, Kozlovskaya IB, Roy RR and Edgerton VR. Myonuclear domain and myosin phenotype in human soleus after bed rest with or without loading. J Appl Physiol 87: 1776-1785, 1999.

Oishi Y, Ishihara A, Yamamoto H and Miyamoto E. Hindlimb suspension induces the expression of multiple myosin heavy chain isoforms in single fibres of the rat soleus muscle. Acta Physiol Scand 162: 127-134, 1998.

Paddon-Jones D, Sheffield-Moore M, Urban RJ, Sanford AP, Aarsland A, Wolfe RR and Ferrando AA. Essential amino acid and carbohydrate supplementation ameliorates muscle protein loss in humans during 28 days bedrest. J Clin Endocrinol Metab 89: 4351-4358, 2004.

Paddon-Jones D, Wolfe RR and Ferrando AA. Amino acid supplementation for reversing bed rest and steroid myopathies. J Nutr 135: 1809S-1812S, 2005.

Pavol MJ and Grabiner MD. Knee strength variability between individuals across ranges of motion and hip angles. Med Sci Sports Exerc 32: 985-992, 2000.

Petrofsky JS and Lind AR. The influence of temperature on the amplitude and frequency components of the EMG during brief and sustained isometric contractions. Eur J Appl Physiol Occup Physiol 44: 189-200, 1980.

Ploutz-Snyder LL, Tesch PA, Crittenden DJ and Dudley GA. Effect of unweighting on skeletal muscle use during exercise. J Appl Physiol 79: 168-175, 1995.

Portero P, Vanhoutte C and Goubel F. Surface electromyogram power spectrum changes in human leg muscles following 4 weeks of simulated microgravity. Eur J Appl Physiol Occup Physiol 73: 340-345, 1996.

Rasch PJ and Morehouse LE. Effect of static and dynamic exercises on muscular strength and hypertrophy. J Appl Physiol 11: 29-34, 1957.

Reeves ND, Narici MV and Maganaris CN. Strength training alters the viscoelastic properties of tendons in elderly humans. Muscle Nerve 28: 74-81, 2003.

Reeves ND, Maganaris CN, Ferretti G and Narici MV. Influence of 90-Day Simulated Microgravity on Human Tendon Mechanical Properties and the Effect of Resistive Countermeasures. J Appl Physiol 98: 2278-2286, 2005.

Rittweger J, Frost HM, Schiessl H, Alkner B, Tesch P and Felsenberg D. Muscle atrophy and bone loss after 90 days' bed rest and the effects of flywheel resistive exercise and pamidronate: Results from the LTBR study. Bone 36: 1019-1029, 2005.

Rohmert W. [Determination of the recovery pause for static work of man]. Int Z Angew Physiol 18: 123-164, 1960.

Rudnick J, Puttmann B, Tesch PA, Alkner B, Schoser BG, Salanova M, Kirsch K, Gunga HC, Schiffl G, Luck G and Blottner D. Differential expression of nitric oxide synthases (NOS 1-3) in human skeletal muscle following exercise countermeasure during 12 weeks of bed rest. Faseb J 18: 1228-1230, 2004.

Ruegg DG, Kakebeeke TH, Gabriel JP and Bennefeld M. Conduction velocity of nerve and muscle fiber action potentials after a space mission or a bed rest. Clin Neurophysiol 114: 86-93, 2003.

Rummel JA, Sawin CF, Michel EL, Buderer MC and Thornton WT. Exercise and long duration speceflight through 84 days. J Am Med Womens Assoc 30: 173-187, 1975.

Rutherford OM and Jones DA. The role of learning and coordination in strength training. Eur J Appl Physiol Occup Physiol 55: 100-105, 1986.

Rutherford OM. Muscular coordination and strength training. Implications for injury rehabilitation. Sports Med 5: 196-202, 1988.

Sale D and MacDougall D. Specificity in strength training: a review for the coach and athlete. Can J Appl Sport Sci 6: 87-92, 1981.

Sale D. Specificity of training. Can J Sport Sci 17: 71, 1992.

Sale DG, Martin JE and Moroz DE. Hypertrophy without increased isometric strength after weight training. Eur J Appl Physiol Occup Physiol 64: 51-55, 1992.

Sale DG. Neural adaptation to strength training. Strength and Power in Sports. Ed. Komi PV. Oxford, Blackwell Science: 281-314, 2003.

Schmitt D and Elmann-Larsen B. Bed-rest studies for the international space station. ESA Bull 108: 94-100, 2001.

Schneider VS, LeBlanc AD and Taggart LC. Physiological adaption to space flight; Bone and mineral metabolism. Space physiology and medicine. Ed. Nicogossian, Huntoon and Pool. Malvern, Lea & Febiger: 327-333, 1994.

Schulze K, Gallagher P and Trappe S. Resistance training preserves skeletal muscle function during unloading in humans. Med Sci Sports Exerc 34: 303-313, 2002.

Shackelford LC, LeBlanc AD, Driscoll TB, Evans HJ, Rianon NJ, Smith SM, Spector E, Feeback DL and Lai D. Resistance exercise as a countermeasure to disuse-induced bone loss. J Appl Physiol 97: 119-129, 2004.

Sleivert GG, Backus RD and Wenger HA. The influence of a strength-sprint training sequence on multi-joint power output. Med Sci Sports Exerc 27: 1655-1665, 1995.

Stein TP, Leskiw MJ, Schluter MD, Donaldson MR and Larina I. Protein kinetics during and after long-duration spaceflight on MIR. Am J Physiol 276: E1014-1021, 1999.

Suzuki Y, Murakami T, Haruna Y, Kawakubo K, Goto S, Makita Y, Ikawa S and Gunji A. Effects of 10 and 20 days bed rest on leg muscle mass and strength in young subjects. Acta Physiol Scand Suppl 616: 5-18, 1994.

Talmadge RJ. Myosin heavy chain isoform expression following reduced neuromuscular activity: potential regulatory mechanisms. Muscle Nerve 23: 661-679, 2000.

Taniguchi Y. Lateral specificity in resistance training: the effect of bilateral and unilateral training. Eur J Appl Physiol Occup Physiol 75: 144-150, 1997.

Taylor HL, Erickson L, Henschel A and Keys A. The effect of bed rest on the blood volume of normal young men. Am J Physiol 144: 227-232, 1945.

Tesch PA and Karlsson J. Muscle fiber types and size in trained and untrained muscles of elite athletes. J Appl Physiol 59: 1716-1720, 1985.

Tesch PA, Lindborg BP and Colliander EB. Evaluation of a dynamometer measuring torque of uniand bilateral concentric and eccentric muscle action. Clin Physiol 10: 1-9, 1990.

Tesch PA and Alkner BA. Acute and chronic muscle metabolic adaptations to strength training. Strength and Power in Sports. Ed. Komi PV. Oxford, Blackwell Science: 265-280, 2003.

Tesch PA, Trieschmann JT and Ekberg A. Plantar flexor muscle atrophy in men and women in response to five wk simulated spaceflight. 14th Humans in Space Symposium, Banff, Alberta, Canada, 2003.

Tesch PA, Ekberg A, Lindqvist DM and Trieschmann JT. Muscle hypertrophy following five wk resistance training using a non-gravity dependent loading principle. Acta Physiol Scand 180: 89-98, 2004a.

Tesch PA, Trieschmann JT and Ekberg A. Hypertrophy of chronically unloaded muscle subjected to resistance exercise. J Appl Physiol 96: 1451-1458, 2004b.

Tesch PA, Berg HE, Bring D, Evans H and LeBlanc A. Effects of 17-day spaceflight on knee extensor muscle function and size. Eur J Appl Physiol 93: 463-468, 2005.

Thomason DB and Booth FW. Atrophy of the soleus muscle by hindlimb unweighting. J Appl Physiol 68: 1-12, 1990.

Thorstensson A, Grimby G and Karlsson J. Force-velocity relations and fiber composition in human knee extensor muscles. J Appl Physiol 40: 12-16, 1976a.

Thorstensson A, Karlsson J, Viitasalo JH, Luhtanen P and Komi PV. Effect of strength training on EMG of human skeletal muscle. Acta Physiol Scand 98: 232-236, 1976b.

Trappe S, Williamson D, Godard M, Porter D, Rowden G and Costill D. Effect of resistance training on single muscle fiber contractile function in older men. J Appl Physiol 89: 143-152, 2000.

Trappe S, Godard M, Gallagher P, Carroll C, Rowden G and Porter D. Resistance training improves single muscle fiber contractile function in older women. Am J Physiol Cell Physiol 281: C398-406, 2001.

Trappe S, Trappe T, Gallagher P, Harber M, Alkner B and Tesch P. Human single muscle fibre function with 84 day bed-rest and resistance exercise. J Physiol 557: 501-513, 2004a.

Trappe TA, Raue U and Tesch PA. Human soleus muscle protein synthesis following resistance exercise. Acta Physiol Scand 182: 189-196, 2004b.

Vaernes RJ, Bergan T, Warncke M, Ursin H, Aakvaag A and Hockey R. European isolation and confinement study. Workload and stress: effects on psychosomatic and psychobiological reaction patterns. Adv Space Biol Med 3: 95-120, 1993.

Van Cutsem M, Duchateau J and Hainaut K. Changes in single motor unit behaviour contribute to the increase in contraction speed after dynamic training in humans. J Physiol 513: 295-305, 1998.

Weiss LW, Clark FC and Howard DG. Effects of heavy-resistance triceps surae muscle training on strength and muscularity of men and women. Phys Ther 68: 208-213, 1988.

Westing SH, Seger JY, Karlson E and Ekblom B. Eccentric and concentric torque-velocity characteristics of the quadriceps femoris in man. Eur J Appl Physiol Occup Physiol 58: 100-104, 1988.

Vico L, Collet P, Guignandon A, Lafage-Proust MH, Thomas T, Rehaillia M and Alexandre C. Effects of long-term microgravity exposure on cancellous and cortical weight-bearing bones of cosmonauts. Lancet 355: 1607-1611, 2000.

Widrick JJ, Trappe SW, Costill DL and Fitts RH. Force-velocity and force-power properties of single muscle fibers from elite master runners and sedentary men. Am J Physiol 271: C676-683, 1996.

Widrick JJ, Knuth ST, Norenberg KM, Romatowski JG, Bain JL, Riley DA, Karhanek M, Trappe SW, Trappe TA, Costill DL and Fitts RH. Effect of a 17 day spaceflight on contractile properties of human soleus muscle fibres. J Physiol 516: 915-930, 1999.

Williamson DL, Godard MP, Porter DA, Costill DL and Trappe SW. Progressive resistance training reduces myosin heavy chain coexpression in single muscle fibers from older men. J Appl Physiol 88: 627-633, 2000.

Williamson DL, Gallagher PM, Carroll CC, Raue U and Trappe SW. Reduction in hybrid single muscle fiber proportions with resistance training in humans. J Appl Physiol 91: 1955-1961, 2001.

Wilson G and Murphy A. The efficacy of isokinetic, isometric and vertical jump tests in exercise science. Aust J Sci Med Sport 27: 20-24, 1995.

Winter DA and Yack HJ. EMG profiles during normal human walking: stride-to-stride and intersubject variability. Electroencephalogr Clin Neurophysiol 67: 402-411, 1987.

Zange J, Muller K, Schuber M, Wackerhage H, Hoffmann U, Gunther RW, Adam G, Neuerburg JM, Sinitsyn VE, Bacharev AO and Belichenko OI. Changes in calf muscle performance, energy metabolism, and muscle volume caused by long-term stay on space station MIR. Int J Sports Med 18 Suppl 4: S308-309, 1997.

Zhou MY, Klitgaard H, Saltin B, Roy RR, Edgerton VR and Gollnick PD. Myosin heavy chain isoforms of human muscle after short-term spaceflight. J Appl Physiol 78: 1740-1744, 1995.