

Analisis Univariat

Umur

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	10	16	32.0	32.0	32.0
	11	32	64.0	64.0	96.0
	12	2	4.0	4.0	100.0
	Total	50	100.0	100.0	

Jenis Kelamin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Laki-laki	22	44.0	44.0	44.0
	Perempuan	28	56.0	56.0	100.0
	Total	50	100.0	100.0	

Status Gizi

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Kurus	4	8.0	8.0	8.0
	Normal	34	68.0	68.0	76.0
	Gemuk	12	24.0	24.0	100.0
	Total	50	100.0	100.0	

Uang Saku

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	ya	50	100.0	100.0	100.0

Diberi Uang Saku

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Setiap hari	48	96.0	96.0	96.0
1-2 kali seminggu	1	2.0	2.0	98.0
5-6 kali seminggu	1	2.0	2.0	100.0
Total	50	100.0	100.0	

Jumlah Uang Saku

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 1000	1	2.0	2.0	2.0
5000	7	14.0	14.0	16.0
7000	3	6.0	6.0	22.0
8000	2	4.0	4.0	26.0
10000	31	62.0	62.0	88.0
12000	2	4.0	4.0	92.0
13000	1	2.0	2.0	94.0
15000	2	4.0	4.0	98.0
20000	1	2.0	2.0	100.0
Total	50	100.0	100.0	

Uang Saku yang Dihabiskan

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 1000	1	2.0	2.0	2.0
2000	2	4.0	4.0	6.0
3000	5	10.0	10.0	16.0
4000	2	4.0	4.0	20.0
5000	24	48.0	48.0	68.0
6000	3	6.0	6.0	74.0
7000	3	6.0	6.0	80.0
8000	6	12.0	12.0	92.0
9000	1	2.0	2.0	94.0
10000	3	6.0	6.0	100.0
Total	50	100.0	100.0	

Kebiasaan Jajan

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid sering	29	58.0	58.0	58.0
jarang	21	42.0	42.0	100.0
Total	50	100.0	100.0	

Kualitas Jajanan

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Baik	14	28.0	28.0	28.0
Kurang	36	72.0	72.0	100.0
Total	50	100.0	100.0	

Analisis Bivariat

Kebiasaan Jajan*status gizi

Crosstab

Count

		status gizi			Total
		Kurus	Normal	Gemuk	
kebiasaan jajan	sering	2	21	6	29
	jarang	2	13	6	21
Total		4	34	12	50

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	.618 ^a	2	.734
Likelihood Ratio	.615	2	.735
Linear-by-Linear Association	.112	1	.738
N of Valid Cases	50		

a. 2 cells (33.3%) have expected count less than 5. The minimum expected count is 1.68.

Kualitas Jajanan*status gizi

Crosstab

Count

		status gizi			Total
		Kurus	Normal	Gemuk	
kualitas jajanan	Baik	1	9	4	14
	Kurang	3	25	8	36
Total		4	34	12	50

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	.227 ^a	2	.893
Likelihood Ratio	.222	2	.895
Linear-by-Linear Association	.191	1	.662
N of Valid Cases	50		

a. 3 cells (50.0%) have expected count less than 5. The minimum expected count is 1.12.