

Analisa Univariat

Jenis Kelamin

	Frequency	Percent	Valid Percent	Cumulative Percent
Laki-laki	58	53.2	53.2	53.2
Valid Perempuan	51	46.8	46.8	100.0
Total	109	100.0	100.0	

KM

	Frequency	Percent	Valid Percent	Cumulative Percent
Tidak Beragam	58	53.2	53.2	53.2
Valid Beragam	51	46.8	46.8	100.0
Total	109	100.0	100.0	

Tingkat Kesukaan Makan Siang

	Frequency	Percent	Valid Percent	Cumulative Percent
Tidak Suka	52	47.7	47.7	47.7
Valid Suka	57	52.3	52.3	100.0
Total	109	100.0	100.0	

Sisa Makan Siang

	Frequency	Percent	Valid Percent	Cumulative Percent
Sisa Banyak	57	52.3	52.3	52.3
Valid Sisa Sedikit	52	47.7	47.7	100.0
Total	109	100.0	100.0	

Zat Gizi yang Hilang dari Sisa Makan Siang

	N	Minimum	Maximum	Mean	Std. Deviation
Sisa Energi	109	.0	219.3	51.174	44.0856
Sisa Protein	109	.0	10.9	2.497	2.2165
Sisa Vitamin A	109	.0	47.0	6.055	11.3240
Sisa Fe	109	.0	1.6	.424	.3049
Sisa Serat	109	.0	.5	.151	.1168
Valid N (listwise)	109				

Biaya Sisa Makan Siang

	N	Minimum	Maximum	Mean	Std. Deviation
Biaya Sisa Makanan Pokok	109	0	394	65.41	88.036
Biaya Sisa Lauk Hewani	109	0	676	166.55	171.091
Biaya Sisa Lauk Nabati	109	0	650	126.29	154.713
Valid N (listwise)	109				

Tingkat Kesukaan Lauk Hewani berdasarkan jenis kelamin Laki-laki

	Frequency	Percent	Valid Percent	Cumulative Percent
tidak suka	4	6.9	6.9	6.9
Valid suka	54	93.1	93.1	100.0
Total	58	100.0	100.0	

Tingkat Kesukaan Lauk Nabati berdasarkan jenis kelamin Laki-laki

	Frequency	Percent	Valid Percent	Cumulative Percent
tidak suka	13	22.4	22.4	22.4
Valid suka	45	77.6	77.6	100.0
Total	58	100.0	100.0	

Tingkat Kesukaan Lauk Hewani berdasarkan jenis kelamin Perempuan

	Frequency	Percent	Valid Percent	Cumulative Percent
tidak suka	5	9.8	9.8	9.8
Valid suka	46	90.2	90.2	100.0
Total	51	100.0	100.0	

Tingkat Kesukaan Lauk Nabati berdasarkan jenis kelamin Perempuan

	Frequency	Percent	Valid Percent	Cumulative Percent
tidak suka	21	41.2	41.2	41.2
Valid suka	30	58.8	58.8	100.0
Total	51	100.0	100.0	

Tingkat Kesukaan Lauk Hewani berdasarkan Kebiasaan Makan Tidak Beragam

	Frequency	Percent	Valid Percent	Cumulative Percent
tidak suka	5	8.6	8.6	8.6
Valid suka	53	91.4	91.4	100.0
Total	58	100.0	100.0	

Tingkat Kesukaan Lauk Nabati berdasarkan Kebiasaan Makan Tidak Beragam

	Frequency	Percent	Valid Percent	Cumulative Percent
tidak suka	23	39.7	39.7	39.7
Valid suka	35	60.3	60.3	100.0
Total	58	100.0	100.0	

Tingkat Kesukaan Lauk Hewani berdasarkan Kebiasaan Makan Beragam

	Frequency	Percent	Valid Percent	Cumulative Percent
tidak suka	4	7.8	7.8	7.8
Valid suka	47	92.2	92.2	100.0
Total	51	100.0	100.0	

Tingkat Kesukaan Lauk Nabati berdasarkan Kebiasaan Makan Beragam

	Frequency	Percent	Valid Percent	Cumulative Percent
tidak suka	11	21.6	21.6	21.6
Valid suka	40	78.4	78.4	100.0
Total	51	100.0	100.0	

Sisa Lauk Hewani Berdasarkan Tingkat Kesukaan "Tidak Suka"

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid banyak	9	100.0	100.0	100.0

Sisa Lauk Hewani Berdasarkan Tingkat Kesukaan "Suka"

	Frequency	Percent	Valid Percent	Cumulative Percent
banyak	34	34.0	34.0	34.0
Valid sedikit	66	66.0	66.0	100.0
Total	100	100.0	100.0	

Sisa Lauk Nabati Berdasarkan Tingkat Kesukaan "Tidak Suka"

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid banyak	34	100.0	100.0	100.0

Sisa Lauk Nabati Berdasarkan Tingkat Kesukaan "Suka"

	Frequency	Percent	Valid Percent	Cumulative Percent
banyak	51	68.0	68.0	68.0
Valid sedikit	24	32.0	32.0	100.0
Total	75	100.0	100.0	

Analisa Bivariat

Jenis Kelamin * Tingkat Kesukaan Makan Siang Crosstabulation

		Tingkat Kesukaan Makan Siang		Total	
		Tidak Suka	Suka		
Jenis Kelamin	Laki-laki	Count	22	36	58
		% within Jenis Kelamin	37.9%	62.1%	100.0%
	Perempuan	Count	30	21	51
		% within Jenis Kelamin	58.8%	41.2%	100.0%
Total	Count	52	57	109	
	% within Jenis Kelamin	47.7%	52.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.748 ^a	1	.029		
Continuity Correction ^b	3.948	1	.047		
Likelihood Ratio	4.780	1	.029		
Fisher's Exact Test				.036	.023
Linear-by-Linear Association	4.705	1	.030		
N of Valid Cases	109				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 24.33.

b. Computed only for a 2x2 table

KM * Tingkat Kesukaan Makan Siang Crosstabulation

		Tingkat Kesukaan Makan Siang		Total	
		Tidak Suka	Suka		
KM	Tidak Beragam	Count	32	26	58
		% within KM	55.2%	44.8%	100.0%
	Beragam	Count	20	31	51
		% within KM	39.2%	60.8%	100.0%
Total	Count	52	57	109	
	% within KM	47.7%	52.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.770 ^a	1	.096		
Continuity Correction ^b	2.167	1	.141		
Likelihood Ratio	2.784	1	.095		
Fisher's Exact Test				.125	.070
Linear-by-Linear Association	2.744	1	.098		
N of Valid Cases	109				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 24.33.

b. Computed only for a 2x2 table

Tingkat Kesukaan Makan Siang * Sisa Makan Siang Crosstabulation

			Sisa Makan Siang		Total
			Sisa Banyak	Sisa Sedikit	
Tingkat Kesukaan Makan Siang	Tidak Suka	Count	40	12	52
		% within Tingkat Kesukaan Makan Siang	76.9%	23.1%	100.0%
	Suka	Count	17	40	57
		% within Tingkat Kesukaan Makan Siang	29.8%	70.2%	100.0%
Total		Count	57	52	109
		% within Tingkat Kesukaan Makan Siang	52.3%	47.7%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	24.179 ^a	1	.000		
Continuity Correction ^b	22.328	1	.000		
Likelihood Ratio	25.227	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	23.957	1	.000		
N of Valid Cases	109				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 24.81.

b. Computed only for a 2x2 table