

Analisis Bivariat

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
kategori_Pengetahuan_gizi * Kebiasaan_jajan	50	100,0%	0	0,0%	50	100,0%

kategori_Pengetahuan_gizi * Kebiasaan_jajan Crosstabulation

		Kebiasaan_jajan		Total
		Sering(4)	Jarang(<4)	
<70(kurang)	Count	11	1	12
	Expected Count	7,0	5,0	12,0
	% within kategori_Pengetahuan_gizi	91,7%	8,3%	100,0%
>70(Baik)	Count	18	20	38
	Expected Count	22,0	16,0	38,0
	% within kategori_Pengetahuan_gizi	47,4%	52,6%	100,0%
Total	Count	29	21	50
	Expected Count	29,0	21,0	50,0
	% within kategori_Pengetahuan_gizi	58,0%	42,0%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	7,347 ^a	1	,007		
Continuity Correction ^b	5,641	1	,018		
Likelihood Ratio	8,571	1	,003		
Fisher's Exact Test				,008	,006
Linear-by-Linear Association	7,200	1	,007		
N of Valid Cases	50				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 5,04.

b. Computed only for a 2x2 table

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
besar_uang_saku * Kebiasaan_jajan	50	100,0%	0	0,0%	50	100,0%

besar_uang_saku * Kebiasaan_jajan Crosstabulation

		Kebiasaan_jajan		Total
		Sering(4)	Jarang(<4)	
Tinggi	Count	22	8	30
	Expected Count	17,4	12,6	30,0
	% within besar_uang_saku	73,3%	26,7%	100,0%
Rendah	Count	7	13	20
	Expected Count	11,6	8,4	20,0
	% within besar_uang_saku	35,0%	65,0%	100,0%
Total	Count	29	21	50
	Expected Count	29,0	21,0	50,0
	% within besar_uang_saku	58,0%	42,0%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1-sided)
Pearson Chi-Square	7,239 ^a	1	,007		
Continuity Correction ^b	5,751	1	,016		
Likelihood Ratio	7,336	1	,007		
Fisher's Exact Test				,010	,008
Linear-by-Linear Association	7,094	1	,008		
N of Valid Cases	50				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 8,40.

b. Computed only for a 2x2 table

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Pekerjaan_Ayah * Kebiasaan_jajan	50	100,0%	0	0,0%	50	100,0%

Pekerjaan_Ayah * Kebiasaan_jajan Crosstabulation

			Kebiasaan_jajan		Total
			Sering (>4)	Jarang (<4)	
Tidak Bekerja	Count		3	9	12
	Expected Count		7,0	5,0	12,0
	% within Pekerjaan_Ayah		25,0%	75,0%	100,0%
Bekerja	Count		26	12	38
	Expected Count		22,0	16,0	38,0
	% within Pekerjaan_Ayah		68,4%	31,6%	100,0%
Total	Count		29	21	50
	Expected Count		29,0	21,0	50,0
	% within Pekerjaan_Ayah		58,0%	42,0%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	7,059 ^a	1	,008		
Continuity Correction ^b	5,389	1	,020		
Likelihood Ratio	7,135	1	,008		
Fisher's Exact Test				,017	,010
Linear-by-Linear Association	6,917	1	,009		
N of Valid Cases	50				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 5,04.

b. Computed only for a 2x2 table

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Pekerjaan_Ibu * Kebiasaan_jajan	50	100,0%	0	0,0%	50	100,0%

Pekerjaan_Ibu * Kebiasaan_jajan Crosstabulation

		Kebiasaan_jajan		Total
		Sering (>4)	Jarang (<4)	
Tidak bekerja	Count	9	13	22
	Expected Count	12,8	9,2	22,0
	% within Pekerjaan_Ibu	40,9%	59,1%	100,0%
Bekerja	Count	20	8	28
	Expected Count	16,2	11,8	28,0
	% within Pekerjaan_Ibu	71,4%	28,6%	100,0%
Total	Count	29	21	50
	Expected Count	29,0	21,0	50,0
	% within Pekerjaan_Ibu	58,0%	42,0%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4,711 ^a	1	,030		
Continuity Correction ^b	3,541	1	,060		
Likelihood Ratio	4,759	1	,029		
Fisher's Exact Test				,044	,030
Linear-by-Linear Association	4,617	1	,032		
N of Valid Cases	50				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 9,24.

b. Computed only for a 2x2 table

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Pendidikan_Ayah * Kebiasaan_jajan	50	100,0%	0	0,0%	50	100,0%

Pendidikan_Ayah * Kebiasaan_jajan Crosstabulation

		Kebiasaan_jajan		Total
		Sering (>4)	Jarang (<4)	
Rendah (SD-SMP)	Count	19	9	28
	Expected Count	16,2	11,8	28,0
	% within Pendidikan_Ayah	67,9%	32,1%	100,0%
Tinggi (SMA-Lainnya)	Count	10	12	22
	Expected Count	12,8	9,2	22,0
	% within Pendidikan_Ayah	45,5%	54,5%	100,0%
Total	Count	29	21	50
	Expected Count	29,0	21,0	50,0
	% within Pendidikan_Ayah	58,0%	42,0%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2,538 ^a	1	,111		
Continuity Correction ^b	1,702	1	,192		
Likelihood Ratio	2,548	1	,110		
Fisher's Exact Test				,152	,096
Linear-by-Linear Association	2,487	1	,115		
N of Valid Cases	50				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 9,24.

b. Computed only for a 2x2 table

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Pendidikan_Ibu * Kebiasaan_jajan	50	100,0%	0	0,0%	50	100,0%

Pendidikan_Ibu * Kebiasaan_jajan Crosstabulation

			Kebiasaan_jajan		Total
			Sering(4)	Jarang(<4)	
Pendidikan_Ibu	Rendah (SD-SMP)	Count	20	10	30
		Expected Count	17,4	12,6	30,0
		% within Pendidikan_Ibu	66,7%	33,3%	100,0%
	Tinggi (SMA-Lainnya)	Count	9	11	20
		Expected Count	11,6	8,4	20,0
		% within Pendidikan_Ibu	45,0%	55,0%	100,0%
Total		Count	29	21	50
		Expected Count	29,0	21,0	50,0
		% within Pendidikan_Ibu	58,0%	42,0%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2,313 ^a	1	,128		
Continuity Correction ^b	1,509	1	,219		
Likelihood Ratio	2,313	1	,128		
Fisher's Exact Test				,154	,110
Linear-by-Linear Association	2,266	1	,132		
N of Valid Cases	50				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 8,40.

b. Computed only for a 2x2 table

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Pendapatan_orangtua * Kebiasaan_jajan	50	100,0%	0	0,0%	50	100,0%

Pendapatan_orangtua * Kebiasaan_jajan Crosstabulation

		Kebiasaan_jajan		Total
		Sering(4)	Jarang(<4)	
Rendah	Count	9	14	23
	Expected Count	13,3	9,7	23,0
	% within Pendapatan_orangtua	39,1%	60,9%	100,0%
Tinggi	Count	20	7	27
	Expected Count	15,7	11,3	27,0
	% within Pendapatan_orangtua	74,1%	25,9%	100,0%
Total	Count	29	21	50
	Expected Count	29,0	21,0	50,0
	% within Pendapatan_orangtua	58,0%	42,0%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1-sided)
Pearson Chi-Square	6,226 ^a	1	,013		
Continuity Correction ^b	4,874	1	,027		
Likelihood Ratio	6,337	1	,012		
Fisher's Exact Test				,021	,013
Linear-by-Linear Association	6,101	1	,014		
N of Valid Cases	50				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 9,66.

b. Computed only for a 2x2 table

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Pengaruh_teman_sebaya * Kebiasaan_jajan	50	100,0%	0	0,0%	50	100,0%

Pengaruh_teman_sebaya * Kebiasaan_jajan Crosstabulation

			Kebiasaan_jajan		Total
			Sering(4)	Jarang(<4)	
Pengaruh_teman_sebaya	Ada pengaruh	Count	23	10	33
		Expected Count	19,1	13,9	33,0
		% within Pengaruh_teman_sebaya	69,7%	30,3%	100,0%
	Tidak ada pengaruh	Count	6	11	17
		Expected Count	9,9	7,1	17,0
		% within Pengaruh_teman_sebaya	35,3%	64,7%	100,0%
Total	Count	29	21	50	
	Expected Count	29,0	21,0	50,0	
	% within Pengaruh_teman_sebaya	58,0%	42,0%	100,0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5,451 ^a	1	,020		
Continuity Correction ^b	4,131	1	,042		
Likelihood Ratio	5,470	1	,019		
Fisher's Exact Test				,033	,021
Linear-by-Linear Association	5,342	1	,021		
N of Valid Cases	50				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 7,14.

b. Computed only for a 2x2 table

Frequencies


Statistics

nominal_pendapatan

N	Valid	50
	Missing	0
Mean		2180000,00
Std. Error of Mean		164180,090
Median		2500000,00
Std. Deviation		1160928,552
Variance		1347755102040,816
Skewness		,216
Std. Error of Skewness		,337
Minimum		700000
Maximum		5000000

nominal_pendapatan

	Frequency	Percent	Valid Percent	Cumulative Percent
700000	3	6,0	6,0	6,0
800000	1	2,0	2,0	8,0
900000	1	2,0	2,0	10,0
1000000	15	30,0	30,0	40,0
1500000	2	4,0	4,0	44,0
2000000	1	2,0	2,0	46,0
Valid 2500000	6	12,0	12,0	58,0
3000000	11	22,0	22,0	80,0
3200000	1	2,0	2,0	82,0
3500000	6	12,0	12,0	94,0
4000000	2	4,0	4,0	98,0
5000000	1	2,0	2,0	100,0
Total	50	100,0	100,0	



Descriptives

Descriptive Statistics

	N	Minimum	Maximum	Mean	Std. Deviation
nominal_pendapatan	50	700000	5000000	2180000,00	1160928,552
Valid N (listwise)	50				