

LAMPIRAN SPSS

Univariate - Frequency Table

Umur

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	< 35 tahun	27	90,0	90,0	90,0
	>= 35 tahun	3	10,0	10,0	100,0
	Total	30	100,0	100,0	

Jenis Kelamin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Perempuan	27	90,0	90,0	90,0
	Laki-laki	3	10,0	10,0	100,0
	Total	30	100,0	100,0	

Pendidikan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	D3	11	36,7	36,7	36,7
	S1	19	63,3	63,3	100,0
	Total	30	100,0	100,0	

Lama Kerja

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	< 5 tahun	20	66,7	66,7	66,7
	>= 5 tahun	10	33,3	33,3	100,0
	Total	30	100,0	100,0	

Pengetahuan Perawat

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Kurang Baik	8	26,7	26,7	26,7
	Baik	22	73,3	73,3	100,0
	Total	30	100,0	100,0	

Sikap Perawat

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Kurang Baik	16	53,3	53,3	53,3
	Baik	14	46,7	46,7	100,0
	Total	30	100,0	100,0	

Kepatuhan Perawat

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Patuh	13	43,3	43,3	43,3
	Patuh	17	56,7	56,7	100,0
	Total	30	100,0	100,0	

Bivariate – Crosstabs

Umur * Kepatuhan Perawat

Crosstab

		Kepatuhan Perawat		Total
		Tidak Patuh	Patuh	
Umur < 35 tahun	Count	11	16	27
	% within Umur	40,7%	59,3%	100,0%
Umur >= 35 tahun	Count	2	1	3
	% within Umur	66,7%	33,3%	100,0%
Total	Count	13	17	30
	% within Umur	43,3%	56,7%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,739 ^b	1	,390		
Continuity Correction ^a	,060	1	,806		
Likelihood Ratio	,736	1	,391		
Fisher's Exact Test				,565	,397
Linear-by-Linear Association	,714	1	,398		
N of Valid Cases	30				

a. Computed only for a 2x2 table

b. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,30.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Umur (< 35 tahun / >= 35 tahun)	,344	,028	4,273
For cohort Kepatuhan Perawat = Tidak Patuh	,611	,243	1,534
For cohort Kepatuhan Perawat = Patuh	1,778	,348	9,079
N of Valid Cases	30		

Pendidikan * Kepatuhan Perawat

Crosstab

			Kepatuhan Perawat		Total
			Tidak Patuh	Patuh	
Pendidikan	D3	Count	4	7	11
		% within Pendidikan	36,4%	63,6%	100,0%
S1	Count	9	10	19	
	% within Pendidikan	47,4%	52,6%	100,0%	
Total	Count	13	17	30	
	% within Pendidikan	43,3%	56,7%	100,0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,344 ^b	1	,558		
Continuity Correction ^a	,042	1	,838		
Likelihood Ratio	,346	1	,556		
Fisher's Exact Test				,708	,421
Linear-by-Linear Association	,332	1	,564		
N of Valid Cases	30				

a. Computed only for a 2x2 table

b. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,77.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Pendidikan (D3 / S1)	,635	,138	2,912
For cohort Kepatuhan Perawat = Tidak Patuh	,768	,308	1,915
For cohort Kepatuhan Perawat = Patuh	1,209	,652	2,242
N of Valid Cases	30		

Lama Kerja * Kepatuhan Perawat

Crosstab

		Kepatuhan Perawat		Total
		Tidak Patuh	Patuh	
Lama Kerja	< 5 tahun	Count 10	10	20
		% within Lama Kerja 50,0%	50,0%	100,0%
	>= 5 tahun	Count 3	7	10
		% within Lama Kerja 30,0%	70,0%	100,0%
Total		Count 13	17	30
		% within Lama Kerja 43,3%	56,7%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1,086 ^b	1	,297		
Continuity Correction ^a	,424	1	,515		
Likelihood Ratio	1,111	1	,292		
Fisher's Exact Test				,440	,259
Linear-by-Linear Association	1,050	1	,306		
N of Valid Cases	30				

a. Computed only for a 2x2 table

b. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,33.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Lama Kerja (< 5 tahun / >= 5 tahun)	2,333	,466	11,693
For cohort Kepatuhan Perawat = Tidak Patuh	1,667	,587	4,731
For cohort Kepatuhan Perawat = Patuh	,714	,393	1,298
N of Valid Cases	30		

Pengetahuan Perawat * Kepatuhan Perawat

Crosstab

			Kepatuhan Perawat		Total
			Tidak Patuh	Patuh	
Pengetahuan Perawat	Kurang Baik	Count	2	6	8
		% within Pengetahuan Perawat	25.0%	75.0%	100.0%
	Baik	Count	11	11	22
		% within Pengetahuan Perawat	50.0%	50.0%	100.0%
Total		Count	13	17	30
		% within Pengetahuan Perawat	43.3%	56.7%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.493 ^b	1	.222		
Continuity Correction ^a	.649	1	.421		
Likelihood Ratio	1.558	1	.212		
Fisher's Exact Test				.407	.212
Linear-by-Linear Association	1.443	1	.230		
N of Valid Cases	30				

a. Computed only for a 2x2 table

b. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 3.47.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Pengetahuan Perawat (Kurang Baik / Baik)	.333	.055	2.027
For cohort Kepatuhan Perawat = Tidak Patuh	.500	.140	1.782
For cohort Kepatuhan Perawat = Patuh	1.500	.841	2.675
N of Valid Cases	30		

Sikap Perawat * Kepatuhan Perawat

Crosstab

			Kepatuhan Perawat		Total
			Tidak Patuh	Patuh	
Sikap Perawat	Kurang Baik	Count	11	5	16
		% within Sikap Perawat	68,8%	31,3%	100,0%
	Baik	Count	2	12	14
		% within Sikap Perawat	14,3%	85,7%	100,0%
Total		Count	13	17	30
		% within Sikap Perawat	43,3%	56,7%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	9,020 ^b	1	,003		
Continuity Correction ^a	6,938	1	,008		
Likelihood Ratio	9,696	1	,002		
Fisher's Exact Test				,004	,004
Linear-by-Linear Association	8,719	1	,003		
N of Valid Cases	30				

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 6,07.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Sikap Perawat (Kurang Baik / Baik)	13,200	2,112	82,500
For cohort Kepatuhan Perawat = Tidak Patuh	4,813	1,279	18,104
For cohort Kepatuhan Perawat = Patuh	,365	,171	,778
N of Valid Cases	30		