

HASIL SPSS**A. Distribusi frekuensi****1. Jenis Kelamin**

	Frequency	Percent	Valid Percent	Cumulative Percent
Laki-Laki	23	50,0	50,0	50,0
Valid Perempuan	23	50,0	50,0	100,0
Total	46	100,0	100,0	

2. Usia

	Frequency	Percent	Valid Percent	Cumulative Percent
10 Tahun	29	63,0	63,0	63,0
Valid 11 Tahun	17	37,0	37,0	100,0
Total	46	100,0	100,0	

3. Pendidikan Ibu

	Frequency	Percent	Valid Percent	Cumulative Percent
Rendah	15	32,6	32,6	32,6
Valid Tinggi	31	67,4	67,4	100,0
Total	46	100,0	100,0	

4. Pendapatan Orang Tua

	Frequency	Percent	Valid Percent	Cumulative Percent
< UMR	29	63,0	63,0	63,0
Valid > UMR	17	37,0	37,0	100,0
Total	46	100,0	100,0	

5. Riwayat ASI-Eksklusif

	Frequency	Percent	Valid Percent	Cumulative Percent
Tidak	32	69,6	69,6	69,6
Valid Iya	14	30,4	30,4	100,0
Total	46	100,0	100,0	

6. Riwayat MP-ASI

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid MP-ASI Dini	32	69,6	69,6	69,6
Tidak	14	30,4	30,4	100,0
Total	46	100,0	100,0	

7. Pengetahuan Gizi

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Kurang	24	52,2	52,2	52,2
Baik	22	47,8	47,8	100,0
Total	46	100,0	100,0	

8. Picky Eater

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Picky Eater	37	80,4	80,4	80,4
Tidak Picky Eater	9	19,6	19,6	100,0
Total	46	100,0	100,0	

9. Tingkat Kecukupan Energi

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Kurang	40	87,0	87,0	87,0
Cukup	6	13,0	13,0	100,0
Total	46	100,0	100,0	

B. Faktor-Faktor yang Mempengaruhi Kejadian *Picky Eater*

1. Pendidikan ibu dengan *Picky Eater*

		Picky Eater yang Dikategorikan		Total
		Picky Eater	Tidak Picky Eater	
Pendidikan Ibu	Rendah	12	3	15
	Tinggi	25	6	31
	Total	37	9	46

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,003 ^a	1	,959		
Continuity Correction ^b	,000	1	1,000		
Likelihood Ratio	,003	1	,959		
Fisher's Exact Test				1,000	,624
Linear-by-Linear Association	,003	1	,959		
N of Valid Cases	46				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 2,93.

b. Computed only for a 2x2 table

2. Pendapatan Orang Tua dengan *Picky Eater*

		Picky Eater yang Dikategorikan		Total
		Picky Eater	Tidak Picky Eater	
Pendapatan Orang Tua	< UMR	23	6	29
	> UMR	14	3	17
Total		37	9	46

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,063 ^a	1	,802		
Continuity Correction ^b	,000	1	1,000		
Likelihood Ratio	,064	1	,801		
Fisher's Exact Test				1,000	,561
Linear-by-Linear Association	,062	1	,804		
N of Valid Cases	46				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 3,33.

b. Computed only for a 2x2 table

3. Riwayat ASI-Eksklusif dengan *Picky Eater*

		Picky Eater yang Dikategorikan		Total
		Picky Eater	Tidak Picky Eater	
Riwayat ASI-Eksklusif	Tidak	25	7	32
	Iya	12	2	14
Total		37	9	46

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,356 ^a	1	,550		
Continuity Correction ^b	,037	1	,847		
Likelihood Ratio	,373	1	,541		
Fisher's Exact Test				,701	,437
Linear-by-Linear Association	,349	1	,555		
N of Valid Cases	46				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 2,74.

b. Computed only for a 2x2 table

4. Riwayat MP-ASI dengan *Picky Eater*

		Picky Eater yang Dikategorikan		Total
		Picky Eater	Tidak Picky Eater	
Riwayat MP-ASI	MP-ASI Dini	25	7	32
	Tidak	12	2	14
Total		37	9	46

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,356 ^a	1	,550		
Continuity Correction ^b	,037	1	,847		
Likelihood Ratio	,373	1	,541		
Fisher's Exact Test				,701	,437
Linear-by-Linear Association	,349	1	,555		
N of Valid Cases	46				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 2,74.

b. Computed only for a 2x2 table

5. Pengetahuan Gizi dengan *Picky Eater*

		Picky Eater yang Dikategorikan		Total
		Picky Eater	Tidak Picky Eater	
Pengetahuan Gizi yang Dikategorikan	Kurang	18	6	24
	Baik	19	3	22
Total		37	9	46

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,942 ^a	1	,332	,464	,277
Continuity Correction ^b	,358	1	,550		
Likelihood Ratio	,959	1	,327		
Fisher's Exact Test					
Linear-by-Linear Association	,921	1	,337		
N of Valid Cases	46				

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 4,30.

b. Computed only for a 2x2 table

6. *Picky Eater* dengan Tingkat Kecukupan Energi

		Tingkat Kecukupan Energi yang Dikategorikan		Total
		Kurang	Cukup	
Picky Eater yang Dikategorikan	Picky Eater	33	4	37
	Tidak Picky Eater	7	2	9
Total		40	6	46

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,831 ^a	1	,362	,581	,333
Continuity Correction ^b	,130	1	,719		
Likelihood Ratio	,741	1	,389		
Fisher's Exact Test					
Linear-by-Linear Association	,813	1	,367		
N of Valid Cases	46				

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,17.

b. Computed only for a 2x2 table