

Lampiran 8

Hasil Analisis Univariat dan Bivariat**Umur**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Remaja Awal	10	18.5	18.5	18.5
	Remaja Tengah	44	81.5	81.5	100.0
	Total	54	100.0	100.0	

Jenis Kelamin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Laki-laki	40	74.1	74.1	74.1
	Perempuan	14	25.9	25.9	100.0
	Total	54	100.0	100.0	

Kelas Olahraga

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ringan	4	7.4	7.4	7.4
	Sedang	28	51.9	51.9	59.3
	Berat	17	31.5	31.5	90.7
	Berat Sekali	5	9.3	9.3	100.0
	Total	54	100.0	100.0	

Kualitas Menu

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Berkualitas	40	74.1	74.1	74.1
	Berkualitas	14	25.9	25.9	100.0
	Total	54	100.0	100.0	

Energi

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Sangat Kurang	16	29.6	29.6	29.6
	Kurang	14	25.9	25.9	55.6
	Baik	14	25.9	25.9	81.5
	Lebih	10	18.5	18.5	100.0
	Total	54	100.0	100.0	

Keberagaman

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Beragam	6	11.1	11.1	11.1
	Beragam	48	88.9	88.9	100.0
	Total	54	100.0	100.0	

Preferensi Menu I

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Baik	24	44.4	44.4	44.4
	Baik	30	55.6	55.6	100.0
	Total	54	100.0	100.0	

Preferensi Menu II

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Baik	30	55.6	55.6	55.6
	Baik	24	44.4	44.4	100.0
	Total	54	100.0	100.0	

Preferensi Menu III

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Baik	30	55.6	55.6	55.6
	Baik	24	44.4	44.4	100.0
	Total	54	100.0	100.0	

Preferensi Menu IV

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Baik	18	33.3	33.3	33.3
	Baik	36	66.7	66.7	100.0
	Total	54	100.0	100.0	

Preferensi Menu V

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Baik	26	48.1	48.1	48.1
	Baik	28	51.9	51.9	100.0
	Total	54	100.0	100.0	

Preferensi Menu II

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Baik	30	55.6	55.6	55.6
	Baik	24	44.4	44.4	100.0
	Total	54	100.0	100.0	

Preferensi Menu III

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Baik	30	55.6	55.6	55.6
	Baik	24	44.4	44.4	100.0
	Total	54	100.0	100.0	

Preferensi Menu IV

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Baik	18	33.3	33.3	33.3
	Baik	36	66.7	66.7	100.0
	Total	54	100.0	100.0	

Preferensi Menu V

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Baik	26	48.1	48.1	48.1
	Baik	28	51.9	51.9	100.0
	Total	54	100.0	100.0	

Preferensi Menu VI

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Baik	34	63.0	63.0	63.0
	Baik	20	37.0	37.0	100.0
	Total	54	100.0	100.0	

Preferensi Menu VII

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Baik	27	50.0	50.0	50.0
	Baik	27	50.0	50.0	100.0
	Total	54	100.0	100.0	

Status Gizi

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak Normal	51	94.4	94.4	94.4
	Normal	3	5.6	5.6	100.0
	Total	54	100.0	100.0	

Kualitas Menu dengan Status Gizi

Count

		Status Gizi		Total
		Tidak Normal	Normal	
Kualitas Menu	Tidak Berkualitas	37	3	40
	Berkualitas	14	0	14
Total		51	3	54

Preferensi Menu I dengan Status Gizi

Count

		Status Gizi		Total
		Tidak Normal	Normal	
Preferensi Menu I	Tidak Baik	22	2	24
	Baik	29	1	30
Total		51	3	54

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.635 ^a	1	.425		
Continuity Correction ^b	.040	1	.842		
Likelihood Ratio	.636	1	.425		
Fisher's Exact Test				.579	.415
Linear-by-Linear Association	.624	1	.430		
N of Valid Cases	54				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.33.

b. Computed only for a 2x2 table

Crosstab Preferensi Menu II dengan Status Gizi

Count

		Status Gizi		Total
		Tidak Normal	Normal	
Preferensi Menu II	Tidak Baik	28	2	30
	Baik	23	1	24
Total		51	3	54

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.159 ^a	1	.690		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.163	1	.687		
Fisher's Exact Test				1.000	.585
Linear-by-Linear Association	.156	1	.693		
N of Valid Cases	54				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.33.

b. Computed only for a 2x2 table

Preferensi Menu III dengan Status Gizi

Count

		Status Gizi		Total
		Tidak Normal	Normal	
Preferensi Menu III	Tidak Baik	27	3	30
	Baik	24	0	24
Total		51	3	54

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.541 ^a	1	.111		
Continuity Correction ^b	.993	1	.319		
Likelihood Ratio	3.667	1	.055		
Fisher's Exact Test				.245	.164
Linear-by-Linear Association	2.494	1	.114		
N of Valid Cases	54				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.33.

b. Computed only for a 2x2 table

Preferensi Menu IV dengan Status Gizi

Count

		Status Gizi		Total
		Tidak Normal	Normal	
Preferensi Menu IV	Tidak Baik	17	1	18
	Baik	34	2	36
Total		51	3	54

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.000 ^a	1	1.000		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.000	1	1.000		
Fisher's Exact Test				1.000	.712
Linear-by-Linear Association	.000	1	1.000		
N of Valid Cases	54				

- a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.00.
 b. Computed only for a 2x2 table

Preferensi Menu V dengan Status Gizi

Count

		Status Gizi		Total
		Tidak Normal	Normal	
Preferensi Menu V	Tidak Baik	25	1	26
	Baik	26	2	28
Total		51	3	54

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.279 ^a	1	.597		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.285	1	.593		
Fisher's Exact Test				1.000	.528
Linear-by-Linear Association	.274	1	.601		
N of Valid Cases	54				

- a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.44.
 b. Computed only for a 2x2 table

Preferensi Menu VI dengan Status Gizi

Count

		Status Gizi		Total
		Tidak Normal	Normal	
Preferensi Menu VI	Tidak Baik	31	3	34
	Baik	20	0	20
Total		51	3	54

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.869 ^a	1	.172		
Continuity Correction ^b	.565	1	.452		
Likelihood Ratio	2.879	1	.090		
Fisher's Exact Test				.287	.241
Linear-by-Linear Association	1.834	1	.176		
N of Valid Cases	54				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.11.

b. Computed only for a 2x2 table

Preferensi Menu VII dengan Status Gizi

Count

		Status Gizi		Total
		Tidak Normal	Normal	
Preferensi Menu VII	Tidak Baik	25	2	27
	Baik	26	1	27
Total		51	3	54

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.353 ^a	1	.552		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.359	1	.549		
Fisher's Exact Test				1.000	.500
Linear-by-Linear Association	.346	1	.556		
N of Valid Cases	54				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.50.

b. Computed only for a 2x2 table