

NORMALITAS

Tests of Normality

Peerlakuan	Kolmogorov-Smirnov ^a			Shapiro-Wilk			
	Statistic	df	Sig.	Statistic	df	Sig.	
Sebelum	1	.209	6	.200*	.907	6	.415
	2	.223	6	.200*	.908	6	.421
Sesudah	1	.209	6	.200*	.907	6	.415
	2	.254	6	.200*	.866	6	.212
Selisih	1	.407	6	.002	.640	6	.001
	2	.333	6	.036	.827	6	.101

a. Lilliefors Significance Correction

*. This is a lower bound of the true significance.

HOMOGENITAS

	t-test for Equality of Means						
	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
						Lower	Upper
Sebelum	.728	10	.484	.50000	.68718	-1.03114	2.03114
	.728	9.988	.484	.50000	.68718	-1.03140	2.03140

HIPOTESIS I

Paired Samples Test

	Paired Differences					t	df	Sig. (2-tailed)
	Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
				Lower	Upper			
Pair 1 Sebelum1 - Sessudah1	-3.33333	.51640	.21082	-3.87526	-2.79141	-15.811	5	.000

HIPOTESIS II

Paired Samples Test

	Paired Differences					t	df	Sig. (2-tailed)
	Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
				Lower	Upper			
Pair 1 Sebelum2 - Sesudah2	-4.00000	.63246	.25820	-4.66372	-3.33628	-15.492	5	.000

HIPOTESIS III

Independent Samples Test

	Levene's Test for Equality of Variances		t-test for Equality of Means						
	F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
								Lower	Upper
Selisih									
Equal variances assumed	.250	.628	-2.000	10	.073	-.66667	.33333	-1.40938	.07605
Equal variances not assumed			-2.000	9.615	.075	-.66667	.33333	-1.41343	.08009

FREKUENSI

Statistics

		Sebelum1	Sesudah1	Selisih1	Sebelum2	Sesudah2	Selisih2
N	Valid	6	6	6	6	6	6
	Missing	6	6	6	6	6	6
Mean		5.3333	8.6667	3.3333	4.8333	8.8333	4.0000
Std. Deviation		1.21106	1.21106	.51640	1.16905	.75277	.63246
Minimum		4.00	7.00	3.00	3.00	8.00	3.00
Maximum		7.00	10.00	4.00	6.00	10.00	5.00