

**Lampiran 1. Uji Normalitas
One-Sample Kolmogorov-Smirnov Test**

		BI	PG	GM	AF1	AF2	AF_TOTA L	IMT_SD	IMT_U	BF	KEL_G M	KEL_IM T	KEL_BF
n		12	12	12	12	12	12	12	12	12	12	12	12
Normal Parameters ^{a,b}	Mean	1.33	73.42	1.33	1.2102	1.5314	1.4465	-.5683	19.6250	22.99	1.3333	20.4583	23.0833
	Std. Deviation	.492	13.608	.492	.12837	.19070	.18410	.92040	1.87186	3.538	.49237	2.12580	3.36988
Most Extreme Differences	Absolute	.417	.213	.417	.172	.156	.160	.151	.182	.167	.417	.269	.153
	Positive	.417	.171	.417	.103	.156	.142	.097	.182	.167	.417	.269	.153
	Negative	-.249	-.213	-.249	-.172	-.113	-.160	-.151	-.181	-.136	-.249	-.126	-.140
Test Statistic		.417	.213	.417	.172	.156	.160	.151	.182	.167	.417	.269	.153
Asymp. Sig. (2-tailed)		.000 ^c	.139 ^c	.000 ^c	.200 ^{c,d}	.200 ^{c,d}	.200 ^{c,d}	.200 ^{c,d}	.200 ^{c,d}	.200 ^{c,d}	.000 ^c	.016 ^c	.200 ^{c,d}

a. Test distribution is Normal.

b. Calculated from data.

c. Lilliefors Significance Correction.

d. This is a lower bound of the true significance.

Lampiran 2. Analisis Univariat

Descriptive Statistics

	N	Minimum	Maximum	Mean	Std. Deviation
BI	12	1	2	1.33	.492
PG	12	52	88	73.42	13.608
GM	12	1	2	1.33	.492
AF_TOTAL	12	1.18	1.71	1.4465	.18410
IMT_U	12	15.60	23.30	19.6250	1.87186
BF	12	18	28	22.99	3.538
Valid N (listwise)	12				

Statistics

		BI	PG	GM	AF_TOTA L	IMT_U	BF
N	Valid	12	12	12	12	12	12
	Missing	0	0	0	0	0	0
Mean		1.33	73.42	1.33	1.4465	19.6250	22.99
Std. Deviation		.492	13.608	.492	.18410	1.87186	3.538
Skewness		.812	-.560	.812	.207	-.318	-.066
Std. Error of Skewness		.637	.637	.637	.637	.637	.637
Minimum		1	52	1	1.18	15.60	18
Maximum		2	88	2	1.71	23.30	28
Percentiles	25	1.00	57.00	1.00	1.2866	18.5500	19.25
	50	1.00	77.00	1.00	1.4168	19.7500	23.45
	75	2.00	84.00	2.00	1.6425	20.6750	26.30

Frequency Table

BI

	Frequency	Percent	Valid Percent	Cumulative Percent

Valid positif	8	66.7	66.7	66.7
negatif	4	33.3	33.3	100.0
Total	12	100.0	100.0	

Universitas
Esa Unggul

Universitas
Esa Unggul

Universitas
Esa Unggul

Universitas
Esa Unggul

Universitas
Esa Unggul

Universitas
Esa Unggul

Universitas
Esa Unggul

Universitas
Esa Unggul

Universitas
Esa Unggul

PG

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 52	1	8.3	8.3	8.3
56	2	16.7	16.7	25.0
60	1	8.3	8.3	33.3
75	1	8.3	8.3	41.7
76	1	8.3	8.3	50.0
78	1	8.3	8.3	58.3
84	3	25.0	25.0	83.3
88	2	16.7	16.7	100.0
Total	12	100.0	100.0	

GM

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid bulimia	8	66.7	66.7	66.7
binge eating	4	33.3	33.3	100.0
Total	12	100.0	100.0	

AF_TOTAL



	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 1.18	1	8.3	8.3	8.3
1.26	1	8.3	8.3	16.7
1.29	1	8.3	8.3	25.0
1.29	1	8.3	8.3	33.3
1.34	1	8.3	8.3	41.7
1.40	1	8.3	8.3	50.0
1.43	1	8.3	8.3	58.3
1.50	1	8.3	8.3	66.7
1.62	1	8.3	8.3	75.0
1.65	1	8.3	8.3	83.3
1.69	1	8.3	8.3	91.7
1.71	1	8.3	8.3	100.0
Total	12	100.0	100.0	



IMT_U



		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	15.60	1	8.3	8.3	8.3
	17.90	1	8.3	8.3	16.7
	18.30	1	8.3	8.3	25.0
	19.30	2	16.7	16.7	41.7
	19.60	1	8.3	8.3	50.0
	19.90	1	8.3	8.3	58.3
	20.20	1	8.3	8.3	66.7
	20.60	1	8.3	8.3	75.0
	20.70	1	8.3	8.3	83.3
	20.80	1	8.3	8.3	91.7
	23.30	1	8.3	8.3	100.0
Total		12	100.0	100.0	

BF

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	18	1	8.3	8.3	8.3
	19	2	16.7	16.7	25.0
	20	1	8.3	8.3	33.3
	21	1	8.3	8.3	41.7
	23	1	8.3	8.3	50.0
	24	1	8.3	8.3	58.3
	25	1	8.3	8.3	66.7
	26	1	8.3	8.3	75.0
	26	1	8.3	8.3	83.3
	27	1	8.3	8.3	91.7
	28	1	8.3	8.3	100.0
Total		12	100.0	100.0	







Lampiran 3. Analisis Bivariat

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
BI * GM	12	100.0%	0	0.0%	12	100.0%

BI * GM Crosstabulation

Count

		GM		Total
		bulimia	binge eating	
BI	positif	7	1	8
	negatif	1	3	4
Total		8	4	12

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.688 ^a	1	.030	.067	.067
Continuity Correction ^b	2.297	1	.130		
Likelihood Ratio	4.749	1	.029		
Fisher's Exact Test					
Linear-by-Linear Association	4.297	1	.038		
N of Valid Cases	12				

a. 3 cells (75.0%) have expected count less than 5. The minimum expected count is 1.33.

b. Computed only for a 2x2 table

Correlations

			PG	GM
Spearman's rho	PG	Correlation Coefficient	1.000	-.440
		Sig. (2-tailed)	.	.152
		N	12	12
		GM	Correlation Coefficient	-.440
		Sig. (2-tailed)	.152	.
		N	12	12

Group Statistics

GM	N	Mean	Std. Deviation	Std. Error Mean
IMT_U bulimia	8	18.1750	2.45691	.86865
binge eating	4	21.1250	.87702	.43851

Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
IMT_U	Equal variances assumed	.995	.342	-2.282	10	.046	-2.95000	1.29271	-5.83033	-.06967
	Equal variances not assumed			-3.032	9.572	.013	-2.95000	.97306	-5.13133	-.76867

Group Statistics

	GM	N	Mean	Std. Deviation	Std. Error Mean
BF	bulimia	8	21.44	3.118	1.102
	binge eating	4	26.10	1.977	.988

Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
BF	Equal variances assumed	2.612	.137	-2.696	10	.022	-4.663	1.729	-8.516	-.809
	Equal variances not assumed			-3.149	9.083	.012	-4.663	1.480	-8.007	-1.318

Descriptive Statistics

	Mean	Std. Deviation	N
AKT V	1.4983	.20440	12
BF	22.9917	3.53848	12

Correlations

		AKTV	BF
AKT V	Pearson Correlation	1	-.576
	Sig. (2-tailed)		.050
	N	12	12
BF	Pearson Correlation	-.576	1
	Sig. (2-tailed)	.050	
	N	12	12

Descriptive Statistics

	Mean	Std. Deviation	N
AKTV	1.4983	.20440	12
IMT_ U	19.1583	2.48210	12

