

Gambaran Faktor Usia

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid >= 45 tahun	27	42.9	42.9	42.9
< 45 tahun	36	57.1	57.1	100.0
Total	63	100.0	100.0	

Gambaran Faktor Jarak Monitor

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid < 50 cm	41	65.1	65.1	65.1
>= 50 cm	22	34.9	34.9	100.0
Total	63	100.0	100.0	

Gambaran Faktor Tingkat Pencahayaan

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid < 300 lux	45	71.4	71.4	71.4
>= 300 lux	18	28.6	28.6	100.0
Total	63	100.0	100.0	

Gambaran Faktor Kelainan Refraksi

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid ada kelainan refraksi	48	76.2	76.2	76.2
tidak ada kelainan refraksi	15	23.8	23.8	100.0
Total	63	100.0	100.0	

Gambaran Faktor Istirahat Mata

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid tidak istirahat	14	22.2	22.2	22.2
istirahat	49	77.8	77.8	100.0
Total	63	100.0	100.0	

Gambaran Keluhan Kelelahan Mata

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid mengalami keluhan	42	66.7	66.7	66.7
tidak mengalami keluhan	21	33.3	33.3	100.0
Total	63	100.0	100.0	

Hubungan Usia Dengan Keluhan Kelelahan Mata

Crosstab

		Total Keluhan		Total	
		Mengalami Keluhan	Tidak Mengalami Keluhan		
Usia	≥ 45 tahun	Count	24	3	27
		% within Usia	88.9%	11.1%	100.0%
		% within Total Keluhan	57.1%	14.3%	42.9%
	< 45 tahun	Count	18	18	36
		% within Usia	50.0%	50.0%	100.0%
		% within Total Keluhan	42.9%	85.7%	57.1%
Total	Count	42	21	63	
	% within Usia	66.7%	33.3%	100.0%	
	% within Total Keluhan	100.0%	100.0%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	10.500 ^a	1	.001		
Continuity Correction ^b	8.823	1	.003		
Likelihood Ratio	11.457	1	.001		
Fisher's Exact Test				.001	.001
Linear-by-Linear Association	10.333	1	.001		
N of Valid Cases	63				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 9,00.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Usia (≥ 45 tahun / < 45 tahun)	8.000	2.040	31.373
For cohort Total Keluhan = mengalami keluhan	1.778	1.249	2.530
For cohort Total Keluhan = tidak mengalami keluhan	.222	.073	.678
N of Valid Cases	63		

Hubungan Jarak Monitor Dengan Keluhan Kelelahan Mata

Crosstab

		Total Keluhan		Total	
		Mengalami Keluhan	Tidak Mengalami Keluhan		
Jarak Monitor	< 50 cm	Count	32	9	41
		% within Jarak Monitor	78.0%	22.0%	100.0%
		% within Total Keluhan	76.2%	42.9%	65.1%
	≥ 50 cm	Count	10	12	22
		% within Jarak Monitor	45.5%	54.5%	100.0%
		% within Total Keluhan	23.8%	57.1%	34.9%
Total	Count	42	21	63	
	% within Jarak Monitor	66.7%	33.3%	100.0%	
	% within Total Keluhan	100.0%	100.0%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	6.845 ^a	1	.009		
Continuity Correction ^b	5.457	1	.019		
Likelihood Ratio	6.729	1	.009		
Fisher's Exact Test				.012	.010
Linear-by-Linear Association	6.736	1	.009		
N of Valid Cases	63				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 7,33.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Jarak Monitor (< 50 cm / >= 50 cm)	4.267	1.394	13.058
For cohort Total Keluhan = mengalami keluhan	1.717	1.056	2.791
For cohort Total Keluhan = tidak mengalami keluhan	.402	.201	.804
N of Valid Cases	63		

Hubungan Tingkat Pencahayaan Dengan Keluhan Kelelahan Mata

Crosstab

		Total Keluhan		Total	
		Mengalami Keluhan	Tidak Mengalami Keluhan		
Tingkat Pencahayaan	< 300 lux	Count	34	11	45
		% within Tingkat Pencahayaan	75.6%	24.4%	100.0%
		% within Total Keluhan	81.0%	52.4%	71.4%
	>= 300 lux	Count	8	10	18
		% within Tingkat Pencahayaan	44.4%	55.6%	100.0%
		% within Total Keluhan	19.0%	47.6%	28.6%
Total		Count	42	21	63
		% within Tingkat Pencahayaan	66.7%	33.3%	100.0%
		% within Total Keluhan	100.0%	100.0%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5.600 ^a	1	.018		
Continuity Correction ^b	4.288	1	.038		
Likelihood Ratio	5.417	1	.020		
Fisher's Exact Test				.036	.020
Linear-by-Linear Association	5.511	1	.019		
N of Valid Cases	63				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 6,00.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Tingkat Pencahayaan (< 300 lux / >= 300 lux)	3.864	1.221	12.224
For cohort Total Keluhan = mengalami keluhan	1.700	.988	2.925
For cohort Total Keluhan = tidak mengalami keluhan	.440	.228	.851
N of Valid Cases	63		

Hubungan Kelainan Refraksi Dengan Keluhan Kelelahan Mata

Crosstab

		Total Keluhan		Total	
		Mengalami Keluhan	Tidak Mengalami Keluhan		
Kelainan Refraksi	ada kelainan refraksi	Count	36	12	48
		% within Kelainan Refraksi	75.0%	25.0%	100.0%
		% within Total Keluhan	85.7%	57.1%	76.2%
	tidak ada kelainan refraksi	Count	6	9	15
		% within Kelainan Refraksi	40.0%	60.0%	100.0%
		% within Total Keluhan	14.3%	42.9%	23.8%
Total	Count	42	21	63	
	% within Kelainan Refraksi	66.7%	33.3%	100.0%	
	% within Total Keluhan	100.0%	100.0%	100.0%	

Chi Square Test

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	6.300 ^a	1	.012		
Continuity Correction ^b	4.823	1	.028		
Likelihood Ratio	6.026	1	.014		
Fisher's Exact Test				.025	.015
Linear-by-Linear Association	6.200	1	.013		
N of Valid Cases	63				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5,00.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Kelainan Refraksi (ada kelainan refraksi / tidak ada kelainan refraksi)	4.500	1.326	15.277
For cohort Total Keluhan = mengalami keluhan	1.875	.988	3.559
For cohort Total Keluhan = tidak mengalami keluhan	.417	.219	.791
N of Valid Cases	63		

Hubungan Istirahat Mata Dengan Keluhan Kelelahan Mata

Crosstab

			Total Keluhan		Total
			Mengalami Keluhan	Tidak Mengalami Keluhan	
Istirahat Mata	tidak istirahat	Count	11	3	14
		% within Istirahat Mata	78.6%	21.4%	100.0%
		% within Total Keluhan	26.2%	14.3%	22.2%
	istirahat	Count	31	18	49
		% within Istirahat Mata	63.3%	36.7%	100.0%
		% within Total Keluhan	73.8%	85.7%	77.8%
Total	Count	42	21	63	
	% within Istirahat Mata	66.7%	33.3%	100.0%	
	% within Total Keluhan	100.0%	100.0%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.148 ^a	1	.284	.350	.230
Continuity Correction ^b	.563	1	.453		
Likelihood Ratio	1.215	1	.270		
Fisher's Exact Test					
Linear-by-Linear Association	1.130	1	.288		
N of Valid Cases	63				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,67.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Istirahat Mata (tidak istirahat / istirahat)	2.129	.524	8.655
For cohort Total Keluhan = mengalami keluhan	1.242	.878	1.757
For cohort Total Keluhan = tidak mengalami keluhan	.583	.200	1.698
N of Valid Cases	63		