

4.select case dan crostabulation

A. rokok 1 dan 4

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.068 ^a	1	.795		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.068	1	.794		
Fisher's Exact Test				1.000	.567
Linear-by-Linear Association	.065	1	.799		
N of Valid Cases	23				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 4.30.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for perilaku merokok untuk tabel 2x3 (bukan perokok / perokok berat)	1.250	.233	6.715
For cohort penyakit pada pekerja = tidak ispa	1.111	.507	2.435
For cohort penyakit pada pekerja = ISPA	.889	.362	2.184
N of Valid Cases	23		

B. rokok 2 dan 4

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	3.792 ^a	1	.052		
Continuity Correction ^b	2.514	1	.113		
Likelihood Ratio	3.717	1	.054		
Fisher's Exact Test				.075	.058
Linear-by-Linear Association	3.694	1	.055		
N of Valid Cases	39				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 4.31.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for perilaku merokok untuk tabel 2x3 (perokok ringan / perokok berat)	4.000	.953	16.791
For cohort penyakit pada pekerja = tidak ispa	1.600	.915	2.799
For cohort penyakit pada pekerja = ISPA	.400	.156	1.027
N of Valid Cases	39		

C. rokok 3 dan 4

Chi-Square Tests

	Value	df	Asymptotic Significance (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	5.266 ^a	1	.022		
Continuity Correction ^b	3.492	1	.062		
Likelihood Ratio	5.804	1	.016		
Fisher's Exact Test				.036	.028
Linear-by-Linear Association	5.064	1	.024		
N of Valid Cases	26				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 3.69.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for perilaku merokok untuk tabel 2x3 (perokok sedang / perokok berat)	11.000	1.103	109.674
For cohort penyakit pada pekerja = tidak ispa	1.833	1.057	3.180
For cohort penyakit pada pekerja = ISPA	.167	.024	1.169
N of Valid Cases	26		