

Hasil Pengolahan dengan SPSS V 21

UMUR

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	BERESIKO	59	70,2	70,2	70,2
	TIDAK BERESIKO	25	29,8	29,8	100,0
	Total	84	100,0	100,0	

JENIS KELAMIN

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	L	38	45,2	45,2	45,2
	P	46	54,8	54,8	100,0
	Total	84	100,0	100,0	

OBESITAS

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	OBESITAS > 27	47	56,0	56,0	56,0
	TIDAK OBESITAS <27	37	44,0	44,0	100,0
	Total	84	100,0	100,0	

HIPERTENSI

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	HIPERTENSI >140MMHG	19	22,6	22,6	22,6
	NORMAL <140MMHG	65	77,4	77,4	100,0
	Total	84	100,0	100,0	

RIWAYAT KELUARGA

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	YA	37	44,0	44,0
	TIDAK	47	56,0	100,0
	Total	84	100,0	

AKTIVITAS FISIK

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	RINGAN SEDANG	72	85,7	85,7
	BERAT	12	14,3	100,0
	Total	84	100,0	

DIABETES MELITUS

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	YA	53	63,1	63,1
	TIDAK	31	36,9	100,0
	Total	84	100,0	

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
UMUR * DIABETES MELITUS	84	100,0%	0	0,0%	84	100,0%

UMUR * DIABETES MELITUS Crosstabulation

		DIABETES MELITUS		Total	
		YA	TIDAK		
UMUR	BERESIKO	Count	36	23	59
		% within UMUR	61,0%	39,0%	100,0%
UMUR	TIDAK BERESIKO	Count	17	8	25
		% within UMUR	68,0%	32,0%	100,0%
Total		Count	53	31	84
		% within UMUR	63,1%	36,9%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,368 ^a	1	,544		
Continuity Correction ^b	,129	1	,719		
Likelihood Ratio	,372	1	,542		
Fisher's Exact Test				,626	,363
N of Valid Cases	84				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 9,23.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for UMUR (BERESIKO / TIDAK BERESIKO)	,737	,274	1,982
For cohort DIABETES MELITUS = YA	,897	,640	1,258
For cohort DIABETES MELITUS = TIDAK	1,218	,633	2,344
N of Valid Cases	84		

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
JENIS KELAMIN * DIABETES MELITUS	84	100,0%	0	0,0%	84	100,0%

JENIS KELAMIN * DIABETES MELITUS Crosstabulation

		DIABETES MELITUS		Total	
		YA	TIDAK		
JENIS KELAMIN	L	Count	26	12	38
		% within JENIS KELAMIN	68,4%	31,6%	100,0%
	P	Count	27	19	46
		% within JENIS KELAMIN	58,7%	41,3%	100,0%
Total		Count	53	31	84
		% within JENIS KELAMIN	63,1%	36,9%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,845 ^a	1	,358		
Continuity Correction ^b	,479	1	,489		
Likelihood Ratio	,850	1	,357		
Fisher's Exact Test				,375	,245
N of Valid Cases	84				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 14,02.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for JENIS KELAMIN (L / P)	1,525	,619	3,755
For cohort DIABETES MELITUS = YA	1,166	,843	1,613
For cohort DIABETES MELITUS = TIDAK	,765	,428	1,367
N of Valid Cases	84		

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
OBESITAS * DIABETES MELITUS	84	100,0%	0	0,0%	84	100,0%

OBESITAS * DIABETES MELITUS Crosstabulation

		DIABETES MELITUS		Total	
		YA	TIDAK		
OBESITAS	OBESITAS > 27	Count	45	2	47
		% within OBESITAS	95,7%	4,3%	100,0%
TIDAK OBESITAS <27	TIDAK OBESITAS <27	Count	8	29	37
		% within OBESITAS	21,6%	78,4%	100,0%
Total		Count	53	31	84
		% within OBESITAS	63,1%	36,9%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	48,848 ^a	1	,000		
Continuity Correction ^b	45,717	1	,000		
Likelihood Ratio	55,444	1	,000		
Fisher's Exact Test				,000	,000
N of Valid Cases	84				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 13,65.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for OBESITAS (OBESITAS > 27 / TIDAK OBESITAS <27)	81,563	16,169	411,423
For cohort DIABETES MELITUS = YA	4,428	2,391	8,202
For cohort DIABETES MELITUS = TIDAK	,054	,014	,213
N of Valid Cases	84		

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
HIPERTENSI * DIABETES MELITUS	84	100,0%	0	0,0%	84	100,0%

HIPERTENSI * DIABETES MELITUS Crosstabulation

		DIABETES MELITUS		Total	
		YA	TIDAK		
HIPERTENSI	HIPERTENSI >140MMHG	Count	17	2	19
		% within HIPERTENSI	89,5%	10,5%	100,0%
	NORMAL <140MMHG	Count	36	29	65
		% within HIPERTENSI	55,4%	44,6%	100,0%
Total		Count	53	31	84
		% within HIPERTENSI	63,1%	36,9%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	7,337 ^a	1	,007		
Continuity Correction ^b	5,946	1	,015		
Likelihood Ratio	8,478	1	,004		
Fisher's Exact Test				,007	,005
N of Valid Cases	84				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 7,01.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for HIPERTENSI (HIPERTENSI >140MMHG / NORMAL <140MMHG)	6,847	1,461	32,088
For cohort DIABETES MELITUS = YA	1,615	1,237	2,110
For cohort DIABETES MELITUS = TIDAK	,236	,062	,900
N of Valid Cases	84		

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
RIWAYAT KELUARGA * DIABETES MELITUS	84	100,0%	0	0,0%	84	100,0%

RIWAYAT KELUARGA * DIABETES MELITUS Crosstabulation

		DIABETES MELITUS		Total
		YA	TIDAK	
RIWAYAT KELUARGA	YA	Count 29	8	37
		% within RIWAYAT KELUARGA 78,4%	21,6%	100,0%
	TIDAK	Count 24	23	47
		% within RIWAYAT KELUARGA 51,1%	48,9%	100,0%
Total	Count 53	31	84	
	% within RIWAYAT KELUARGA 63,1%	36,9%	100,0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	6,633 ^a	1	,010		
Continuity Correction ^b	5,512	1	,019		
Likelihood Ratio	6,851	1	,009		
Fisher's Exact Test				,013	,009
N of Valid Cases	84				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 13,65.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for RIWAYAT KELUARGA (YA / TIDAK)	3,474	1,318	9,159
For cohort DIABETES MELITUS = YA	1,535	1,107	2,129
For cohort DIABETES MELITUS = TIDAK	,442	,224	,872
N of Valid Cases	84		

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
AKTIVITAS FISIK * DIABETES MELITUS	84	100,0%	0	0,0%	84	100,0%

AKTIVITAS FISIK * DIABETES MELITUS Crosstabulation

			DIABETES MELITUS		Total
			YA	TIDAK	
AKTIVITAS FISIK	RINGAN SEDANG	Count	43	29	72
		% within AKTIVITAS FISIK	59,7%	40,3%	100,0%
BERAT		Count	10	2	12
		% within AKTIVITAS FISIK	83,3%	16,7%	100,0%
Total		Count	53	31	84
		% within AKTIVITAS FISIK	63,1%	36,9%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2,463 ^a	1	,117		
Continuity Correction ^b	1,553	1	,213		
Likelihood Ratio	2,732	1	,098		
Fisher's Exact Test				,196	,104
N of Valid Cases	84				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,43.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for AKTIVITAS FISIK (RINGAN SEDANG / BERAT)	,297	,061	1,454
For cohort DIABETES MELITUS = YA	,717	,522	,983
For cohort DIABETES MELITUS = TIDAK	2,417	,661	8,832
N of Valid Cases	84		