

LAMPIRAN 5 : OUTPUT PENGOLAHAN DATA

1. Univariat

a. Usia

usia

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid dewasa	14	43.8	43.8	43.8
Valid lansia	18	56.3	56.3	100.0
Total	32	100.0	100.0	

b. Status Gizi

StatusGizi

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Kurang	5	15.6	15.6	15.6
Valid Lebih	2	6.3	6.3	21.9
Valid Normal	25	78.1	78.1	100.0
Total	32	100.0	100.0	

c. IDWG

IDWG

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Baik	14	43.8	43.8	43.8
Valid Tinggi	18	56.3	56.3	100.0
Total	32	100.0	100.0	

d. Cairan

Cairan

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Baik	20	62.5	62.5	62.5
Valid Lebih	12	37.5	37.5	100.0
Total	32	100.0	100.0	

e. Natrium

Natrium					
	Frequency	Percent	Valid Percent	Cumulative Percent	
Valid	Baik	19	59.4	59.4	59.4
	Lebih	13	40.6	40.6	100.0
	Total	32	100.0	100.0	

f. Kalium

Kalium					
	Frequency	Percent	Valid Percent	Cumulative Percent	
Valid	Baik	25	78.1	78.1	78.1
	Lebih	7	21.9	21.9	100.0
	Total	32	100.0	100.0	

g. Lama Hemodialisa

LamaHemodialisa					
	Frequency	Percent	Valid Percent	Cumulative Percent	
Valid	kurang dari sama dengan 1 tahun	14	43.8	43.8	43.8
	Lebih dari 1 tahun	18	56.3	56.3	100.0
	Total	32	100.0	100.0	

RATA-RATA

Statistics					
		IDWGkg	AsupanCairan	AsupanNatrium	AsupanKalium
N	Valid	32	32	32	32
	Missing	0	0	0	0
Mean		2.5438	743.1844	1430.2188	2280.3521

2. Bivariat

a. Hubungan tingkat kecukupan cairan dan IDWG

Kalium

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Baik	25	78.1	78.1	78.1
Valid Lebih	7	21.9	21.9	100.0
Total	32	100.0	100.0	

Cairan * IDWG Crosstabulation

Count

		IDWG		Total
		Baik	Tinggi	
Cairan	Baik	12	8	20
	Lebih	2	10	12
Total		14	18	32

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5.723 ^a	1	.017		
Continuity Correction ^b	4.097	1	.043		
Likelihood Ratio	6.126	1	.013		
Fisher's Exact Test				.028	.020
N of Valid Cases	32				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5.25.

b. Computed only for a 2x2 table

Mantel-Haenszel Common Odds Ratio Estimate

Estimate		7.500
ln(Estimate)		2.015
Std. Error of ln(Estimate)		.899
Asymp. Sig. (2-sided)		.025
Common Odds Ratio	Lower Bound	1.288
	Upper Bound	43.687
Asymp. 95% Confidence Interval	Lower Bound	.253
	Upper Bound	3.777
	ln(Common Odds Ratio)	

The Mantel-Haenszel common odds ratio estimate is asymptotically normally distributed under the common odds ratio of 1.000 assumption. So is the natural log of the estimate.

b. Hubungan tingkat kecukupan Natrium dan IDWG

Natrium * IDWG Crosstabulation

Count

		IDWG		Total
		Baik	Tinggi	
Natrium	Baik	12	7	19
	Lebih	2	11	13
Total		14	18	32

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	7.158 ^a	1	.007		
Continuity Correction ^b	5.349	1	.021		
Likelihood Ratio	7.690	1	.006		
Fisher's Exact Test				.012	.009
N of Valid Cases	32				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5.69.

b. Computed only for a 2x2 table

Mantel-Haenszel Common Odds Ratio Estimate

Estimate		9.429
ln(Estimate)		2.244
Std. Error of ln(Estimate)		.904
Asymp. Sig. (2-sided)		.013
	Common Odds Ratio	Lower Bound 1.603
		Upper Bound 55.447
Asymp. 95% Confidence Interval	ln(Common Odds Ratio)	Lower Bound .472
		Upper Bound 4.015

The Mantel-Haenszel common odds ratio estimate is asymptotically normally distributed under the common odds ratio of 1.000 assumption. So is the natural log of the estimate.

c. Hubungan tingkat kecukupan kalium dan IDWG

Kalium * IDWG Crosstabulation

Count

		IDWG		Total
		Baik	Tinggi	
Kalium	Baik	13	12	25
	Lebih	1	6	7
Total		14	18	32

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	3.161 ^a	1	.075		
Continuity Correction ^b	1.814	1	.178		
Likelihood Ratio	3.501	1	.061		
Fisher's Exact Test				.104	.087
N of Valid Cases	32				

a. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 3.06.

b. Computed only for a 2x2 table

Mantel-Haenszel Common Odds Ratio Estimate

Estimate	6.500	
ln(Estimate)	1.872	
Std. Error of ln(Estimate)	1.152	
Asymp. Sig. (2-sided)	.104	
	Lower Bound	.680
	Upper Bound	62.149
Asymp. 95% Confidence Interval	Lower Bound	-.386
	Upper Bound	4.130

The Mantel-Haenszel common odds ratio estimate is asymptotically normally distributed under the common odds ratio of 1.000 assumption. So is the natural log of the estimate.

d. Hubungan Lama Hemodialisa dan IDWG

LamaHemodialisa * IDWG Crosstabulation

Count

		IDWG		Total
		Baik	Tinggi	
LamaHemodialisa	kurang dari sama dengan 1 tahun	8	6	14
	Lebih dari 1 tahun	6	12	18
Total		14	18	32

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.814 ^a	1	.178		
Continuity Correction ^b	.976	1	.323		
Likelihood Ratio	1.824	1	.177		
Fisher's Exact Test				.283	.162
N of Valid Cases	32				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 6.13.

b. Computed only for a 2x2 table

Mantel-Haenszel Common Odds Ratio Estimate

Estimate		2.667
ln(Estimate)		.981
Std. Error of ln(Estimate)		.736
Asymp. Sig. (2-sided)		.183
Common Odds Ratio	Lower Bound	.630
	Upper Bound	11.283
Asymp. 95% Confidence Interval	Lower Bound	-.462
	Upper Bound	2.423

The Mantel-Haenszel common odds ratio estimate is asymptotically normally distributed under the common odds ratio of 1.000 assumption. So is the natural log of the estimate.