

OUTPUT SPSS

Diagnosa

	Frequenc y	Percent	Valid Percent	Cumulative Percent
Valid diagnosa TB paru	60	66.7	66.7	66.7
diagnosa tidak TB paru	30	33.3	33.3	100.0
Total	90	100.0	100.0	

umurdikategorikan

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 0-5tahun	40	44.4	44.4	44.4
6-14tahun	50	55.6	55.6	100.0
Total	90	100.0	100.0	

Jeniskelamin

	Frequenc y	Percent	Valid Percent	Cumulative Percent
Valid laki-laki	56	62.2	62.2	62.2
perempuan	34	37.8	37.8	100.0
Total	90	100.0	100.0	

SIB

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid tidak	15	16.7	16.7	16.7
iya	75	83.3	83.3	100.0
Total	90	100.0	100.0	

STATUS Gizi Zscoredikategorikan

	Frequenc y	Percent	Valid Percent	Cumulative Percent
Valid kurus	36	40.0	40.0	40.0
tidak kurus	54	60.0	60.0	60.0
Total	90	100.0	100.0	

Statusekonomi

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid buruk	37	41.1	41.1	41.1
baik	53	58.9	58.9	100.0
Total	90	100.0	100.0	

Riwayat

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid ada	47	52.2	52.2	52.2
tidak	43	47.8	47.8	100.0
Total	90	100.0	100.0	

Kebiasaan

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid ada	78	86.7	86.7	86.7
tidak	12	13.3	13.3	100.0
Total	90	100.0	100.0	

umurdikategorikan * Diagnosa Crosstabulation

		Diagnosa		Total	
		diagnosa TB paru	diagnosa tidak TB paru		
Umurdikategorikan	0-5tahun	Count % within umurdikategorikan	20 50.0%	20 50.0%	40 100.0%
	6-14tahun	Count % within umurdikategorikan	40 80.0%	10 20.0%	50 100.0%
Total		Count % within umurdikategorikan	60 66.7%	30 33.3%	90 100.0%

Chi-Square Tests^c

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	9.000 ^a	1	.003	.004	.003	
Continuity Correction ^b	7.701	1	.006			
Likelihood Ratio	9.081	1	.003	.004	.003	
Fisher's Exact Test				.004	.003	
Linear-by-Linear Association	8.900 ^d	1	.003	.004	.003	.002
N of Valid Cases	90					

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 13.33.

b. Computed only for a 2x2 table

c. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

d. The standardized statistic is -2.983.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for umurdikategorikan (0-5tahun / 6-14tahun)	.250	.099	.633
For cohort Diagnosa = diagnosa TB paru	.625	.445	.878
For cohort Diagnosa = diagnosa tidak TB paru	2.500	1.325	4.718
N of Valid Cases	90		

Jeniskelamin * Diagnosa Crosstabulation

		Diagnosa		Total
		diagnosa TB paru	diagnosa tidak TB paru	
Jeniskelamin laki-laki	Count	41	15	56
	% within Jeniskelamin	73.2%	26.8%	100.0%
perempuan	Count	19	15	34
	% within Jeniskelamin	55.9%	44.1%	100.0%
Total	Count	60	30	90
	% within Jeniskelamin	66.7%	33.3%	100.0%

Chi-Square Tests^c

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2.860 ^a	1	.091	.110	.073	
Continuity Correction ^b	2.133	1	.144			
Likelihood Ratio	2.825	1	.093	.110	.073	
Fisher's Exact Test				.110	.073	
Linear-by-Linear Association	2.828 ^d	1	.093	.110	.073	.045
N of Valid Cases	90					

- a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 11.33.
 b. Computed only for a 2x2 table
 c. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.
 d. The standardized statistic is 1.682.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Jeniskelamin (laki-laki / perempuan)	2.158	.878	5.302
For cohort Diagnosa = diagnosa TB paru	1.310	.934	1.837
For cohort Diagnosa = diagnosa tidak TB paru	.607	.342	1.079
N of Valid Cases	90		

SIB * Diagnosa Crosstabulation

			Diagnosa		Total
			diagnosa TB paru	diagnosa tidak TB paru	
SIB	tidak	Count	12	3	15
		% within SIB	80.0%	20.0%	100.0%
	iya	Count	48	27	75
		% within SIB	64.0%	36.0%	100.0%
Total		Count	60	30	90
		% within SIB	66.7%	33.3%	100.0%

Chi-Square Tests^c

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1.440 ^a	1	.230	.369	.186	
Continuity Correction ^b	.810	1	.368			
Likelihood Ratio	1.548	1	.213	.255	.186	
Fisher's Exact Test				.369	.186	
Linear-by-Linear Association	1.424 ^d	1	.233	.369	.186	.124
N of Valid Cases	90					

- a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 5.00.
- b. Computed only for a 2x2 table
- c. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.
- d. The standardized statistic is 1.193.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for SIB (tidak / iya)	2.250	.583	8.681
For cohort Diagnosa = diagnosa TB paru	1.250	.922	1.695
For cohort Diagnosa = diagnosa tidak TB paru	.556	.193	1.597
N of Valid Cases	90		

Zscoredikategorikan * Diagnosa Crosstabulation

		Diagnosa		Total	
		diagnosa TB paru	diagnosa tidak TB paru		
Zscoredikategorikan	kurus	Count	41	15	56
		% within Zscoredikategorikan	73.2%	26.8%	100.0%
	tidakurus	Count	19	15	34
		% within Zscoredikategorikan	55.9%	44.1%	100.0%
Total		Count	60	30	90
		% within Zscoredikategorikan	66.7%	33.3%	100.0%

Chi-Square Tests^c

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2.860 ^a	1	.091	.110	.073	
Continuity Correction ^b	2.133	1	.144			
Likelihood Ratio	2.825	1	.093	.110	.073	
Fisher's Exact Test				.110	.073	
Linear-by-Linear Association	2.828 ^d	1	.093	.110	.073	.045
N of Valid Cases	90					

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 11.33.

b. Computed only for a 2x2 table

c. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

d. The standardized statistic is 1.682.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Zscoredikategorikan (kurus / tidak kurus)	2.158	.878	5.302
For cohort Diagnosa = diagnosa TB paru	1.310	.934	1.837
For cohort Diagnosa = diagnosa tidak TB paru	.607	.342	1.079
N of Valid Cases	90		

Statusekonomi * Diagnosa Crosstabulation

		Diagnosa		Total
		diagnosa TB paru	diagnosa tidak TB paru	
Statusekonomi	buruk	Count 26 % within Statusekonomi 70.3%	Count 11 29.7%	37 100.0%
	Baik	Count 34 % within Statusekonomi 64.2%	Count 19 35.8%	53 100.0%
Total		Count 60 % within Statusekonomi 66.7%	Count 30 33.3%	90 100.0%

Chi-Square Tests^c

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	.367 ^a	1	.545	.651	.354	
Continuity Correction ^b	.143	1	.705			
Likelihood Ratio	.370	1	.543	.651	.354	
Fisher's Exact Test				.651	.354	
Linear-by-Linear Association	.363 ^d	1	.547	.651	.354	.151
N of Valid Cases	90					

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 12.33.

b. Computed only for a 2x2 table

c. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

d. The standardized statistic is .603.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Statusekonomi (buruk / baik)	1.321	.536	3.253
For cohort Diagnosa = diagnosa TB paru	1.095	.819	1.465
For cohort Diagnosa = diagnosa tidak TB paru	.829	.449	1.530
N of Valid Cases	90		

Riwayat * Diagnosa Crosstabulation

		Diagnosa		Total
		diagnosa TB paru	diagnosa tidak TB paru	
Riwayat ada	Count	45	2	47
	% within Riwayat	95.7%	4.3%	100.0%
tidak	Count	15	28	43
	% within Riwayat	34.9%	65.1%	100.0%
Total	Count	60	30	90
	% within Riwayat	66.7%	33.3%	100.0%

Chi-Square Tests^c

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	37.429 ^a	1	.000	.000	.000	
Continuity Correction ^b	34.741	1	.000			
Likelihood Ratio	42.413	1	.000	.000	.000	
Fisher's Exact Test				.000	.000	
Linear-by-Linear Association	37.014 ^d	1	.000	.000	.000	.000
N of Valid Cases	90					

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 14.33.

b. Computed only for a 2x2 table

c. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

d. The standardized statistic is 6.084.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Riwayat (ada / tidak)	42.000	8.923	197.683
For cohort Diagnosa = diagnosa TB paru	2.745	1.816	4.147
For cohort Diagnosa = diagnosa tidak TB paru	.065	.017	.258
N of Valid Cases	90		

Kebiasaan * Diagnosa Crosstabulation

			Diagnosa		Total
			diagnosa TB paru	diagnosa tidak TB paru	
Kebiasaan	Ada	Count	54	24	78
		% within Kebiasaan	69.2%	30.8%	100.0%
	Tidak	Count	6	6	12
		% within Kebiasaan	50.0%	50.0%	100.0%
Total		Count	60	30	90
		% within Kebiasaan	66.7%	33.3%	100.0%

Chi-Square Tests^c

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	1.731 ^a	1	.188	.324	.162	
Continuity Correction ^b	.974	1	.324			
Likelihood Ratio	1.647	1	.199	.324	.162	
Fisher's Exact Test				.204	.162	
Linear-by-Linear Association	1.712 ^d	1	.191	.324	.162	.109
N of Valid Cases	90					

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 4.00.

b. Computed only for a 2x2 table

c. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

d. The standardized statistic is 1.308.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Kebiasaan (ada / tidak)	2.250	.658	7.694
For cohort Diagnosa = diagnosa TB paru	1.385	.772	2.485
For cohort Diagnosa = diagnosa tidak TB paru	.615	.319	1.186
N of Valid Cases	90		