

Statistics

Perilaku Merokok

N	Valid	178
	Missing	0
Mean		1.47
Median		1.00
Mode		1
Std. Deviation		.500
Minimum		1
Maximum		2

Perilaku Merokok

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid merokok	95	53.4	53.4	53.4
Valid tidak merokok	83	46.6	46.6	100.0
Total	178	100.0	100.0	

UJI NORMALITAS

A. Variabel Pengetahuan

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Pengetahuan	.266	178	.000	.812	178	.000

Distribusi data tidak normal, nilai sig.(nilai p) < 0.05
0.000 < 0.05

a. Lilliefors Significance Correction

Descriptives

		Statistic	Std. Error
Pengetahuan	Mean	18.7079	.09407
	95% Confidence Interval for Mean		
	Lower Bound	18.5222	
	Upper Bound	18.8935	
	5% Trimmed Mean	18.8427	
	Median	19.0000	
	Variance	1.575	
	Std. Deviation	1.25507	
	Minimum	13.00	
	Maximum	20.00	
	Range	7.00	
	Interquartile Range	2.00	
	Skewness	-1.634	.182
Kurtosis	3.858	.362	

Nilai Median 19

B. Variabel Pola Asuh Orang Tua

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
polaasuhortu	.121	178	.000	.957	178	.000

Distribusi data tidak normal, nilai sig.(nilai p) < 0.05
0.000 < 0.05

a. Lilliefors Significance Correction

Descriptives

		Statistic	Std. Error
	Mean	26.2303	.27838
95% Confidence Interval for Mean	Lower Bound	25.6810	
	Upper Bound	26.7797	
	5% Trimmed Mean	26.3308	
	Median	27.0000	
	Variance	13.794	
polaasuhortu	Std. Deviation	3.71404	
	Minimum	17.00	
	Maximum	32.00	
	Range	15.00	
	Interquartile Range	6.00	
	Skewness	-.372	.182
	Kurtosis	-.805	.362

Nilai Median 27

C. Variabel Pengaruh Teman Sebaya

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Pengaruhtemanse baya	.090	178	.001	.970	178	.001

Distribusi data tidak normal, nilai sig.(nilai p) < 0.05
0.001 < 0.05

a. Lilliefors Significance Correction

Descriptives

		Statistic	Std. Error
pengaruhtemansebaya	Mean	22.1966	.24790
	95% Confidence Interval for Mean		
	Lower Bound	21.7074	
	Upper Bound	22.6858	
	5% Trimmed Mean	22.2615	
	Median	22.0000	
	Variance	10.939	
	Std. Deviation	3.30734	
	Minimum	14.00	
	Maximum	28.00	
	Range	14.00	
	Interquartile Range	6.00	
	Skewness	-.183	.182
	Kurtosis	-.801	.362

Nilai Median 22

UNIVARIAT

A. Variabel Pengetahuan

Statistics

pengetahuan dikategorikan

N	Valid	178
	Missing	0
Mean		1.67
Median		2.00
Mode		2
Std. Deviation		.470
Minimum		1
Maximum		2

pengetahuan dikategorikan

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid rendah	58	32.6	32.6	32.6
Valid tinggi	120	67.4	67.4	100.0
Total	178	100.0	100.0	

B. Variabel Pola Asuh Orang Tua

Statistics

pola asuh ortu dikategorikan

N	Valid	178
	Missing	0
Mean		1.53
Median		2.00
Mode		2
Std. Deviation		.500
Minimum		1
Maximum		2

pola asuh ortu dikategorikan

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid beresiko	83	46.6	46.6	46.6
Valid tidak beresiko	95	53.4	53.4	100.0
Total	178	100.0	100.0	

C. Variabel Pengaruh Teman Sebaya

Statistics

pengaruh teman sebaya

dikategorikan

N	Valid	178
	Missing	0
Mean		1.58
Median		2.00
Mode		2
Std. Deviation		.494
Minimum		1
Maximum		2

pengaruh teman sebaya dikategorikan

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid berpengaruh	74	41.6	41.6	41.6
Valid tidak berpengaruh	104	58.4	58.4	100.0
Total	178	100.0	100.0	

BIVARIAT

A. Pengetahuan – Perilaku Merokok

pengetahuan dikategorikan * Perilaku Merokok Crosstabulation

		Perilaku Merokok		Total
		merokok	tidak merokok	
pengetahuan dikategorikan	rendah	Count 37	21	58
		Expected Count 31.0	27.0	58.0
		% within pengetahuan dikategorikan 63.8%	36.2%	100.0%
	tinggi	Count 58	62	120
		Expected Count 64.0	56.0	120.0
		% within pengetahuan dikategorikan 48.3%	51.7%	100.0%
Total		Count 95	83	178
		Expected Count 95.0	83.0	178.0
		% within pengetahuan dikategorikan 53.4%	46.6%	100.0%

Chi-Square Tests

	Value	Df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	3.755 ^a	1	.053	.057	.037
Continuity Correction ^b	3.160	1	.075		
Likelihood Ratio	3.795	1	.051		
Fisher's Exact Test					
Linear-by-Linear Association	3.734	1	.053		
N of Valid Cases	178				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 27.04.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for pengetahuan dikategorikan (rendah / tinggi)	1.883	.989	3.587
For cohort Perilaku Merokok = merokok	1.320	1.010	1.725
For cohort Perilaku Merokok = tidak merokok	.701	.478	1.028
N of Valid Cases	178		

B. Pola Asuh Ortu – Perilaku Merokok

pola asuh ortu dikategorikan * Perilaku Merokok Crosstabulation

		Perilaku Merokok		Total
		merokok	tidak merokok	
pola asuh ortu dikategorikan	Beresiko	Count 31	52	83
		Expected Count 44.3	38.7	83.0
		% within pola asuh ortu dikategorikan 37.3%	62.7%	100.0%
	tidak beresiko	Count 64	31	95
		Expected Count 50.7	44.3	95.0
		% within pola asuh ortu dikategorikan 67.4%	32.6%	100.0%
Total	Count	95	83	178
	Expected Count	95.0	83.0	178.0
	% within pola asuh ortu dikategorikan	53.4%	46.6%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	16.040 ^a	1	.000		
Continuity Correction ^b	14.857	1	.000		
Likelihood Ratio	16.267	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	15.950	1	.000		
N of Valid Cases	178				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 38.70.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for pola asuh ortu dikategorikan (beresiko / tidak beresiko)	.289	.156	.536
For cohort Perilaku Merokok = merokok	.554	.406	.757
For cohort Perilaku Merokok = tidak merokok	1.920	1.376	2.679
N of Valid Cases	178		

C. Pengaruh Teman Sebaya – Perilaku Merokok
 pengaruh teman sebaya dikategorikan * Perilaku Merokok Crosstabulation

		Perilaku Merokok		Total
		merokok	tidak merokok	
berpengaruh	Count	28	46	74
	Expected Count	39.5	34.5	74.0
	% within pengaruh teman sebaya dikategorikan	37.8%	62.2%	100.0%
tidak berpengaruh	Count	67	37	104
	Expected Count	55.5	48.5	104.0
	% within pengaruh teman sebaya dikategorikan	64.4%	35.6%	100.0%
Total	Count	95	83	178
	Expected Count	95.0	83.0	178.0
	% within pengaruh teman sebaya dikategorikan	53.4%	46.6%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	12.279 ^a	1	.000		
Continuity Correction ^b	11.234	1	.001		
Likelihood Ratio	12.391	1	.000		
Fisher's Exact Test				.001	.000
Linear-by-Linear Association	12.210	1	.000		
N of Valid Cases	178				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 34.51.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for pengaruh teman sebaya dikategorikan (berpengaruh / tidak berpengaruh)	.336	.181	.624
For cohort Perilaku Merokok = merokok	.587	.424	.813
For cohort Perilaku Merokok = tidak merokok	1.747	1.277	2.391
N of Valid Cases	178		