

Output  
VARIABEL PERILAKU MEROKOK

**Descriptives**

		Statistic	Std. Error	
Total Perilaku merokok	Mean	.57	.052	
	95% Confidence Interval for Mean	Lower Bound	.47	
		Upper Bound	.67	
	5% Trimmed Mean	.58		
	Median	1.00		
	Variance	.248		
	Std. Deviation	.498		
	Minimum	0		
	Maximum	1		
	Range	1		
	Interquartile Range	1		
	Skewness	-.287	.250	
	Kurtosis	-1.960	.495	

**Tests of Normality**

	Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Total Perilaku merokok	.376	93	.000	.629	93	.000

a. Lilliefors Significance Correction

**Total Perilaku merokok**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	TIDAK MEROKOK	40	36.7	43.0	43.0
	MEROKOK	53	48.6	57.0	100.0
	Total	93	85.3	100.0	
Missing	System	16	14.7		
Total		109	100.0		

## VARIABEL PENGETAHUAN

### Descriptives

		Statistic	Std. Error	
Total Pengetahuan	Mean	5.4301	.12458	
	95% Confidence Interval for Mean	Lower Bound	5.1827	
		Upper Bound	5.6775	
	5% Trimmed Mean	5.3668		
	Median	5.0000		
	Variance	1.443		
	Std. Deviation	1.20143		
	Minimum	4.00		
	Maximum	8.00		
	Range	4.00		
	Interquartile Range	1.50		
	Skewness	.610	.250	
	Kurtosis	-.475	.495	

### Tests of Normality

	Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Total Pengetahuan	.231	93	.000	.880	93	.000

a. Lilliefors Significance Correction

### Total Pengetahuan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	4.00	23	21.1	24.7	24.7
	5.00	32	29.4	34.4	59.1
	6.00	20	18.3	21.5	80.6
	7.00	11	10.1	11.8	92.5
	8.00	7	6.4	7.5	100.0
	Total	93	85.3	100.0	
Missing	System	16	14.7		
Total		109	100.0		

pengetahuan kategorik

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Pengetahuan rendah	70	64.2	75.3	75.3
	Pengetahuan tinggi	23	21.1	24.7	100.0
	Total	93	85.3	100.0	
Missing	System	16	14.7		
Total		109	100.0		

Analisa bivariat pengetahuan dengan perilaku merokok

pengetahuan kategorik \* perilaku kategorik Crosstabulation

			perilaku kategorik		Total
			merokok	tidak merokok	
pengetahuan kategorik	Pengetahuan rendah	Count	39	31	70
		% within pengetahuan kategorik	55.7%	44.3%	100.0%
	Pengetahuan tinggi	Count	14	9	23
		% within pengetahuan kategorik	60.9%	39.1%	100.0%
Total		Count	53	40	93
		% within pengetahuan kategorik	57.0%	43.0%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.188 <sup>a</sup>	1	.665		
Continuity Correction <sup>b</sup>	.036	1	.849		
Likelihood Ratio	.189	1	.664		
Fisher's Exact Test				.809	.427
Linear-by-Linear Association	.186	1	.667		
N of Valid Cases	93				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 9.89.

b. Computed only for a 2x2 table

**Risk Estimate**

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for pengetahuan kategorik (Pengetahuan rendah / Pengetahuan tinggi)	.809	.309	2.114
For cohort perilaku kategorik = merokok	.915	.621	1.350
For cohort perilaku kategorik = tidak merokok	1.132	.638	2.008
N of Valid Cases	93		

**Variabel sikap**

**Descriptives**

		Statistic	Std. Error	
total sikap	Mean	9.3118	.32968	
	95% Confidence Interval for Mean	Lower Bound	8.6570	
		Upper Bound	9.9666	
	5% Trimmed Mean	9.2909		
	Median	10.0000		
	Variance	10.108		
	Std. Deviation	3.17934		
	Minimum	4.00		
	Maximum	15.00		
	Range	11.00		
	Interquartile Range	5.00		
	Skewness	-.024	.250	
	Kurtosis	-1.101	.495	

**sikap kategorik**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Sikap buruk	46	42.2	49.5	49.5
	sikap baik	47	43.1	50.5	100.0
	Total	93	85.3	100.0	
Missing	System	16	14.7		
Total		109	100.0		

**Tests of Normality**

Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
Statistic	df	Sig.	Statistic	df	Sig.

total sikap	.122	93	.002	.951	93	.002
-------------	------	----	------	------	----	------

a. Lilliefors Significance Correction

## Analisa bivariat sikap dengan perilaku merokok

**sikap kategorik \* perilaku kategorik Crosstabulation**

			perilaku kategorik		Total
			merokok	tidak merokok	
sikap kategorik	Sikap buruk	Count	39	7	46
		% within sikap kategorik	84.8%	15.2%	100.0%
	sikap baik	Count	14	33	47
		% within sikap kategorik	29.8%	70.2%	100.0%
Total		Count	53	40	93
		% within sikap kategorik	57.0%	43.0%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	28.685 <sup>a</sup>	1	.000		
Continuity Correction <sup>b</sup>	26.485	1	.000		
Likelihood Ratio	30.617	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	28.377	1	.000		
N of Valid Cases	93				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 19.78.

b. Computed only for a 2x2 table

**Risk Estimate**

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for sikap kategorik (Sikap buruk / sikap baik)	13.133	4.741	36.378
For cohort perilaku kategorik = merokok	2.846	1.805	4.489
For cohort perilaku kategorik = tidak merokok	.217	.107	.440
N of Valid Cases	93		

## Varaibel pengaruh teman sebaya

### Descriptives

		Statistic	Std. Error	
total pengaruh teman sebaya	Mean	1.3763	.11016	
	95% Confidence Interval for Mean	Lower Bound	1.1576	
		Upper Bound	1.5951	
	5% Trimmed Mean	1.3268		
	Median	1.0000		
	Variance	1.129		
	Std. Deviation	1.06234		
	Minimum	.00		
	Maximum	4.00		
	Range	4.00		
	Interquartile Range	1.00		
	Skewness	.471	.250	
	Kurtosis	-.382	.495	

### Tests of Normality

	Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
total pengaruh teman sebaya	.219	93	.000	.891	93	.000

a. Lilliefors Significance Correction

### pengaruh teman sebaya kategorik

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ya mempengaruhi	72	66.1	77.4	77.4
	Tidak mempengaruhi	21	19.3	22.6	100.0
	Total	93	85.3	100.0	
Missing	System	16	14.7		
Total		109	100.0		

## Analisa bivariat pengaruh teman sebaya dengan perilaku merokok

**pengaruh teman sebaya kategorik \* perilaku kategorik Crosstabulation**

			perilaku kategorik		Total
			merokok	tidak merokok	
pengaruh teman sebaya kategorik	Ya mempengaruhi	Count	36	36	72
		% within pengaruh teman sebaya kategorik	50.0%	50.0%	100.0%
	Tidak mempengaruhi	Count	17	4	21
		% within pengaruh teman sebaya kategorik	81.0%	19.0%	100.0%
Total		Count	53	40	93
		% within pengaruh teman sebaya kategorik	57.0%	43.0%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	6.355 <sup>a</sup>	1	.012	.013	.010
Continuity Correction <sup>b</sup>	5.155	1	.023		
Likelihood Ratio	6.839	1	.009		
Fisher's Exact Test					
Linear-by-Linear Association	6.286	1	.012		
N of Valid Cases	93				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 9.03.

b. Computed only for a 2x2 table

**Risk Estimate**

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for pengaruh teman sebaya kategorik (Ya mempengaruhi / Tidak mempengaruhi)	.235	.072	.768
For cohort perilaku kategorik = merokok	.618	.453	.843
For cohort perilaku kategorik = tidak merokok	2.625	1.055	6.531
N of Valid Cases	93		

## Pengaruh iklan rokok

### Descriptives

		Statistic	Std. Error	
total pengaruh iklan rokok	Mean	4.2688	.24469	
	95% Confidence Interval for Mean	Lower Bound	3.7828	
		Upper Bound	4.7548	
	5% Trimmed Mean	4.2987		
	Median	4.0000		
	Variance	5.568		
	Std. Deviation	2.35972		
	Minimum	.00		
	Maximum	8.00		
	Range	8.00		
	Interquartile Range	3.50		
	Skewness	-.198	.250	
	Kurtosis	-.955	.495	

### Tests of Normality

	Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
total pengaruh iklan rokok	.134	93	.000	.947	93	.001

a. Lilliefors Significance Correction

### pengaruh iklan rokok kategorik

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ya berpengaruh	59	54.1	63.4	63.4
	Tidak berpengaruh	34	31.2	36.6	100.0
	Total	93	85.3	100.0	
Missing	System	16	14.7		
Total		109	100.0		

## Analisa bivariat pengaruh iklan rokok dengan perilaku merokok

**pengaruh iklan rokok kategorik \* perilaku kategorik Crosstabulation**

			perilaku kategorik		Total
			merokok	tidak merokok	
pengaruh iklan rokok kategorik	Ya	Count	52	7	59
	berpengaruh	% within pengaruh iklan rokok kategorik	88.1%	11.9%	100.0%
h	Tidak	Count	1	33	34
	berpengaruh	% within pengaruh iklan rokok kategorik	2.9%	97.1%	100.0%
Total		Count	53	40	93
		% within pengaruh iklan rokok kategorik	57.0%	43.0%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	63.870 <sup>a</sup>	1	.000	.000	.000
Continuity Correction <sup>b</sup>	60.442	1	.000		
Likelihood Ratio	75.102	1	.000		
Fisher's Exact Test					
Linear-by-Linear Association	63.184	1	.000		
N of Valid Cases	93				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 14.62.

b. Computed only for a 2x2 table

**Risk Estimate**

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for pengaruh iklan rokok kategorik (Ya berpengaruh / Tidak berpengaruh)	245.143	28.836	2084.025
For cohort perilaku kategorik = merokok	29.966	4.336	207.112

For cohort perilaku kategorik = tidak merokok	.122	.061	.246
N of Valid Cases	93		

## Akses mendapatkan rokok

### Descriptives

		Statistic	Std. Error	
total akses mendapatkan rokok	Mean	4.2581	.08220	
	95% Confidence Interval for Mean	Lower Bound	4.0948	
		Upper Bound	4.4213	
	5% Trimmed Mean	4.3226		
	Median	4.0000		
	Variance	.628		
	Std. Deviation	.79267		
	Minimum	2.00		
	Maximum	5.00		
	Range	3.00		
	Interquartile Range	1.00		
	Skewness	-.900	.250	
	Kurtosis	.372	.495	

### Tests of Normality

	Kolmogorov-Smirnov <sup>a</sup>			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
total akses mendapatkan rokok	.266	93	.000	.790	93	.000

a. Lilliefors Significance Correction

### akses mendapatkan rokok kategorik

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tersedia	79	72.5	84.9	84.9
	Tidak tersedia	14	12.8	15.1	100.0
	Total	93	85.3	100.0	
Missing	System	16	14.7		
Total		109	100.0		

## Analisa bivariat akses mendapatkan rokok dengan perilaku merokok

akses mendapatkan rokok kategorik \* perilaku kategorik Crosstabulation

			perilaku kategorik		Total
			merokok	tidak merokok	
akses mendapatkan rokok kategorik	Tersedia	Count % within akses mendapatkan rokok kategorik	49 62.0%	30 38.0%	79 100.0%
	Tidak tersedia	Count % within akses mendapatkan rokok kategorik	4 28.6%	10 71.4%	14 100.0%
Total		Count % within akses mendapatkan rokok kategorik	53 57.0%	40 43.0%	93 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5.430 <sup>a</sup>	1	.020	.037	.021
Continuity Correction <sup>b</sup>	4.151	1	.042		
Likelihood Ratio	5.448	1	.020		
Fisher's Exact Test					
Linear-by-Linear Association	5.372	1	.020		
N of Valid Cases	93				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 6.02.

b. Computed only for a 2x2 table

Risk Estimate

Value	95% Confidence Interval	
	Lower	Upper

Odds Ratio for akses mendapatkan rokok kategorik (Tersedia / Tidak tersedia)	4.083	1.175	14.186
For cohort perilaku kategorik = merokok	2.171	.932	5.059
For cohort perilaku kategorik = tidak merokok	.532	.344	.821
N of Valid Cases	93		