

**TotalDM**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ya	62	69.7	72.1	72.1
	TidakDM	24	27.0	27.9	100.0
	Total	86	96.6	100.0	
Missing	System	3	3.4		
Total		89	100.0		

**USIA\_KATEGORIK**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	56-65	44	49.4	51.2	51.2
	46-55	42	47.4	48.8	100.0
	Total	86	96,6	100.0	
Missing	System	3	3.4		
Total		89	100.0		

**USIA\_2KATEGORIK**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	46-55	42	47.4	48.8	100.0
	56-65	44	49.4	51.2	51.2
	Total	86	96,6	100,0	
Missing	System	3	3,4		
Total		89	100,0		

**TotalJK**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Perempuan	52	58,4	60,5	60,5
	laki-laki	34	38,2	39,5	100,0
	Total	86	96,6	100,0	
Missing	System	3	3,4		
Total		89	100,0		

**Aktifitas fisik kategorik**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Aktivitas kurang	31	34,8	36,0	36,0
	aktivitas baik	55	61,8	64,0	100,0
	Total	86	96,6	100,0	
Missing	System	3	3,4		
Total		89	100,0		

**Obesitas kategorik**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	ya >30kg	78	87,6	90,7	90,7
	tidak, <30kg	8	9,0	9,3	100,0
	Total	86	96,6	100,0	
Missing	System	3	3,4		
Total		89	100,0		

**Merokok kategorik**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	perokok	65	73,0	75,6	75,6
	tidak merokok	21	23,6	24,4	100,0

Total	86	96,6	100,0
Missing System	3	3,4	
Total	89	100,0	

#### Usia\_Kategorik \* TotalDM Crosstabulation

			TotalDM		Total
			Ya	TidakDM	
Usia_Kategorik	45-55	Count	33	11	44
		% within Usia_Kategorik	75.0%	25.0%	100.0%
	56-65	Count	29	13	42
		% within Usia_Kategorik	69.0%	31.0%	100.0%
Total		Count	62	24	86
		% within Usia_Kategorik	72.1%	27.9%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.378 <sup>a</sup>	1	.538		
Continuity Correction <sup>b</sup>	.140	1	.708		
Likelihood Ratio	.379	1	.538		
Fisher's Exact Test				.633	.354
Linear-by-Linear Association	.374	1	.541		
N of Valid Cases	86				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 11,72.

b. Computed only for a 2x2 table

#### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Usia_Kategorik (,00 / 1,00)	1.345	.523	3.461
For cohort TotalDM = Ya	1.086	.834	1.415
For cohort TotalDM = TidakDM	.808	.408	1.598
N of Valid Cases	86		

**TotalJK \* DM kategorik Crosstabulation**

			DM kategorik		Total
			Ya	TidakDM	
TotalJK	Perempuan	Count	42	10	52
		% within TotalJK	80,8%	19,2%	100,0%
	laki-laki	Count	20	14	34
		% within TotalJK	58,8%	41,2%	100,0%
Total		Count	62	24	86
		% within TotalJK	72,1%	27,9%	100,0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4,921 <sup>a</sup>	1	,027	,048	,025
Continuity Correction <sup>b</sup>	3,891	1	,049		
Likelihood Ratio	4,853	1	,028		
Fisher's Exact Test					
Linear-by-Linear Association	4,864	1	,027		
N of Valid Cases	86				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 9,49.

b. Computed only for a 2x2 table

**Risk Estimate**

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for TotalJK (Perempuan / laki-laki)	2,940	1,114	7,760
For cohort DM kategorik = Ya	1,373	1,006	1,874
For cohort DM kategorik = TidakDM	,467	,235	,928
N of Valid Cases	86		

**Aktifitas fisik kategorik \* DM kategorik Crosstabulation**

			DM kategorik		Total
			Ya	TidakDM	
Aktifitas fisik kategorik	Aktivitas kurang	Count	27	4	31

	% within Aktifitas fisik kategorik	87,1%	12,9%	100,0%
aktivitas baik	Count	35	20	55
	% within Aktifitas fisik kategorik	63,6%	36,4%	100,0%
Total	Count	62	24	86
	% within Aktifitas fisik kategorik	72,1%	27,9%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5,424 <sup>a</sup>	1	,020		
Continuity Correction <sup>b</sup>	4,320	1	,038		
Likelihood Ratio	5,892	1	,015		
Fisher's Exact Test				,025	,016
Linear-by-Linear Association	5,361	1	,021		
N of Valid Cases	86				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 8,65.

b. Computed only for a 2x2 table

#### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Aktifitas fisik kategorik (Aktifitas kurang / aktivitas baik)	3,857	1,179	12,617
For cohort DM kategorik = Ya	1,369	1,075	1,742
For cohort DM kategorik = TidakDM	,355	,133	,945
N of Valid Cases	86		

#### Obesitas kategorik \* DM kategorik Crosstabulation

			DM kategorik		Total
			Ya	TidakDM	
Obesitas kategorik	ya >30kg	Count	60	18	78
		% within Obesitas kategorik	76,9%	23,1%	100,0%

tidak, <30kg	Count	2	6	8
	% within Obesitas kategorik	25,0%	75,0%	100,0%
Total	Count	62	24	86
	% within Obesitas kategorik	72,1%	27,9%	100,0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	9,723 <sup>a</sup>	1	,002		
Continuity Correction <sup>b</sup>	7,313	1	,007		
Likelihood Ratio	8,567	1	,003		
Fisher's Exact Test				,005	,005
Linear-by-Linear Association	9,610	1	,002		
N of Valid Cases	86				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 2,23.

b. Computed only for a 2x2 table

#### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Obesitas kategorik (ya >30kg / tidak, <30kg)	10,000	1,855	53,911
For cohort DM kategorik = Ya	3,077	,921	10,281
For cohort DM kategorik = TidakDM	,308	,174	,544
N of Valid Cases	86		

#### Merokok kategorik \* DM kategorik Crosstabulation

			DM kategorik		Total
			Ya	TidakDM	
Merokok kategorik	perokok	Count	48	17	65
		% within Merokok kategorik	73,8%	26,2%	100,0%
	tidak merokok	Count	14	7	21
		% within Merokok kategorik	66,7%	33,3%	100,0%
Total		Count	62	24	86

% within Merokok kategorik	72,1%	27,9%	100,0%
----------------------------	-------	-------	--------

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,407 <sup>a</sup>	1	,524		
Continuity Correction <sup>b</sup>	,128	1	,720		
Likelihood Ratio	,397	1	,529		
Fisher's Exact Test				,580	,354
Linear-by-Linear Association	,402	1	,526		
N of Valid Cases	86				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5,86.

b. Computed only for a 2x2 table

#### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Merokok kategorik (perokok / tidak merokok)	1,412	,488	4,086
For cohort DM kategorik = Ya	1,108	,792	1,549
For cohort DM kategorik = TidakDM	,785	,378	1,628
N of Valid Cases	86		