

## Lampiran III Hasil Output Spss

### Analisis Univariat

#### Frequencies

##### Kejadian Diare

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Buruk	36	30.8	30.8	30.8
	Baik	81	69.2	69.2	100.0
	Total	117	100.0	100.0	

#### Frequencies

##### Stop BABS

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Buruk	87	74.4	74.4	74.4
	Baik	30	25.6	25.6	100.0
	Total	117	100.0	100.0	

#### Frequencies

##### Cuci Tangan Pakai Sabun

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Buruk	32	27.4	27.4	27.4
	Baik	85	72.6	72.6	100.0
	Total	117	100.0	100.0	

#### Frequencies

##### Pengelolaan Air Minum dan Makanan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Buruk	43	36.8	36.8	36.8
	Baik	74	63.2	63.2	100.0
	Total	117	100.0	100.0	

## Frequencies

### Pengelolaan Sampah

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Buruk	48	41.0	41.0	41.0
	Baik	69	59.0	59.0	100.0
	Total	117	100.0	100.0	

## Frequencies

### Pengelolaan Limbah Cair

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Buruk	39	33.3	33.3	33.3
	Baik	78	66.7	66.7	100.0
	Total	117	100.0	100.0	

## Analisis Bivariat

### Hubungan Stop BABS dengan Kejadian Diare

			D		Total
			diare	tidak diare	
stop babs 11 kategorik	buruk	Count	28	59	87
		Expected Count	26.8	60.2	87.0
		% within stop babs 11 kategorik	32.2%	67.8%	100.0%
	baik	Count	8	22	30
		Expected Count	9.2	20.8	30.0
		% within stop babs 11 kategorik	26.7%	73.3%	100.0%
Total	Count	36	81	117	
	Expected Count	36.0	81.0	117.0	
	% within stop babs 11 kategorik	30.8%	69.2%	100.0%	

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.319 <sup>a</sup>	1	.572		
Continuity Correction <sup>b</sup>	.112	1	.737		
Likelihood Ratio	.325	1	.569		
Fisher's Exact Test				.651	.374
Linear-by-Linear Association	.316	1	.574		
N of Valid Cases	117				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 9.23.

b. Computed only for a 2x2 table

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for stop babs 11 kategorik (buruk / baik)	1.305	.517	3.294
For cohort D = diare	1.207	.619	2.352
For cohort D = tidak diare	.925	.713	1.199
N of Valid Cases	117		

## Crosstabs

### Hubungan Cuci Tangan Pakai Sabun dengan Diare

			D		Total
			diare	tidak diare	
ctps 13 kategorik	buruk	Count	30	2	32
		Expected Count	9.8	22.2	32.0
		% within ctps 13 kategorik	93.8%	6.3%	100.0%
	baik	Count	6	79	85
		Expected Count	26.2	58.8	85.0
		% within ctps 13 kategorik	7.1%	92.9%	100.0%
Total	Count	36	81	117	
	Expected Count	36.0	81.0	117.0	
	% within ctps 13 kategorik	30.8%	69.2%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	82.019 <sup>a</sup>	1	.000		
Continuity Correction <sup>b</sup>	78.000	1	.000		
Likelihood Ratio	86.095	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	81.318	1	.000		
N of Valid Cases	117				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 9.85.

b. Computed only for a 2x2 table

## Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for ctps 13 kategorik (buruk / baik)	197.500	37.756	1033.103
For cohort D= diare	13.281	6.109	28.873
For cohort D = tidak diare	.067	.018	.258
N of Valid Cases	117		

## Crosstabs

### Hubungan Pengelolaan Air Minum dan Makanan dengan Diare

			D		Total
			diare	tidak diare	
pamm rt 1 kategorik	buruk	Count	35	8	43
		Expected Count	13.2	29.8	43.0
		% within pamm rt 1 kategorik	81.4%	18.6%	100.0%
	baik	Count	1	73	74
		Expected Count	22.8	51.2	74.0
		% within pamm rt 1 kategorik	1.4%	98.6%	100.0%
Total	Count	36	81	117	
	Expected Count	36.0	81.0	117.0	
	% within pamm rt 1 kategorik	30.8%	69.2%	100.0%	

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	81.801 <sup>a</sup>	1	.000		
Continuity Correction <sup>b</sup>	78.086	1	.000		
Likelihood Ratio	92.522	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	81.101	1	.000		
N of Valid Cases	117				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 13.23.

b. Computed only for a 2x2 table

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for pamm rt 1 kategorik (buruk / baik)	319.375	38.429	2654.247
For cohort D = diare	60.233	8.553	424.167
For cohort D = tidak diare	.189	.101	.353
N of Valid Cases	117		

## Crosstabs

### Hubungan Pengelolaan Sampah dengan Diare

			D		Total
			diare	tidak diare	
peng sam 20 kategorik	buruk	Count	23	25	48
		Expected Count	14.8	33.2	48.0
		% within peng sam 20 kategorik	47.9%	52.1%	100.0%
	baik	Count	13	56	69
		Expected Count	21.2	47.8	69.0
		% within peng sam 20 kategorik	18.8%	81.2%	100.0%
Total	Count	36	81	117	
	Expected Count	36.0	81.0	117.0	
	% within peng sam 20 kategorik	30.8%	69.2%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	11.235 <sup>a</sup>	1	.001		
Continuity Correction <sup>b</sup>	9.911	1	.002		
Likelihood Ratio	11.197	1	.001		
Fisher's Exact Test				.001	.001
Linear-by-Linear Association	11.139	1	.001		
N of Valid Cases	117				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 14.77.

b. Computed only for a 2x2 table

## Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for peng sam 20 kategorik (buruk / baik)	3.963	1.732	9.066
For cohort D = diare	2.543	1.436	4.505
For cohort D = tidak diare	.642	.478	.861
N of Valid Cases	117		

## Crosstabs

### Hubungan Pengelolaan Limbah Cair dengan Diare

			D		Total
			diare	tidak diare	
peng lim 18 kategorik	buruk	Count	25	14	39
		Expected Count	12.0	27.0	39.0
		% within peng lim 18 kategorik	64.1%	35.9%	100.0%
	baik	Count	11	67	78
		Expected Count	24.0	54.0	78.0
		% within peng lim 18 kategorik	14.1%	85.9%	100.0%
Total	Count	36	81	117	
	Expected Count	36.0	81.0	117.0	
	% within peng lim 18 kategorik	30.8%	69.2%	100.0%	

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	35.389 <sup>a</sup>	1	.000		
Continuity Correction <sup>b</sup>	32.906	1	.000		
Likelihood Ratio	35.045	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	35.086	1	.000		
N of Valid Cases	117				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 12.00.

b. Computed only for a 2x2 table

### Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for peng lim 2 kategorik (buruk / baik)	13.600	5.312	34.821
For cohort D = diare	5.200	2.798	9.664
For cohort D = tidak diare	.382	.243	.601
N of Valid Cases	117		