

Lampiran IV

UJI SPSS

1. Hasil Uji Normalitas *Kolmogorov-Shapiro wilk*

Tests of Normality

imt		Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
tug	underweight < 18,5	.248	9	.116	.913	9	.338
	normal range 18,5-22,9	.283	4	.	.863	4	.272
	overweight 23-29,9	.230	7	.200 [*]	.814	7	.056
	obesitas > 30	.260	4	.	.827	4	.161

a. Lilliefors Significance Correctin

*. This is alower bound of the true significance

2. Hasil Uji *Pearson Correlation*

Correlations

		imt	tug
imt	Pearson Correlation	1	.972 ^{**}
	Sig. (2-tailed)		.000
	N	24	24
tug	Pearson Correlation	.972 ^{**}	1
	Sig. (2-tailed)	.000	
	N	24	24

**.. Correction is significant at the 0.01 level (2-tailed)

3. Hasil Uji *Sperman's rho*

Correlations

		imt	tug
Spearman's rho	imt	Correlation Coefficient	1.000
		Sig. (2-tailed)	.000
		N	24
	tug	Correlation Coefficient	.956 ^{**}
		Sig. (2-tailed)	.000
		N	24

**.. Correlation is significant at the 0.01 level (2-tailed)