

Statistics

ca_mamae

| | | |
|---|---------|-----|
| N | Valid | 120 |
| | Missing | 0 |

ca_mamae

| | Frequen cy | Percent | Valid Percent | Cumulative Percent |
|-----------------------|---------------|---------|------------------|-----------------------|
| Valid kanker payudara | 60 | 50.0 | 50.0 | 50.0 |
| tidak kanker payudara | 60 | 50.0 | 50.0 | 100.0 |
| Total | 120 | 100.0 | 100.0 | |

Statistics

USIA

| | | |
|---|---------|-----|
| N | Valid | 120 |
| | Missing | 0 |

USIA

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|----------------|-----------|---------|---------------|--------------------|
| Valid beresiko | 93 | 77.5 | 77.5 | 77.5 |
| tdk beresiko | 27 | 22.5 | 22.5 | 100.0 |
| Total | 120 | 100.0 | 100.0 | |

Statistics

IMT

| | | |
|---|---------|-----|
| N | Valid | 120 |
| | Missing | 0 |

IMT

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|----------------|-----------|---------|---------------|--------------------|
| Valid OBESITAS | 77 | 64.2 | 64.2 | 64.2 |
| TIDAK OBESITAS | 43 | 35.8 | 35.8 | 100.0 |
| Total | 120 | 100.0 | 100.0 | |

Statistics

imt_univariat

| | | |
|---|---------|-----|
| N | Valid | 120 |
| | Missing | 0 |

imt_univariat

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|----------------------|-----------|---------|---------------|--------------------|
| Valid kurus | 5 | 4.2 | 4.2 | 4.2 |
| normal | 27 | 22.5 | 22.5 | 26.7 |
| berat badan berlebih | 11 | 9.2 | 9.2 | 35.8 |
| obesitas | 77 | 64.2 | 64.2 | 100.0 |
| Total | 120 | 100.0 | 100.0 | |

Statistics

paritas

| | | |
|---|---------|-----|
| N | Valid | 120 |
| | Missing | 0 |

paritas

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|----------------|-----------|---------|---------------|--------------------|
| Valid beresiko | 59 | 49.2 | 49.2 | 49.2 |
| tidak beresiko | 61 | 50.8 | 50.8 | 100.0 |
| Total | 120 | 100.0 | 100.0 | |

Statistics

status_menikah

| | | |
|---|---------|-----|
| N | Valid | 120 |
| | Missing | 0 |

status_menikah

| | Frequency | Percent | Valid Percent | Cumulative Percent |
|---------------------|-----------|---------|---------------|--------------------|
| Valid menikah | 90 | 75.0 | 75.0 | 75.0 |
| Valid belum menikah | 30 | 25.0 | 25.0 | 100.0 |
| Total | 120 | 100.0 | 100.0 | |

Case Processing Summary

| | Cases | | | | | |
|---------------------------|-------|---------|---------|---------|-------|---------|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| USIA * KANKER_PAYUDARA | 120 | 100.0% | 0 | 0.0% | 120 | 100.0% |

USIA * KANKER_PAYUDARA Crosstabulation

| | | KANKER_PAYUDARA | | Total |
|---------------|--------------------------|-----------------|-----------------------|--------|
| | | kanker payudara | tidak kanker payudara | |
| USIA beresiko | Count | 54 | 39 | 93 |
| | Expected Count | 46.5 | 46.5 | 93.0 |
| | % within KANKER_PAYUDARA | 90.0% | 65.0% | 77.5% |
| tdk beresiko | Count | 6 | 21 | 27 |
| | Expected Count | 13.5 | 13.5 | 27.0 |
| | % within KANKER_PAYUDARA | 10.0% | 35.0% | 22.5% |
| Total | Count | 60 | 60 | 120 |
| | Expected Count | 60.0 | 60.0 | 120.0 |
| | % within KANKER_PAYUDARA | 100.0% | 100.0% | 100.0% |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|---------------------|----|-----------------------|----------------------|----------------------|
| Pearson Chi-Square | 10.753 ^a | 1 | .001 | | |
| Continuity Correction ^b | 9.367 | 1 | .002 | | |
| Likelihood Ratio | 11.256 | 1 | .001 | | |
| Fisher's Exact Test | | | | .002 | .001 |
| Linear-by-Linear Association | 10.663 | 1 | .001 | | |

| | | | | | |
|------------------|-----|--|--|--|--|
| N of Valid Cases | 120 | | | | |
|------------------|-----|--|--|--|--|

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 13.50.

b. Computed only for a 2x2 table

Symmetric Measures

| | | Value | Approx. Sig. |
|--------------------|-------------------------|-------|--------------|
| Nominal by Nominal | Contingency Coefficient | .287 | .001 |
| N of Valid Cases | | 120 | |

Risk Estimate

| | Value | 95% Confidence Interval | |
|--|-------|-------------------------|--------|
| | | Lower | Upper |
| Odds Ratio for USIA (beresiko / tdk beresiko) For cohort KANKER_PAYUDARA = kanker payudara | 4.846 | 1.789 | 13.126 |
| For cohort KANKER_PAYUDARA = kanker payudara | 2.613 | 1.264 | 5.403 |
| For cohort KANKER_PAYUDARA = tidak kanker payudara | .539 | .394 | .737 |
| N of Valid Cases | 120 | | |

Case Processing Summary

| | Cases | | | | | |
|--------------------------|-------|---------|---------|---------|-------|---------|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| IMT * KANKER_PAYUDARA | 120 | 100.0% | 0 | 0.0% | 120 | 100.0% |

IMT * KANKER_PAYUDARA Crosstabulation

| | | | KANKER_PAYUDARA | | Total |
|-------------------|----------|-----------------------------|------------------------|-----------------------------|-------|
| | | | KANKER PAYUDAR A | TIDAK KANKER PAYUDARA | |
| IMT | OBESITAS | Count | 45 | 32 | 77 |
| | | Expected Count | 38.5 | 38.5 | 77.0 |
| | | % within KANKER_PAYUDARA | 75.0% | 53.3% | 64.2% |
| TIDAK OBESITAS | | Count | 15 | 28 | 43 |
| | | Expected Count | 21.5 | 21.5 | 43.0 |
| | | % within KANKER_PAYUDARA | 25.0% | 46.7% | 35.8% |
| Total | | Count | 60 | 60 | 120 |
| | | Expected Count | 60.0 | 60.0 | 120.0 |

| | | | |
|-----------------------------|--------|--------|--------|
| % within KANKER_PAYUDARA | 100.0% | 100.0% | 100.0% |
|-----------------------------|--------|--------|--------|

Chi-Square Tests

| | Value | Df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|--------------------|----|--------------------------|-------------------------|-------------------------|
| Pearson Chi-Square | 6.125 ^a | 1 | .013 | | |
| Continuity Correction ^b | 5.219 | 1 | .022 | | |
| Likelihood Ratio | 6.198 | 1 | .013 | | |
| Fisher's Exact Test | | | | .022 | .011 |
| Linear-by-Linear Association | 6.074 | 1 | .014 | | |
| N of Valid Cases | 120 | | | | |

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 21.50.

b. Computed only for a 2x2 table

Symmetric Measures

| | | Value | Approx. Sig. |
|--------------------|-------------------------|-------|--------------|
| Nominal by Nominal | Contingency Coefficient | .220 | .013 |
| N of Valid Cases | | 120 | |

Risk Estimate

| | Value | 95% Confidence Interval | |
|--|-------|-------------------------|-------|
| | | Lower | Upper |
| Odds Ratio for IMT (OBESITAS / TIDAK OBESITAS) For cohort KANKER_PAYUDARA = KANKER PAYUDARA | 2.625 | 1.211 | 5.691 |
| For cohort KANKER_PAYUDARA = TIDAK KANKER PAYUDARA | 1.675 | 1.069 | 2.627 |
| N of Valid Cases | .638 | .453 | .900 |
| | 120 | | |

Case Processing Summary

| | Cases | | | | | |
|------------------------------|-------|---------|---------|---------|-------|---------|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| paritas * kanker_payudara | 120 | 100.0% | 0 | 0.0% | 120 | 100.0% |

paritas * kanker_payudara Crosstabulation

| | | kanker_payudara | | Total |
|------------------|--------------------------|-----------------|---------------------|--------|
| | | kanker payudara | non kanker payudara | |
| paritas beresiko | Count | 27 | 32 | 59 |
| | Expected Count | 29.5 | 29.5 | 59.0 |
| | % within kanker_payudara | 45.0% | 53.3% | 49.2% |
| tidak beresiko | Count | 33 | 28 | 61 |
| | Expected Count | 30.5 | 30.5 | 61.0 |
| | % within kanker_payudara | 55.0% | 46.7% | 50.8% |
| Total | Count | 60 | 60 | 120 |
| | Expected Count | 60.0 | 60.0 | 120.0 |
| | % within kanker_payudara | 100.0% | 100.0% | 100.0% |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|-------------------|----|-----------------------|----------------------|----------------------|
| Pearson Chi-Square | .834 ^a | 1 | .361 | | |
| Continuity Correction ^b | .533 | 1 | .465 | | |
| Likelihood Ratio | .835 | 1 | .361 | | |
| Fisher's Exact Test | | | | .465 | .233 |
| Linear-by-Linear Association | .827 | 1 | .363 | | |

| | | | | |
|------------------|-----|--|--|--|
| N of Valid Cases | 120 | | | |
|------------------|-----|--|--|--|

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 29.50.

b. Computed only for a 2x2 table

Symmetric Measures

| | | Value | Approx. Sig. |
|--------------------|-------------------------|-------|--------------|
| Nominal by Nominal | Contingency Coefficient | .083 | .361 |
| N of Valid Cases | | 120 | |

Risk Estimate

| | Value | 95% Confidence Interval | |
|--|-------|-------------------------|-------|
| | | Lower | Upper |
| Odds Ratio for paritas (beresiko / tidak beresiko) For cohort | .716 | .349 | 1.468 |
| kanker_payudara = kanker payudara For cohort | .846 | .589 | 1.214 |
| kanker_payudara = non kanker payudara For cohort | 1.182 | .825 | 1.693 |
| N of Valid Cases | | 120 | |

Case Processing Summary

| | Cases | | | | | |
|-------------------------------------|-------|---------|---------|---------|-------|---------|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| status_menikah * kanker_payudara | 120 | 100.0% | 0 | 0.0% | 120 | 100.0% |

status_menikah * kanker_payudara Crosstabulation

| | | | kanker_payudara | | Total |
|----------------|---------|--------------------------|-----------------|-----------------------|--------|
| | | | kanker payudara | tidak kanker payudara | |
| status_menikah | menikah | Count | 47 | 43 | 90 |
| | | Expected Count | 45.0 | 45.0 | 90.0 |
| | | % within kanker_payudara | 78.3% | 71.7% | 75.0% |
| belum menikah | menikah | Count | 13 | 17 | 30 |
| | | Expected Count | 15.0 | 15.0 | 30.0 |
| | | % within kanker_payudara | 21.7% | 28.3% | 25.0% |
| Total | | Count | 60 | 60 | 120 |
| | | Expected Count | 60.0 | 60.0 | 120.0 |
| | | % within kanker_payudara | 100.0% | 100.0% | 100.0% |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|-------------------|----|-----------------------|----------------------|----------------------|
| Pearson Chi-Square | .711 ^a | 1 | .399 | | |
| Continuity Correction ^b | .400 | 1 | .527 | | |
| Likelihood Ratio | .713 | 1 | .399 | | |
| Fisher's Exact Test | | | | .528 | .264 |
| Linear-by-Linear Association | .705 | 1 | .401 | | |

| | | | | | |
|------------------|-----|--|--|--|--|
| N of Valid Cases | 120 | | | | |
|------------------|-----|--|--|--|--|

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 15.00.

b. Computed only for a 2x2 table

Symmetric Measures

| | | Value | Approx. Sig. |
|--------------------|-------------------------|-------|--------------|
| Nominal by Nominal | Contingency Coefficient | .077 | .399 |
| N of Valid Cases | | 120 | |

Risk Estimate

| | Value | 95% Confidence Interval | |
|---|-------|-------------------------|-------|
| | | Lower | Upper |
| Odds Ratio for status_menikah (menikah / belum menikah) | 1.429 | .622 | 3.285 |
| For cohort kanker_payudara = kanker payudara | 1.205 | .765 | 1.898 |
| For cohort kanker_payudara = tidak kanker payudara | .843 | .576 | 1.233 |
| N of Valid Cases | | 120 | |