

Statistics

		UMUR	RESPONDEN UMUR	KLASIFIKASI OBESITAS BERDASARKAN IMT	PENDIDIKAN	PENDIDIKAN REponden	PEKERJAAN RESPONDEN	TEKANAN DARAH	RIWAYAT HIPERTENSI	KEBIASAAN ROKOK
N	Valid	290	290	290	290	290	290	290	290	290
	Missing	0	0	0	0	0	0	0	0	0

RESPONDEN UMUR

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid BERSIKO (lebih dari sama dengan 45 tahun)	201	69.3	69.3	69.3
TIDAK BERSIKO (kurang dari 45 tahun)	89	30.7	30.7	100.0
Total	290	100.0	100.0	

Statistics

KATEGORI OBESITAS BERDASARKAN IMT

N	Valid	290
	Missing	0

KATEGORI OBESITAS BERDASARKAN IMT

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid OBESITAS	84	29.0	29.0	29.0
TIDAK OBESITAS	206	71.0	71.0	100.0
Total	290	100.0	100.0	

PENDIDIKAN

	Frequency	Percent	Valid Percent	Cumulative Percent
DIII	11	3.8	3.8	3.8
PT	32	11.0	11.0	14.8
SD	54	18.6	18.6	33.4
Valid SMA	60	20.7	20.7	54.1
SMP	133	45.9	45.9	100.0
Total	290	100.0	100.0	

PENDIDIKAN RESPONDEN

	Frequency	Percent	Valid Percent	Cumulative Percent
PENDIDIKAN DASAR	187	64.5	64.5	64.5
Valid PENDIDIKAN LANJUTAN	103	35.5	35.5	100.0
Total	290	100.0	100.0	

JENIS PEKERJAAN RESPONDEN

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid SWASTA	189	65.2	65.2	65.2
PNS	49	16.9	16.9	82.1
TIDAK BEKERJA	52	17.9	17.9	100.0
Total	290	100.0	100.0	

HIPERTENSI PRIMER

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid HIPERTENSI	195	67.2	67.2	67.2
TIDAK HIPERTENSI	95	32.8	32.8	100.0
Total	290	100.0	100.0	

RIWAYAT HIPERTENSI

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid ADA	186	64.1	64.1	64.1
TIDAK ADA	104	35.9	35.9	100.0
Total	290	100.0	100.0	

Statistics

KEBIASAAN ROKOK

N	Valid	290
	Missing	0

KEBIASAAN ROKOK

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid PEROKOK	16	5.5	5.5	5.5
BUKAN PEROKOK	274	94.5	94.5	100.0
Total	290	100.0	100.0	

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
RESPONDEN UMUR * HIPERTENSI PRIMER	290	100.0%	0	0.0%	290	100.0%
RIWAYAT HIPERTENSI * HIPERTENSI PRIMER	290	100.0%	0	0.0%	290	100.0%
KEBIASAAN ROKOK * HIPERTENSI PRIMER	290	100.0%	0	0.0%	290	100.0%
KLASIFIKASI OBESITAS BERDASARKAN IMT * HIPERTENSI PRIMER	290	100.0%	0	0.0%	290	100.0%
PENDIDIKAN RESPONDEN * HIPERTENSI PRIMER	290	100.0%	0	0.0%	290	100.0%
PEKERJAAN RESPONDEN * HIPERTENSI PRIMER	290	100.0%	0	0.0%	290	100.0%

RESPONDEN UMUR * HIPERTENSI PRIMER

Crosstab

			HIPERTENSI PRIMER		Total
			HIPERTENSI	TIDAK HIPERTENSI	
RESPONDEN UMUR	BERSIKO (lebih dari sama dengan 45 tahun)	Count	145	56	201
		Expected Count	135.2	65.8	201.0
		% within RESPONDEN UMUR	72.1%	27.9%	100.0%
	TIDAK BERSIKO (kurang dari 45 tahun)	Count	50	39	89
		Expected Count	59.8	29.2	89.0
		% within RESPONDEN UMUR	56.2%	43.8%	100.0%
Total	Count	195	95	290	
	Expected Count	195.0	95.0	290.0	
	% within RESPONDEN UMUR	67.2%	32.8%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	7.133 ^a	1	.008		
Continuity Correction ^b	6.427	1	.011		
Likelihood Ratio	6.971	1	.008		
Fisher's Exact Test				.010	.006
Linear-by-Linear Association	7.108	1	.008		
N of Valid Cases	290				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 29.16.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for RESPONDEN UMUR (BERSIKO (lebih dari sama dengan 45 tahun) / TIDAK BERSIKO (kurang dari 45 tahun))	2.020	1.201	3.397
For cohort HIPERTENSI PRIMER = HIPERTENSI	1.284	1.049	1.572
For cohort HIPERTENSI PRIMER = TIDAK HIPERTENSI	.636	.460	.879
N of Valid Cases	290		

RIWAYAT HIPERTENSI KELUARGA * HIPERTENSI PRIMER

Crosstab

		HIPERTENSI PRIMER		Total	
		HIPERTENSI	TIDAK HIPERTENSI		
RIWAYAT HIPERTENSI	ADA	Count	156	30	186
		Expected Count	125.1	60.9	186.0
		% within RIWAYAT HIPERTENSI	83.9%	16.1%	100.0%
	TIDAK ADA	Count	39	65	104
		Expected Count	69.9	34.1	104.0
		% within RIWAYAT HIPERTENSI	37.5%	62.5%	100.0%
Total		Count	195	95	290
		Expected Count	195.0	95.0	290.0
		% within RIWAYAT HIPERTENSI	67.2%	32.8%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	65.115 ^a	1	.000		
Continuity Correction ^b	63.026	1	.000		
Likelihood Ratio	64.868	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	64.890	1	.000		
N of Valid Cases	290				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 34.07.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for RIWAYAT HIPERTENSI (ADA / TIDAK ADA)	8.667	4.965	15.127
For cohort HIPERTENSI PRIMER = HIPERTENSI	2.237	1.731	2.889
For cohort HIPERTENSI PRIMER = TIDAK HIPERTENSI	.258	.180	.370
N of Valid Cases	290		

KEBIASAAN ROKOK * HIPERTENSI PRIMER

Crosstab

			HIPERTENSI PRIMER		Total
			HIPERTENSI	TIDAK HIPERTENSI	
KEBIASAAN ROKOK	PEROKOK	Count	7	9	16
		Expected Count	10.8	5.2	16.0
		% within KEBIASAAN ROKOK	43.8%	56.2%	100.0%
BUKAN PEROKOK		Count	188	86	274
		Expected Count	184.2	89.8	274.0
		% within KEBIASAAN ROKOK	68.6%	31.4%	100.0%
Total		Count	195	95	290
		Expected Count	195.0	95.0	290.0
		% within KEBIASAAN ROKOK	67.2%	32.8%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.243 ^a	1	.039		
Continuity Correction ^b	3.189	1	.074		
Likelihood Ratio	3.950	1	.047		
Fisher's Exact Test				.054	.040
Linear-by-Linear Association	4.228	1	.040		
N of Valid Cases	290				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 5.24.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for KEBIASAAN ROKOK (PEROKOK / BUKAN PEROKOK)	.356	.128	.987
For cohort HIPERTENSI PRIMER = HIPERTENSI	.638	.364	1.118
For cohort HIPERTENSI PRIMER = TIDAK HIPERTENSI	1.792	1.124	2.857
N of Valid Cases	290		

KLASIFIKASI OBESITAS BERDASARKAN IMT * HIPERTENSI PRIMER

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
KATEGORI OBESITAS BERDASARKAN IMT * HIPERTENSI PRIMER	290	100.0%	0	0.0%	290	100.0%

KATEGORI OBESITAS BERDASARKAN IMT * HIPERTENSI PRIMER Crosstabulation

			HIPERTENSI PRIMER		Total
			HIPERTENSI	TIDAK HIPERTENSI	
KATEGORI OBESITAS BERDASARKAN IMT	OBESITAS	Count	54	30	84
		Expected Count	56.5	27.5	84.0
		% within KATEGORI OBESITAS BERDASARKAN IMT	64.3%	35.7%	100.0%
	TIDAK OBESITAS	Count	141	65	206
		Expected Count	138.5	67.5	206.0
		% within KATEGORI OBESITAS BERDASARKAN IMT	68.4%	31.6%	100.0%
Total		Count	195	95	290
		Expected Count	195.0	95.0	290.0
		% within KATEGORI OBESITAS BERDASARKAN IMT	67.2%	32.8%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.469 ^a	1	.493		
Continuity Correction ^b	.299	1	.584		
Likelihood Ratio	.465	1	.495		
Fisher's Exact Test				.494	.291
Linear-by-Linear Association	.467	1	.494		
N of Valid Cases	290				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 27.52.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for KATEGORI OBESITAS BERDASARKAN IMT (OBESITAS / TIDAK OBESITAS)	.830	.486	1.416
For cohort HIPERTENSI PRIMER = HIPERTENSI	.939	.781	1.129
For cohort HIPERTENSI PRIMER = TIDAK HIPERTENSI	1.132	.797	1.607
N of Valid Cases	290		

PENDIDIKAN RESPONDEN * HIPERTENSI PRIMER

Crosstab

		HIPERTENSI PRIMER		Total	
		HIPERTENSI	TIDAK HIPERTENSI		
PENDIDIKAN RESPONDEN	PENDIDIKAN DASAR	Count	142	45	187
		Expected Count	125.7	61.3	187.0
		% within PENDIDIKAN RESPONDEN	75.9%	24.1%	100.0%
PENDIDIKAN RESPONDEN	PENDIDIKAN LANJUTAN	Count	53	50	103
		Expected Count	69.3	33.7	103.0
		% within PENDIDIKAN RESPONDEN	51.5%	48.5%	100.0%
Total		Count	195	95	290
		Expected Count	195.0	95.0	290.0
		% within PENDIDIKAN REponden	67.2%	32.8%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	18.069 ^a	1	.000		
Continuity Correction ^b	16.974	1	.000		
Likelihood Ratio	17.743	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	18.006	1	.000		
N of Valid Cases	290				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 33.74.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for PENDIDIKAN RESPONDEN (PENDIDIKAN DASAR / PENDIDIKAN LANJUTAN)	2.977	1.785	4.965
For cohort HIPERTENSI PRIMER = HIPERTENSI	1.476	1.203	1.810
For cohort HIPERTENSI PRIMER = TIDAK HIPERTENSI	.496	.359	.685
N of Valid Cases	290		

PEKERJAAN RESPONDEN * HIPERTENSI PRIMER

Crosstab

			HIPERTENSI PRIMER		Total
			HIPERTENSI	TIDAK HIPERTENSI	
PEKERJAAN RESPONDEN	SWASTA	Count	136	53	189
		Expected Count	127.1	61.9	189.0
		% within PEKERJAAN RESPONDEN	72.0%	28.0%	100.0%
	PNS	Count	26	23	49
		Expected Count	32.9	16.1	49.0
		% within PEKERJAAN RESPONDEN	53.1%	46.9%	100.0%
	TIDAK BEKERJA	Count	33	19	52
		Expected Count	35.0	17.0	52.0
		% within PEKERJAAN RESPONDEN	63.5%	36.5%	100.0%
Total	Count	195	95	290	
	Expected Count	195.0	95.0	290.0	
	% within PEKERJAAN RESPONDEN	67.2%	32.8%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.719 ^a	2	.035
Likelihood Ratio	6.521	2	.038
Linear-by-Linear Association	3.038	1	.081
N of Valid Cases	290		

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 16.05.

Risk Estimate

	Value
Odds Ratio for PEKERJAAN RESPONDEN (SWASTA / PNS)	^a

a. Risk Estimate statistics cannot be computed. They are only computed for a 2*2 table without empty cells.

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
KATEGORI OBESITAS BERDASARKAN IMT * HIPERTENSI PRIMER * RIWAYAT HIPERTENSI	290	100.0%	0	0.0%	290	100.0%

KATEGORI OBESITAS BERDASARKAN IMT * HIPERTENSI PRIMER * RIWAYAT HIPERTENSI Crosstabulation

RIWAYAT HIPERTENSI		HIPERTENSI PRIMER		Total	
		HIPERTENSI	TIDAK HIPERTENSI		
ADA	KATEGORI OBESITAS BERDASARKAN IMT	Count	37	7	44
		Expected Count	36.9	7.1	44.0
		% within KATEGORI OBESITAS BERDASARKAN IMT	84.1%	15.9%	100.0%
TIDAK OBESITAS		Count	119	23	142
		Expected Count	119.1	22.9	142.0

		% within KATEGORI			
		OBESITAS BERDASARKAN	83.8%	16.2%	100.0%
		IMT			
		Count	156	30	186
		Expected Count	156.0	30.0	186.0
Total		% within KATEGORI			
		OBESITAS BERDASARKAN	83.9%	16.1%	100.0%
		IMT			
		Count	17	23	40
		Expected Count	15.0	25.0	40.0
	OBESITAS	% within KATEGORI			
		OBESITAS BERDASARKAN	42.5%	57.5%	100.0%
		IMT			
		Count	22	42	64
		Expected Count	24.0	40.0	64.0
	TIDAK ADA	% within KATEGORI			
		OBESITAS BERDASARKAN	34.4%	65.6%	100.0%
		IMT			
		Count	39	65	104
		Expected Count	39.0	65.0	104.0
	TIDAK OBESITAS	% within KATEGORI			
		OBESITAS BERDASARKAN	37.5%	62.5%	100.0%
		IMT			
		Count	54	30	84
		Expected Count	56.5	27.5	84.0
Total	KATEGORI OBESITAS BERDASARKAN IMT	OBESITAS			

Total	TIDAK OBESITAS	% within KATEGORI OBESITAS BERDASARKAN IMT	64.3%	35.7%	100.0%
		Count	141	65	206
		Expected Count	138.5	67.5	206.0
		% within KATEGORI OBESITAS BERDASARKAN IMT	68.4%	31.6%	100.0%
		Count	195	95	290
	Expected Count	195.0	95.0	290.0	
	OBESITAS	% within KATEGORI OBESITAS BERDASARKAN IMT	67.2%	32.8%	100.0%
		Count	141	65	206
		Expected Count	138.5	67.5	206.0
		% within KATEGORI OBESITAS BERDASARKAN IMT	68.4%	31.6%	100.0%
Count		195	95	290	

Chi-Square Tests

RIWAYAT HIPERTENSI	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.002 ^c	1	.964		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.002	1	.964		
Fisher's Exact Test				1.000	.586
Linear-by-Linear Association	.002	1	.964		
N of Valid Cases	186				

TIDAK ADA	Pearson Chi-Square	.693 ^d	1	.405	
	Continuity Correction ^b	.390	1	.532	
	Likelihood Ratio	.690	1	.406	
	Fisher's Exact Test			.414	.265
	Linear-by-Linear Association	.687	1	.407	
	N of Valid Cases	104			
Total	Pearson Chi-Square	.469 ^a	1	.493	
	Continuity Correction ^b	.299	1	.584	
	Likelihood Ratio	.465	1	.495	
	Fisher's Exact Test			.494	.291
	Linear-by-Linear Association	.467	1	.494	
	N of Valid Cases	290			

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 27.52.

b. Computed only for a 2x2 table

c. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 7.10.

d. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 15.00.

Risk Estimate

RIWAYAT HIPERTENSI	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for KATEGORI OBESITAS BERDASARKAN IMT (OBESITAS / TIDAK OBESITAS)	1.022	.406	2.571
For cohort HIPERTENSI PRIMER = HIPERTENSI	1.003	.866	1.163
For cohort HIPERTENSI PRIMER = TIDAK HIPERTENSI	.982	.452	2.133
N of Valid Cases	186		
Odds Ratio for KATEGORI OBESITAS BERDASARKAN IMT (OBESITAS / TIDAK OBESITAS)	1.411	.627	3.178
For cohort HIPERTENSI PRIMER = HIPERTENSI	1.236	.754	2.027
For cohort HIPERTENSI PRIMER = TIDAK HIPERTENSI	.876	.636	1.207
N of Valid Cases	104		
Odds Ratio for KATEGORI OBESITAS BERDASARKAN IMT (OBESITAS / TIDAK OBESITAS)	.830	.486	1.416

ADA

TIDAK ADA

Total

For cohort HIPERTENSI PRIMER = HIPERTENSI	.939	.781	1.129
For cohort HIPERTENSI PRIMER = TIDAK HIPERTENSI	1.132	.797	1.607
N of Valid Cases	290		