

USIA RESPONDEN

Crosstab

		Pernah LBP tidaknya		Total
		Pernah	Tidak Pernah	
Usia Responden	Beresiko Jika Usia lebih 30 Tahun	Count 19	15	34
		% within Usia Responden 55.9%	44.1%	100.0%
Usia Responden	Tidak Beresiko Jika Usia Dibawah 30 Tahun	Count 7	9	16
		% within Usia Responden 43.8%	56.3%	100.0%
Total		Count 26	24	50
		% within Usia Responden 52.0%	48.0%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	.642 ^a	1	.423		
Continuity Correction ^b	.248	1	.019		
Likelihood Ratio	.642	1	.423		
Fisher's Exact Test				.547	.309
Linear-by-Linear Association	.629	1	.428		
N of Valid Cases	50				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 7,68.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Usia Responden (Beresiko Jika Usia lebih 30 Tahun / Tidak Beresiko Jika Usia Dibawah 30 Tahun)	1.629	.492	5.393
For cohort Pernah LBP tidaknya = Pernah	1.277	.680	2.400
For cohort Pernah LBP tidaknya = Tidak Pernah	.784	.442	1.393
N of Valid Cases	50		

IMT (STATUS GIZI)

Crosstab

		Pernah LBP tidaknya		Total	
		Pernah	Tidak Pernah		
Status Gizi Responden	Beresiko Jika IMT lebih 25	Count	15	9	24
		% within Status Gizi Responden	62.5%	37.5%	100.0%
	Tidak Beresiko Jika IMT Kurang 25	Count	11	15	26
		% within Status Gizi Responden	42.3%	57.7%	100.0%
Total		Count	26	24	50
		% within Status Gizi Responden	52.0%	48.0%	100.0%

Chi-Square Tests

	Value	Df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.039 ^a	1	.153		
Continuity Correction ^b	1.310	1	.025		
Likelihood Ratio	2.054	1	.152		
Fisher's Exact Test				.171	.126
Linear-by-Linear Association	1.998	1	.158		
N of Valid Cases	50				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 11,52.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Status Gizi Responden (Beresiko Jika IMT lebih 25 / Tidak Beresiko Jika IMT Kurang 25)	2.273	.730	7.071
For cohort Pernah LBP tidaknya = Pernah	1.477	.856	2.549
For cohort Pernah LBP tidaknya = Tidak Pernah	.650	.352	1.199
N of Valid Cases	50		

POSISI KERJA

Crosstab

		Pernah LBP tidaknya		Total	
		Pernah	Tidak Pernah		
Nordic Body Map	Beresiko Jika 8&9	Count	24	18	42
		% within Nordic Body Map	57.1%	42.9%	100.0%
	Tidak Beresiko Jika diluar 8&9	Count	2	6	8
		% within Nordic Body Map	25.0%	75.0%	100.0%
Total	Count	26	24	50	
	% within Nordic Body Map	52.0%	48.0%	100.0%	

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.782 ^a	1	.095		
Continuity Correction ^b	1.643	1	.020		
Likelihood Ratio	2.873	1	.090		
Fisher's Exact Test				.132	.100
Linear-by-Linear Association	2.726	1	.099		
N of Valid Cases	50				

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 3,84.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Nordic Body Map (Beresiko Jika 8&9 / Tidak Beresiko Jika diluar 8&9)	4.000	.721	22.183
For cohort Pernah LBP tidaknya = Pernah	2.286	.669	7.808
For cohort Pernah LBP tidaknya = Tidak Pernah	.571	.336	.972
N of Valid Cases	50		

MASA KERJA

Crosstab

			Pernah LBP tidaknya		Total
			Pernah	Tidak Pernah	
Masa Kerja Responden	Beresiko	Count	32	5	37
		% within Masa Kerja	86.5%	13.5%	100.0%
	Tidak Beresiko	Count	10	3	13
		% within Masa Kerja	76.9%	23.1%	100.0%
Total		Count	42	8	50
		% within Masa Kerja	84.0%	16.0%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	.655 ^a	1	.418		
Continuity Correction ^b	.136	1	.012		
Likelihood Ratio	.615	1	.433		
Fisher's Exact Test				.413	.341
Linear-by-Linear Association	.642	1	.423		
N of Valid Cases	50				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 2,08.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Masa Kerja (Beresiko / Tidak Beresiko)	1.920	.388	9.489
For cohort Nordic Body Map = Beresiko Jika 8&9	1.124	.813	1.554
For cohort Nordic Body Map = Tidak Beresiko Jika diluar 8&9	.586	.162	2.115
N of Valid Cases	50		