

DAFTAR PUSTAKA

- Ahmad Bhat Scholar, S. *et al.* (2017) ‘Development of norms of maximal oxygen uptake (VO₂ max.) as an indicator of aerobic fitness of high altitude male youth of Kashmir’, ~ 1037 ~ *International Journal of Physiology*, 2(2), pp. 1037–1040.
- Allen, P. *et al.* (2019) ‘Emerging temporal lobe dysfunction in people at clinical high risk for psychosis’, *Frontiers in Psychiatry*, 10(MAY), pp. 1–12.
- Arfa, K. J. (2017) ‘EXERCISE AND CLINICAL PRACTICE : INTEGRATION ISSUES AND KNOWLEDGE IN MENTAL HEALTH PROFESSIONALS Dissertation presented to the Faculty of the California School of Professional Psychology Alliant International University Los Angeles In partial fulfillment ’.
- Arijaya, D. N. K. (2013) ‘Clozapine Pada Skizofrenia Paranoid Dengan Obesitas’, 53(9), pp. 1–8.
- Artanty, A. and Lufthansa, L. (2017) ‘Pengaruh latihan lari 15 menit terhadap kemampuan vo 2 max’, *Jendela Olahraga*, 2(2), pp. 9–19.
- Ashok, A. H., Baugh, J. and Yeragani, V. K. (2012) ‘Paul Eugen Bleuler and the origin of the term schizophrenia (SCHIZOPRENIEGRUPPE)’, *Indian Journal of Psychiatry*, 54(1), pp. 95–96.
- Azad, M. C. *et al.* (2016) ‘Cardiovascular diseases among patients with schizophrenia’, *Asian Journal of Psychiatry*.
- Barajas, A. *et al.* (2015) ‘Gender differences in individuals at high-risk of psychosis: A comprehensive literature review’, *Scientific World Journal*, 2015.
- Browne, J. *et al.* (2016) ‘Work out by walking: A pilot exercise program for individuals with schizophrenia spectrum disorders’, *Journal of Nervous and Mental Disease*, 204(9), pp. 651–657.
- Cameron, I. M. *et al.* (2017) ‘Obesity in individuals with schizophrenia: a case controlled study in Scotland’, *BJPsych Open*, 3(5), pp. 254–256.
- Coxon, J. P. *et al.* (2018) ‘GABA concentration in sensorimotor cortex following high-intensity exercise and relationship to lactate levels’, *Journal of*

- Physiology*, 596(4), pp. 691–702.
- Curcic, D. *et al.* (2017a) ‘Positive impact of prescribed physical activity on symptoms of schizophrenia: Randomized clinical trial’, *Psychiatria Danubina*, 29(4), pp. 459–465.
- Curcic, D. *et al.* (2017b) ‘Positive impact of prescribed physical activity on symptoms of schizophrenia: Randomized clinical trial’, *Psychiatria Danubina*, 29(4), pp. 459–465.
- Dasar, R. K. (2013) ‘Penyajian Pokok-Pokok Hasil Riset Kesehatan Dasar 2013’.
- Engh, J. A. *et al.* (2015) ‘Effects of high-intensity aerobic exercise on psychotic symptoms and neurocognition in outpatients with schizophrenia: Study protocol for a randomized controlled trial’, *Trials*.
- ‘ExRx’ (no date).
- Firth, J. *et al.* (2015) ‘A systematic review and meta-Analysis of exercise interventions in schizophrenia patients’, *Psychological Medicine*, 45(7), pp. 1343–1361.
- Firth, J. *et al.* (2017) ‘Aerobic exercise improves cognitive functioning in people with schizophrenia: A systematic review and meta-analysis’, *Schizophrenia Bulletin*, 43(3), pp. 546–556.
- Franco, V. C., Zortéa, K. and de Abreu, P. S. B. (2016) ‘Obesity and Clozapine use in Schizophrenia’, *Obesity Research - Open Journal*, 3(2), pp. 24–29.
- Franke, B. *et al.* (2016) ‘Genetic influences on schizophrenia and subcortical brain volumes: Large-scale proof of concept’, *Nature Neuroscience*.
- Franklin, B. A. and Billecke, S. (2012) ‘Putting the benefits and risks of aerobic exercise in perspective’, *Current Sports Medicine Reports*, 11(4), pp. 201–208.
- ‘Schizophrenia, Not a Psychotic Disorder_ Bleuler Revisited’ (no date).
- Kahn, R. S. *et al.* (2015) ‘Schizophrenia’, *Nature Reviews Disease Primers*, 1(1), p. 15067.
- KEMENKES (2018) *KEMENKES (2018) kementrian kesehatan 2016*. JAKARTA.
Available at: http://www.depkes.go.id/resources/download/infoterkini/materi_rakorpop_2018/Hasil_Riskesdas_2018.pdf.kementrian kesehatan 2016. JAKARTA. Available at:

- http://www.depkes.go.id/resources/download/info-terkini/materi_rakorpop_2018/Hasil_Riskesdas_2018.pdf.
- kemenkes 2013 kementerian kesehatan republik indonesia (2013) *No Title RISET KESEHATAN DASAR (RISKESDA)*. JAKARTA.
- Kimhy, D. et al. (2014) ‘Aerobic fitness and body mass index in individuals with schizophrenia: Implications for neurocognition and daily functioning’, *Psychiatry Research*. Elsevier, 220(3), pp. 784–79.
- Krogh, J. et al. (2014) ‘Can Exercise Increase Fitness and Reduce Weight in Patients with Schizophrenia and Depression?’, *Frontiers in Psychiatry*, 5(July), pp. 1–7.
- Lee, S. K., Lee, C. M. and Park, J. H. (2015) ‘Effects of combined exercise on physical fitness and neurotransmitters in children with ADHD: A pilot randomized controlled study’, *Journal of Physical Therapy Science*, 27(9), pp. 2915–2919.
- Manzella, F. (2015) ‘Smoking in schizophrenic patients: A critique of the self-medication hypothesis’, *World Journal of Psychiatry*, 5(1), p. 35.
- Maulana, I. et al. (2019) ‘Penyuluhan Kesehatan Jiwa untuk Meningkatkan Pengetahuan Masyarakat tentang Masalah Kesehatan Jiwa di Lingkungan Sekitarnya’, *Media Karya Kesehatan*, 2(2), pp. 218–225.
- Mousavi Gilani, S. R. and Khazaei Feizabad, A. (2019) ‘The effects of aerobic exercise training on mental health and self-esteem of type 2 diabetes mellitus patients’, *Health Psychology Research*, 7(1), pp. 10–14.
- Mubarik, A. and Tohid, H. (2016) ‘Frontal lobe alterations in schizophrenia: A review’, *Trends in Psychiatry and Psychotherapy*, 38(4), pp. 198–206.
- Perez-Cruzado, D. et al. (2017) ‘Physical fitness and levels of physical activity in people with severe mental illness: A cross-sectional study’, *BMC Sports Science, Medicine and Rehabilitation*. BMC Sports Science, Medicine and Rehabilitation, 9(1), pp. 1–6.
- Scheewe, T. W. et al. (2013) ‘Exercise therapy improves mental and physical health in schizophrenia: A randomised controlled trial’, *Acta Psychiatrica Scandinavica*, 127(6), pp. 464–473.
- Sharkey, J. L. (2016) ‘Exercise and mental health—Implications for treatment: A

- review of the literature', *Counseling and Wellness*, 5, pp. 1–8.
- Sommer, I. E. et al. (2015) 'Schizophrenia, PRIMER', (November).
- Son, S., Jeon, B. and Kim, H. (2016) 'Effects of a walking exercise program for obese individuals with intellectual disability staying in a residential care facility', *Journal of Physical Therapy Science*, 28(3), pp. 788–793.
- Soundy, A. et al. (2014) 'The benefits of walking for individuals with schizophrenia spectrum disorders: A systematic review', *International Journal of Therapy and Rehabilitation*, 21(9), pp. 410–420.
- Stubbs, B. et al. (2018) 'EPA guidance on physical activity as a treatment for severe mental illness: a meta-review of the evidence and Position Statement from the European Psychiatric Association (EPA), supported by the International Organization of Physical Therapists in Mental ', *European Psychiatry*, 54, pp. 124–144.
- Tripathi, A., Kar, S. K. and Shukla, R. (2018) 'Cognitive deficits in schizophrenia: Understanding the biological correlates and remediation strategies', *Clinical Psychopharmacology and Neuroscience*, 16(1), pp. 7–17.
- Vancampfort, D. et al. (2015) 'Aerobic capacity is associated with global functioning in people with schizophrenia', *Journal of Mental Health*. Shadowfax Publishing and Informa UK Limited, 24(4), pp. 214–218.
- Vendelboe, T. V. et al. (2016) 'The crystal structure of human dopamine β -hydroxylase at 2.9 Å resolution', *Science Advances*, 2(4), pp. 1–11.
- Wang, P. W. et al. (2018) 'Effect of aerobic exercise on improving symptoms of individuals with schizophrenia: A single blinded randomized control study', *Frontiers in Psychiatry*, 9(MAY), pp. 1–7.
- Xu, Y. M. et al. (2014) 'Prevalence and correlates of cigarette smoking among Chinese schizophrenia inpatients receiving antipsychotic mono-therapy', *PLoS ONE*, 9(2)..
- Yeh, H. P. et al. (2017) 'Physical and emotional benefits of different exercise environments designed for treadmill running', *International Journal of Environmental Research and Public Health*, 14(7).