

LAMPIRAN

Univariat

Hipertensi

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	hipertensi	110	65.9	65.9	65.9
	normal	57	34.1	34.1	100.0
	Total	167	100.0	100.0	

Usia

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	berisiko	99	59.3	59.3	59.3
	tidak berisiko	68	40.7	40.7	100.0
	Total	167	100.0	100.0	

jenis_kelamin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	laki-laki	102	61.1	61.1	61.1
	perempuan	65	38.9	38.9	100.0
	Total	167	100.0	100.0	

Stress

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	stress	125	74.9	74.9	74.9
	tidak stress	42	25.1	25.1	100.0
	Total	167	100.0	100.0	

kebiasaan_merokok

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid merokok	98	58.7	58.7	58.7
tidak merokok	69	41.3	41.3	100.0
Total	167	100.0	100.0	

Bivariat

usia * hipertensi Crosstabulation

			hipertensi		Total
			hipertensi	normal	
Usia	berisiko	Count	90	9	99
		% within usia	90.9%	9.1%	100.0%
	tidak berisiko	Count	20	48	68
		% within usia	29.4%	70.6%	100.0%
Total		Count	110	57	167
		% within usia	65.9%	34.1%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	67.812 ^a	1	.000		
Continuity Correction ^b	65.104	1	.000		
Likelihood Ratio	71.690	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	67.406	1	.000		
N of Valid Cases	167				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 23.21.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for usia (berisiko / tidak berisiko)	24.000	10.144	56.782
For cohort hipertensi = hipertensi	3.091	2.128	4.490
For cohort hipertensi = normal	.129	.068	.245
N of Valid Cases	167		

jenis_kelamin * hipertensi Crosstabulation

			hipertensi		Total
			hipertensi	normal	
jenis_kelamin	laki-laki	Count	85	17	102
		% within jenis_kelamin	83.3%	16.7%	100.0%
	perempuan	Count	25	40	65
		% within jenis_kelamin	38.5%	61.5%	100.0%
Total		Count	110	57	167
		% within jenis_kelamin	65.9%	34.1%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2- sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	35.556 ^a	1	.000		
Continuity Correction ^b	33.588	1	.000		
Likelihood Ratio	35.866	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	35.343	1	.000		
N of Valid Cases	167				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 22.19.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for jenis_kelamin (laki-laki / perempuan)	8.000	3.887	16.463
For cohort hipertensi = hipertensi	2.167	1.574	2.982
For cohort hipertensi = normal	.271	.168	.435
N of Valid Cases	167		

stress * hipertensi Crosstabulation

			hipertensi		Total
			hipertensi	normal	
Stress	stress	Count	99	26	125
		% within stress	79.2%	20.8%	100.0%
	tidak stress	Count	11	31	42
		% within stress	26.2%	73.8%	100.0%
Total		Count	110	57	167
		% within stress	65.9%	34.1%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	39.293 ^a	1	.000		
Continuity Correction ^b	36.971	1	.000		
Likelihood Ratio	38.269	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	39.058	1	.000		
N of Valid Cases	167				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 14.34.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for stress (stress / tidak stress)	10.731	4.763	24.175
For cohort hipertensi = hipertensi	3.024	1.806	5.064
For cohort hipertensi = normal	.282	.191	.415
N of Valid Cases	167		

kebiasaan_merokok * hipertensi Crosstabulation

			hipertensi		Total
			hipertensi	normal	
kebiasaan_merokok	merokok	Count	89	9	98
		% within kebiasaan_merokok	90.8%	9.2%	100.0%
	tidak merokok	Count	21	48	69
		% within kebiasaan_merokok	30.4%	69.6%	100.0%
Total		Count	110	57	167
		% within kebiasaan_merokok	65.9%	34.1%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	65.665 ^a	1	.000		
Continuity Correction ^b	63.006	1	.000		
Likelihood Ratio	69.469	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	65.272	1	.000		
N of Valid Cases	167				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 23.55.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for kebiasaan_merokok (merokok / tidak merokok)	22.603	9.602	53.208
For cohort hipertensi = hipertensi	2.984	2.077	4.287
For cohort hipertensi = normal	.132	.069	.251
N of Valid Cases	167		