

Hasil olah data dengan SPSS

Statistics

		sampel	sbl1	ssd1	sbl2	ssd2
N	Valid	10	10	10	10	10
	Missing	0	0	0	0	0
Mean		5.50	62.50	92.20	72.30	82.80
Median		5.50	62.00	95.00	74.50	85.50
Mode		1 ^a	62	95	80	77 ^a
Std. Deviation		3.028	8.196	6.408	8.945	7.800
Sum		55	625	922	723	828

a. Multiple modes exist. The smallest value is shown

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
sbl1	.224	10	.167	.919	10	.345
ssd1	.269	10	.039	.918	10	.338
sbl2	.175	10	.200 [*]	.931	10	.459
ssd2	.205	10	.200 [*]	.881	10	.134

a. Lilliefors Significance Correction

*. This is a lower bound of the true significance.

Statistics

		selisih1	selisih2
N	Valid	10	10
	Missing	0	0
Mean		29.70	10.50
Std. Deviation		6.413	1.780

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	Df	Sig.	Statistic	df	Sig.
selisih1	.163	10	.200*	.933	10	.477
selisih2	.211	10	.200*	.947	10	.633

a. Lilliefors Significance Correction

*. This is a lower bound of the true significance.

Paired Samples Test

	Paired Differences					t	df	Sig. (2-tailed)
	Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
				Lower	Upper			
Pair 1 sbl1abd - ssd1abd	-29.700	6.413	2.028	-34.287	-25.113	-14.646	9	.000

Paired Samples Test

	Paired Differences					t	df	Sig. (2-tailed)
	Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
				Lower	Upper			
Pair 1 sbl2abd - ssd2abd	-10.500	1.780	.563	-11.773	-9.227	18.659	9	.000

Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
selisihp1p2abd	Equal variances assumed	12.709	.002	9.123	18	.000	19.200	2.104	14.779	23.621
	Equal variances not assumed			9.123	10.378	.000	19.200	2.104	14.534	23.866