

Statistics

		sebelum1	sesudah1	selisih1	sebelum2	sesudah2	selisih2
N	Valid	7	7	7	7	7	7
	Missing	7	7	7	7	7	7
Mean		59.43	60.71	1.29	59.86	60.71	.86
Std. Deviation		6.106	5.499	1.113	4.880	4.751	.690

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
sebelum1	.284	7	.092	.892	7	.287
sesudah1	.261	7	.164	.895	7	.301
sebelum2	.169	7	.200*	.957	7	.791
sesudah2	.184	7	.200*	.955	7	.773

a. Lilliefors Significance Correction

*. This is a lower bound of the true significance.

Paired Samples Test

	Paired Differences					t	df	Sig. (2-tailed)
	Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
				Lower	Upper			
Pair 1 sebelum1 - sesudah1	-1.286	1.113	.421	-2.315	-.257	-3.057	6	.022
Pair 2 sebelum2 - sesudah2	-.857	.690	.261	-1.495	-.219	-3.286	6	.017

Independent Samples Test

	Levene's Test for Equality of Variances	t-test for Equality of Means								
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
selisihp1p2	Equal variances assumed	2.362	.150	.866	12	.403	.429	.495	-.650	1.507
	Equal variances not assumed			.866	10.021	.407	.429	.495	-.674	1.531

Levene's Test of Equality of Error Variances^a

Dependent Variable:sebelump1p2

F	df1	df2	Sig.
.014	1	12	.907