

## DAFTAR PUSTAKA

- AdemolaGbiri, C., & Okafor, U. A. C. (2013). Comparative Efficacy of Open-chain and Close-chain Kinematics on Proprioception, Muscles Strength and Functional Performances in Individual with Knee Osteoarthritis. *Occupational Medicine & Health Affairs.* Vol: 2013. <https://doi.org/10.4172/2329-6879.1000104>
- Alrushud, A. S., Rushton, A. B., Bhogal, G., Pressdee, F., & Greig, C. A. (2018). Effect of a combined programme of dietary restriction and physical activity on the physical function and body composition of obese middle-aged and older adults with knee OA (DRPA): Protocol for a feasibility study. *BMJ Open*, vol 2018;8:e021051. <https://doi.org/10.1136/bmjopen-2017-021051>
- Bytyqi, D., Shabani, B., Lustig, S., Cheze, L., Karahoda Gjurgjeala, N., & Neyret, P. (2014). Gait knee kinematic alterations in medial osteoarthritis: Three dimensional assessment. *International Orthopaedics.* Vol: 2014 (38):1191-1198. <https://doi.org/10.1007/s00264-014-2312-3>
- Cayir, Y., Aslan, S. M., & Akturk, Z. (2015). The effect of pedometer use on physical activity and body weight in obese women. *European Journal of Sport Science.* Vol. 5, No. 4: 351 - 356 <https://doi.org/10.1080/17461391.2014.940558>
- Cudejko, T., Van Der Esch, M., Schrijvers, J., Richards, R., Van Den Noort, J. C., Wrigley, T., Van Der Leeden, M., Roorda, L. D., Lems, W., Harlaar, J., & Dekker, J. (2018). The immediate effect of a soft knee brace on dynamic knee instability in persons with knee osteoarthritis. *Rheumatology (United Kingdom).* <https://doi.org/10.1093/rheumatology/key162>
- Culvenor, A., Engen, C., Øiestad, B., Engebretsen, L., & Risberg, M. (2014). Defining radiographic knee osteoarthritis: a comparison between the Kellgren & Lawrence classification and oarsi atlas. *Osteoarthritis and Cartilage.* Vol: 22 (2014) S57 - S489. <https://doi.org/10.1016/j.joca.2014.02.500>

Fukuda, T. Y., Fingerhut, D., Moreira, V. C., Camarini, P. M. F., Scodeller, N. F., Duarte, A., Martinelli, M., & Bryk, F. F. (2013). Open kinetic chain exercises in a restricted range of motion after anterior cruciate ligament reconstruction: A randomized controlled clinical trial. *American Journal of Sports Medicine*. Vol. 41 , No.4. <https://doi.org/10.1177/0363546513476482>.

Goodman, C. C., & Fuller, K. S. (2015). Pathology: Implications for the Physical Therapist Fourth Edition. In *Saunders Elsevier*.

Hackinger, S., Trajanoska, K., Styrkarsdottir, U., Zengini, E., Steinberg, J., Ritchie, G. R. S., Hatzikotoulas, K., Gilly, A., Evangelou, E., Kemp, J. P., Evans, D., Ingvarsson, T., Jonsson, H., Thorsteinsdottir, U., Stefansson, K., McCaskie, A. W., Brooks, R. A., Wilkinson, J. M., Rivadeneira, F., & Zeggini, E. (2017). Evaluation of shared genetic aetiology between osteoarthritis and bone mineral density identifies SMAD3 as a novel osteoarthritis risk locus. *Human Molecular Genetics*. Vol. 26, No. 19 : 3850-3858. <https://doi.org/10.1093/hmg/ddx285>

Jagtap, V., & Shanmugam, S. (2014). Effect of Mechanical Traction in Osteoarthritis Knee. *International Journal of Science and Research*.

Kim, H.-H., & Shim, J.-M. (2016). Comparison of Forward and Backward Walking Trainings on Gait Pattern in Adults. *Indian Journal of Science and Technology*. Vol. 9 (43): 0974 - 5645. <https://doi.org/10.17485/ijst/2016/v9i43/105030>

Kisner, C., & Colby, L. A. (2012). Therapeutic Exercise Foundations and Techniques, Sixth Edition, F.A. Davis Company, America.

Konstantinidis, G. A., Aletras, V. H., Kanakari, K. A., Natsis, K., Bellamy, N., & Niakas, D. (2014). Comparative validation of the WOMAC osteoarthritis and Lequesne algofunctional indices in Greek patients with hip or knee osteoarthritis. *Quality of Life Research*. <https://doi.org/10.1007/s11136-013-0490-x>

- Kramer, J. F., & Reid, D. C. (1981). Backward walking: A cinematographic and electromyographic pilot study. *Physiotherapy Canada*.
- Kwon, Y. J., Park, S. J., Jefferson, J., & Kim, K. (2013). The effect of open and closed kinetic chain exercises on dynamic balance ability of normal healthy adults. *Journal of Physical Therapy Science*. Vol: 2013 (25) : 671 - 674. <https://doi.org/10.1589/jpts.25.671>
- Levangie, P. K., & Norkin, C. C. (2011). *Joint Structure and Function A Comprehensive Analysis* (5th ed.). Library of Congress Cataloging.
- Logde, A., & Borkar, P. (2018). Effect of retro walking on hamstring flexibility in normal healthy individual. *International Journal of Physical Education, Sports and Health*. <https://doi.org/10.5455/ijtrr.00000017>
- Malgaonkar, P. P., N, S. K., K, V. B., & Rizvi, S. R. (2014). Short Term Effect of Mulligan's Mobilization Versus Kinesio Taping on Knee Pain and Disability for Osteoarthritis of Knee. *International Journal of Physiotherapy*. <https://doi.org/10.15621/ijphy/2014/v1i4/54566>
- Masouros, S. D., Bull, A. M. J., & Amis, A. A. (2010). (i) Biomechanics of the knee joint. *Orthopaedics and Trauma*. Vol. 4, No. 2. <https://doi.org/10.1016/j.morth.2010.03.005>
- Minerva, R. K., Kumar Alagingi, N., Apparao, P., Graduate Student, P., professor, A., Principal, V., & Author, C. (2016). To Compare the Effectiveness of Maitland versus Mulligan Mobilisation in Idiopathic Adhesive Capsulitis of Shoulder. *International Journal of Health Sciences & Research (Www.Ijhsr.Org)*.
- Parentoni, A. N., Mendonça, V. A., Dos Santos, K. D., Sá, L. F., Ferreira, F. O., Gomes Pereira, D. A., & Lustosa, L. P. (2015). Gait Speed as a Predictor of Respiratory Muscle Function, Strength, and Frailty Syndrome in Community-Dwelling Elderly People. *The Journal of Frailty & Aging*. Vol. 4 (2): 64-68. <https://doi.org/10.14283/jfa.2015.41>

- Rao, R. V., Balthillaya, G., Prabhu, A., & Kamath, A. (2018). Immediate effects of Maitland mobilization versus Mulligan Mobilization with Movement in Osteoarthritis knee- A Randomized Crossover trial. *Journal of Bodywork and Movement Therapies*. <https://doi.org/10.1016/j.jbmt.2017.09.017>
- Roos, P. E., Barton, N., & van Deursen, R. W. M. (2012). Patellofemoral joint compression forces in backward and forward running. *Journal of Biomechanics*. <https://doi.org/10.1016/j.jbiomech.2012.03.020>
- Sambandam, C. E., Sailor, S. N., & Alagesan, J. (2011). Effect of Mulligan Mobilization and Maitland Mobilization in Subjects with Unilateral Tibiofemoral Osteoarthritis - Randomized Controlled Trial Method : *journal of pharmaceutical and biomedical sciences*.
- Sedhom, M. G. (2017). Backward Walking Training Improves Knee Proprioception in Non Athletic Males. *International Journal of Physiotherapy*. <https://doi.org/10.15621/ijphy/2017/v4i1/136161>
- Sitoci, K. H., Hudelmaier, M., & Eckstein, F. (2012). Nocturnal changes in knee cartilage thickness in young healthy adults. *Cells Tissues Organs*. <https://doi.org/10.1159/000333456>.
- Tseng, I. J., Yuan, R. Y., & Jeng, C. (2015). Treadmill training improves forward and backward gait in early Parkinson disease. *American Journal of Physical Medicine and Rehabilitation*. <https://doi.org/10.1097/PHM.0000000000000273>
- Verma, S. (2012). Comparing open kinetic chain with closed kinetic chain exercise on quadriceps strength and functional status of women with osteoarthritic knees. *Sports Medicine Journal / Medicina Sportiva*.
- Wang, J., Xu, J., & An, R. (2019). Effectiveness of backward walking training on balance performance: A systematic review and meta-analysis. In *Gait and Posture*. <https://doi.org/10.1016/j.gaitpost.2019.01.002>