

Tabel nilai penurunan nyeri pada kelompok perlakuan I

Intervensi TENS						
Sampel	Nilai VAS Sebelum I	Nilai VAS Sesudah I	Nilai VAS Sebelum II	Nilai VAS Sesudah II	Selisih	%
1	56	54	42	38	18	32,14
2	53	48	45	43	10	18,87
3	73	46	40	34	39	53,42
4	100	81	71	64	36	36,00
5	81	74	65	52	29	35,80
6	75	63	53	49	26	34,67
7	69	58	53	46	23	33,33
Mean	72,43	60,57	52,71	46,57	25,86	34,89
SD	15,81	13,06	11,70	9,86	10,06	10,10

Tabel nilai penurunan nyeri pada kelompok perlakuan II

Intervensi IFC						
Sampel	Nilai VAS Sebelum I	Nilai VAS Sesudah I	Nilai VAS Sebelum II	Nilai VAS Sesudah II	Selisih	%
1	72	49	33	19	53	73,61
2	55	48	41	19	36	65,45
3	57	45	41	28	29	50,88
4	91	60	39	24	67	73,63
5	83	65	37	22	61	73,49
6	62	53	48	31	31	50
7	61	45	26	15	46	75,41
Mean	68,71	52,14	37,86	22,57	46,14	66,07
SD	13,79	7,71	6,93	5,56	14,88	11,15

Tabel distribusi rata-rata penurunan nyeri pada kelompok perlakuan I dan kelompok perlakuan II

Kelompok Perlakuan	Nilai VAS Sebelum I	Nilai VAS Sesudah II	Nilai VAS Sebelum II	Nilai VAS Sesudah II
Perlakuan I	72,43	60,57	52,71	46,57
Perlakuan II	68,71	52,14	37,86	22,57