


LAMPIRAN ANALISIA UNIVARIAT

1. ANALISA UNIVARIAT

Tabel Frekuensi Menurut Umur

		Kategori Umur			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Dewasa	2903	48.8	48.8	48.8
	Lansia	3040	51.2	51.2	100.0
	Total	5943	100.0	100.0	


Grafik Frekuensi Menurut Umur



Tabel Frekuensi Menurut Jenis Kelamin

		Jenis Kelamin			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Laki-laki	3406	57.3	57.3	57.3
	Perempuan	2537	42.7	42.7	100.0
Total		5943	100.0	100.0	


Grafik Frekuensi Menurut Jenis Kelamin



Tabel Frekuensi Menurut Pola Konsumsi Bumbu Penyedap

Pola Konsumsi Bumbu Penyedap					
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Sering	5666	95.3	95.3	95.3
	Jarang	277	4.7	4.7	100.0
	Total	5943	100.0	100.0	


Grafik Menurut Pola Konsumsi Bumbu Penyedap



Tabel Frekuensi Menurut Pola Konsumsi Makanan Asin

		Pola Konsumsi Makanan Asin			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Sering	2014	33.9	33.9	33.9
	Jarang	3929	66.1	66.1	100.0
	Total	5943	100.0	100.0	


Grafik Menurut Pola Konsumsi Makanan Asin



Tabel Frekuensi Menurut Status Gizi

		Status Gizi			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Kurang	220	3.7	3.7	3.7
	Normal	2988	50.3	50.3	54.0
	Lebih	1235	20.8	20.8	74.8
	Obesitas	1500	25.2	25.2	100.0
	Total	5943	100.0	100.0	


Grafik Menurut Status Gizi



Tabel Menurut Tipe Daerah

		Tipe Daerah			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Pedesaan	2676	45.0	45.0	45.0
	Perkotaan	3267	55.0	55.0	100.0
	Total	5943	100.0	100.0	


Grafik Menurut Tipe Daerah



Tabel Frekuensi Menurut Status Hipertensi

		STATUS_HIPERTENSI			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak	4027	67.8	67.8	67.8
	Ya	1916	32.2	32.2	100.0
Total		5943	100.0	100.0	

Grafik Menurut Status Hipertensi



2. ANALISA BIVARIAT

UJI *CHI-SQUARE* HUBUNGAN UMUR TERHADAP KEJADIAN HIPERTENSI DI PROVINSI BALI

Kategori Umur * STATUS_HIPERTENSI Crosstabulation

			STATUS_HIPERTENSI		Total
			Tidak	Ya	
Kategori Umur	Dewasa	Count	2210	693	2903
		% within Kategori Umur	76.1%	23.9%	100.0%
		% of Total	37.2%	11.7%	48.8%
	Lansia	Count	1817	1223	3040
		% within Kategori Umur	59.8%	40.2%	100.0%
		% of Total	30.6%	20.6%	51.2%
Total	Count	4027	1916	5943	
	% within Kategori Umur	67.8%	32.2%	100.0%	
	% of Total	67.8%	32.2%	100.0%	

Chi-Square Tests

	Value	Df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.819E2 ^a	1	.000		
Continuity Correction ^b	181.151	1	.000		
Likelihood Ratio	183.793	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	181.869	1	.000		
N of Valid Cases ^b	5943				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 935.92.

b. Computed only for a 2x2 table

HUBUNGAN JENIS KELAMIN TERHADAP KEJADIAN HIPERTENSI DI PROVINSI BALI

Jenis Kelamin * STATUS_HIPERTENSI Crosstabulation

			STATUS_HIPERTENSI		Total
			Tidak	Ya	
Jenis Kelamin	Laki-laki	Count	2269	1137	3406
		% within Jenis Kelamin	66.6%	33.4%	100.0%
		% of Total	38.2%	19.1%	57.3%
	Perempuan	Count	1758	779	2537
		% within Jenis Kelamin	69.3%	30.7%	100.0%
		% of Total	29.6%	13.1%	42.7%
Total	Count	4027	1916	5943	
	% within Jenis Kelamin	67.8%	32.2%	100.0%	
	% of Total	67.8%	32.2%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.769 ^a	1	.029		
Continuity Correction ^b	4.647	1	.031		
Likelihood Ratio	4.780	1	.029		
Fisher's Exact Test				.031	.015
Linear-by-Linear Association	4.768	1	.029		
N of Valid Cases ^b	5943				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 817.92.

b. Computed only for a 2x2 table

HUBUNGAN POLA KONSUMSI BUMBU PENYEDAP TERHADAP KEJADIAN HIPERTENSI DI PROVINSI BALI

Pola Konsumsi Bumbu Penyedap * STATUS_HIPERTENSI Crosstabulation

			STATUS_HIPERTENSI		Total
			Tidak	Ya	
Pola Konsumsi Bumbu Penyedap	Sering	Count	3844	1822	5666
		% within Pola Konsumsi Bumbu Penyedap	67.8%	32.2%	100.0%
		% of Total	64.7%	30.7%	95.3%
	Jarang	Count	183	94	277
		% within Pola Konsumsi Bumbu Penyedap	66.1%	33.9%	100.0%
		% of Total	3.1%	1.6%	4.7%
Total		Count	4027	1916	5943
		% within Pola Konsumsi Bumbu Penyedap	67.8%	32.2%	100.0%
		% of Total	67.8%	32.2%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.382 ^a	1	.536		
Continuity Correction ^b	.305	1	.581		
Likelihood Ratio	.379	1	.538		
Fisher's Exact Test				.554	.289
Linear-by-Linear Association	.382	1	.536		
N of Valid Cases ^b	5943				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 89.30.

b. Computed only for a 2x2 table

HUBUNGAN POLA KONSUMSI MAKANAN ASIN TERHADAP KEJADIAN HIPERTENSI DI PROVINSI BALI

Pola Konsumsi Makanan Asin * STATUS_HIPERTENSI Crosstabulation

			STATUS_HIPERTENSI		Total
			Tidak	Ya	
Pola Konsumsi Makanan Asin	Sering	Count	1378	636	2014
		% within Pola Konsumsi Makanan Asin	68.4%	31.6%	100.0%
		% of Total	23.2%	10.7%	33.9%
	Jarang	Count	2649	1280	3929
		% within Pola Konsumsi Makanan Asin	67.4%	32.6%	100.0%
		% of Total	44.6%	21.5%	66.1%
Total	Count		4027	1916	5943
	% within Pola Konsumsi Makanan Asin		67.8%	32.2%	100.0%
	% of Total		67.8%	32.2%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.609 ^a	1	.435		
Continuity Correction ^b	.564	1	.453		
Likelihood Ratio	.610	1	.435		
Fisher's Exact Test				.446	.227
Linear-by-Linear Association	.609	1	.435		
N of Valid Cases ^b	5943				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 649.31.

b. Computed only for a 2x2 table

HUBUNGAN STATUS GIZI TERHADAP KEJADIAN HIPERTENSI DI PROVINSI BALI

Status Gizi * STATUS_HIPERTENSI Crosstabulation

			STATUS_HIPERTENSI		Total
			Tidak	Ya	
Status Gizi	Kurang	Count	173	47	220
		% within Status Gizi	78.6%	21.4%	100.0%
		% of Total	2.9%	.8%	3.7%
	Normal	Count	2198	790	2988
		% within Status Gizi	73.6%	26.4%	100.0%
		% of Total	37.0%	13.3%	50.3%
	Lebih	Count	814	421	1235
		% within Status Gizi	65.9%	34.1%	100.0%
		% of Total	13.7%	7.1%	20.8%
Obesitas	Count	842	658	1500	
	% within Status Gizi	56.1%	43.9%	100.0%	
	% of Total	14.2%	11.1%	25.2%	
Total	Count	4027	1916	5943	
	% within Status Gizi	67.8%	32.2%	100.0%	
	% of Total	67.8%	32.2%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.527E2 ^a	3	.000
Likelihood Ratio	150.675	3	.000
Linear-by-Linear Association	151.015	1	.000
N of Valid Cases	5943		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 70.93.

HUBUNGAN TIPE DAERAH TERHADAP KEJADIAN HIPERTENSI DI PROVINSI BALI

Tipe Daerah * STATUS_HIPERTENSI Crosstabulation

			STATUS_HIPERTENSI		Total
			Tidak	Ya	
Tipe Daerah	Pedesaan	Count	1810	866	2676
		% within Tipe Daerah	67.6%	32.4%	100.0%
		% of Total	30.5%	14.6%	45.0%
	Perkotaan	Count	2217	1050	3267
		% within Tipe Daerah	67.9%	32.1%	100.0%
		% of Total	37.3%	17.7%	55.0%
Total	Count	4027	1916	5943	
	% within Tipe Daerah	67.8%	32.2%	100.0%	
	% of Total	67.8%	32.2%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.033 ^a	1	.855		
Continuity Correction ^b	.024	1	.877		
Likelihood Ratio	.033	1	.855		
Fisher's Exact Test				.867	.439
Linear-by-Linear Association	.033	1	.855		
N of Valid Cases ^b	5943				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 862.73.

b. Computed only for a 2x2 table