

DAFTAR PUSTAKA

- Ambrose, A. F., Paul, G., & Hausdorff, J. M. (2013). *Maturitas risk factors for falls among older adults : a review of the literature*. Journal Elsevier Maturitas, 75(1), 51–61.
<https://doi.org/10.1016/j.maturitas.2013.02.009>
- Ana-Belén Meseguer Henarejos, María, R., Carles-hernández, R., Gómez-conesa, A., & Rubio-aparicio, M. (2019). *Characteristics that affect score reliability in the Berg Balance Scale : A meta-analytic reliability generalization study*. European Journal of Physical and Rehabilitation Medicine EDIZIONI MINERVA MEDICA, 1–61.
<https://doi.org/10.23736/S1973-9087.19.05363-2>
- Beato, M., Dawson, N., Svien, L., & Wharton, T. (2019). *Examining the effects of an otago-based home exercise program on falls and fall risks in an assisted living facility*. Journal of Geriatric Physical Therapy, 42(4), 224–229.
<https://doi.org/10.1519/JPT.0000000000000190>
- Bellelli, G., Buccino, G., Bernardini, B., & Padovani, A. (2010). *Action observation treatment improves recovery of postsurgical orthopedic patients : evidence for a top-down effect*. YAPMR, 91(10), 1489–1494. <https://doi.org/10.1016/j.apmr.2010.07.013>
- Correia, A., Pimenta, C., Alves, M., & Virella, D. (2020). *Better balance: a randomised controlled trial of oculomotor and gaze stability exercises to reduce risk of falling after stroke*. Clinical Rehabilitation. <https://doi.org/10.1177/0269215520956338>
- Cuevas-Trisan, R. (2017). *Balance problems and fall risks in the elderly*. in Physical Medicine and Rehabilitation Clinics of North America (Vol. 28, Issue 4, pp. 727–737). W.B. Saunders. <https://doi.org/10.1016/j.pmr.2017.06.006>
- Dubey, N. K., Ningrum, D. N. A., Dubey, R., Deng, Y. H., Li, Y. C., Wang, P. D., Wang, J. R., Syed-Abdul, S., & Deng, W. P. (2018). *Correlation between diabetes mellitus and knee osteoarthritis: A dry-to-wet lab approach*. International Journal of Molecular Sciences, 19(10), 1–13. <https://doi.org/10.3390/ijms19103021>
- Felson, D. T. (2006). *Osteoarthritis of the knee*. The New England Journal of Medicine, 1–8.
- Frontera, W. R. (2017). *Physiologic changes of the musculoskeletal system with aging: A brief review*. Physical Medicine and Rehabilitation Clinics of North America, 28(4), 705–711.
<https://doi.org/10.1016/j.pmr.2017.06.004>
- Garg, H., Dibble, L. E., Schubert, M. C., Sibthorp, J., Foreman, K. B., & Gappmaier, E. (2018). *Gaze stability, dynamic balance and participation deficits in people with multiple sclerosis at fall-risk*. Anatomical Record, 301(11), 1852–1860. <https://doi.org/10.1002/ar.23852>
- Gupta, G., & Sharma, A. (2018). *Impact of osteoarthritis on balance, perceived fear of fall and quality of life*. Orthopedics and Rheumatology Open Access Journal, 11(5), 7.
<https://doi.org/10.19080/oroaj.2018.11.555824>
- Hinman, R. S., & Crossley, K. M. (2007). *Patellofemoral joint osteoarthritis: An important subgroup of knee osteoarthritis*. Rheumatology, 46(7), 1057–1062.
<https://doi.org/10.1093/rheumatology/kem114>
- Irfan, M., & Gahara, R. (2006). *Beda pengaruh penambahan long axis oscillated traction pada intervensi MWD dan TENS terhadap pengurangan rasa nyeri pada capsular pattern akibat osteoarthritis lutut*. Jurnal Fisioterapi Indonusa, 6(1), 25–35.
- KEMSSOS. (2018). *Standar nasional rehabilitasi sosial lanjut usia*. 24 Mei 2018, 121.
<https://jdih.kemsos.go.id/pencarian/www/storage/repository/PERMENSSOS NOMOR 5 TAHUN 2018 TENTANG STANDAR NASIONAL REHABILITASI SOSIAL LANJUT>

USIA.pdf

- Khanna, T., & Singh, S. (2014). *Effect of gaze stability exercises on balance in elderly*. IOSR Journal of Dental and Medical Sciences, 13(9), 41–48. <https://doi.org/10.9790/0853-13914148>
- Kholifah, S. N. (2016). *Modul keperawatan gerontik* (M. Dwisatyadin (ed.); 1st ed.). KEMENTERIAN KESEHATAN REPUBLIK INDONESIA.
<http://bppsdmk.kemkes.go.id/pusdiksdmk/wp-content/uploads/2017/08/Keperawatan-Gerontik-Komprehensif.pdf>
- Kohn, M. D., Sassoon, A. A., & Fernando, N. D. (2016). *Classifications in brief: kellgren-lawrence classification of osteoarthritis*. Clinical Orthopaedics and Related Research, 474(8), 1886–1893. <https://doi.org/10.1007/s11999-016-4732-4>
- Kuptniratsaikul, V., Praditsuwan, R., Assantachai, P., Ploypitch, T., Udompunturak, S., & Pooliam, J. (2011). *Effectiveness of simple balancing training program in elderly patients with history of frequent falls*. Clinical Interventions in Aging, 6(1), 111–117. <https://doi.org/10.2147/CIA.S17851>
- Lee, J., Yoo, H. N., & Lee, B. H. (2017). *Effects of augmented reality-based Otago exercise on balance, gait, and physical factors in elderly women to prevent falls: A randomized controlled trial*. Journal of Physical Therapy Science, 29(9), 1586–1589. <https://doi.org/10.1589/jpts.29.1586>
- Lespasio, M. J., Piuzzi, N. S., Husni, M. E., Muschler, G. F., Guarino, A., & Mont, M. A. (2017). *Knee osteoarthritis: A primer*. The Permanente Journal, 21, 1–7. <https://doi.org/10.7812/TPP/16-183>
- Lima, C. A., Ricci, N. A., Nogueira, E. C., & Perracini, M. R. (2018). *The berg balance scale as a clinical screening tool to predict fall risk in older adults: a systematic review*. In Physiotherapy (United Kingdom) (Vol. 104, Issue 4, pp. 383–394). Elsevier Ltd. <https://doi.org/10.1016/j.physio.2018.02.002>
- Magnusson, K., Turkiewicz, A., & Englund, M. (2019). *Nature vs nurture in knee osteoarthritis – the importance of age, sex and body mass index*. Osteoarthritis and Cartilage, 27(4), 586–592. <https://doi.org/10.1016/j.joca.2018.12.018>
- Maita, L., Nurlisis, N., & Pitriani, R. (2013). *Karakteristik wanita dengan keluhan masa menopause di wilayah kerja puskesmas rejosari*. Jurnal Kesehatan Komunitas, 2(3), 128–131. <https://doi.org/10.25311/keskom.vol2.iss3.59>
- Maratis, J. (2020). *Pengaruh senam rhythmic auditory stimulation (RAS) dengan kemampuan berjalan lansia di desa kohod kabupaten tanggerang*. Fisioterapi Jurnal, 20(April).
- Mat, S., Ng, C. T., Tan, P. J., Ramli, N., Fadzli, F., Rozalli, F. I., Mazlan, M., Hill, K. D., & Tan, M. P. (2018). *Effect of modified otage exercises on postural balance, fear of falling, and fall risk in older fallers with knee osteoarthritis and impaired gait and balance: A secondary analysis*. PM and R, 10(3), 254–262. <https://doi.org/10.1016/j.pmrj.2017.08.405>
- Matsugi, A., Ueta, Y., Oku, K., Okuno, K., Tamaru, Y., Nomura, S., Tanaka, H., & Mori, N. (2017). *Effect of gaze-stabilization exercises on vestibular function during postural control*. NeuroReport, 28(8), 439–443. <https://doi.org/10.1097/WNR.0000000000000776>
- Miller, C. A. (2012). *Nursing for wellness in older adult sixth edition* (E. Nieginski (ed.); sixth). Wolters Kluwer Health | Lippincott Williams & Wilkins.
- Munawwarah, M. (2020). *Modul pendahuluan fisoterapi kesehatan wanita* (pp. 1–16). Universitas Esa Unggul.
- Munawwarah, M., & N, P. N. (2015). *Pemberian latihan pada lansia dapat meningkatkan*

- keseimbangan dan mengurangi resiko jatuh lansia.* Fisioterapi Jurnal, 15(April).
- Nevitt, M. C. (2015). *Symptoms of knee instability are risk factors for recurrent falls.* Arthritis Care & Research, 1–37. <https://doi.org/10.1002/acr.22811>
- Njoto, I. (2018). *Epidemiologi, Pategensis dan Faktor Resiko Osteoarthritis.* <http://www.ghbook.ir/index.Fakultas Kedokteran Universitas Wijaya Kusuma Surabaya>
- Orimo, H., Ito, H., Suzuki, T., Araki, A., Hosoi, T., & Sawabe, M. (2006). *Reviewing the definition of “elderly.”* Geriatrics and Gerontology International, 6(3), 149–158. <https://doi.org/10.1111/j.1447-0594.2006.00341.x>
- Park, S., & Lee, Y. (2016). *The diagnostic accuracy of the berg balance scale in predicting falls.* Western Journal of Nursing Research, 1 –24. <https://doi.org/10.1177/0193945916670894>
- Pimenta, C., Correia, A., Alves, M., & Virella, D. (2017). *Effects of oculomotor and gaze stability exercises on balance after stroke: Clinical trial protocol.* Porto Biomedical Journal, 2(3), 76–80. <https://doi.org/10.1016/j.pbj.2017.01.003>
- Pratama, S. A., Sukmanintyas, H., & Warlisti, I. V. (2019). *Hubungan hipertensi dengan derajat defek kerusakan kartilago osteoarthritis knee berdasarkan ultrasonografi.* Diponegoro Medical Journal (Jurnal Kedokteran Diponegoro), 8(1), 501–508.
- Priatna, H. (2019). *Pendekatan fisioterapi terkini pada kasus osteoarthritis knee* (p. 46).
- Rasyiqah, F., & Khairani. (2019). *Cognitive function between the risk level of falling in the elderly in banda aceh.* Idea Nursing Journal, X(2), 40–46.
- Roh, M., & Lee, E. (2019). *Effects of gaze stability exercises on cognitive function , dynamic postural ability , balance confidence , and subjective health status in old people with mild cognitive impairment.* Journal of Exercise Rehabilitation, 15(2), 270–274. <https://doi.org/10.12965/jer.1938026.013>
- Smith, T. O., Higson, E., Pearson, M., & Mansfield, M. (2018). *Is there an increased risk of falls and fractures in people with early diagnosed hip and knee osteoarthritis? data from the osteoarthritis initiative.* International Journal of Rheumatic Diseases, 21(6), 1193–1201. <https://doi.org/10.1111/1756-185X.12871>
- Sudrajat, W. A., & Soetardji. (2014). *Efek pemberian latihan keseimbangan dalam mempertahankan kemampuan keseimbangan manula panti wredha rindang asih 1 ungaran.* Journal of Sport Sciences and Fitness, 3(1), 49–54.
- Tinetti, M. E. (2003). *Preventing falls in elderly persons.* New England Journal of Medicine, 348(1), 42–49. <https://doi.org/10.1056/nejmcp020719>
- Van Manen, M. D., Nace, J., & Mont, M. A. (2012). *Management of primary knee osteoarthritis and indications for total knee arthroplasty for general practitioners.* Journal of the American Osteopathic Association, 112(11), 709–715. <https://doi.org/10.7556/jaoa.2012.112.11.709>
- van Schoor, N. M., Dennison, E., Castell, M. V., Cooper, C., Edwards, M. H., Maggi, S., Pedersen, N. L., van der Pas, S., Rijnhart, J. J. M., Lips, P., & Deeg, D. J. H. (2020). *Clinical osteoarthritis of the hip and knee and fall risk: The role of low physical functioning and pain medication.* Seminars in Arthritis and Rheumatism, 50(3), 380–386. <https://doi.org/10.1016/j.semarthrit.2020.02.006>

