

LAMPIRAN 1

HASIL UJI ORGANOLEPTIK MUTU HEDONIK

SAMPEL I			
WARNA	AROMA	TEKSTUR	RASA
2	3	4	3
2	2	4	4
2	2	4	4
2	1	4	5
2	2	4	5
2	2	5	4
2	2	4	3
2	4	5	5
2	2	4	5
2	2	3	3
1	2	3	5
4	4	2	3
2	1	4	3
1	2	4	4
2	4	4	3
2	2	5	3
2	2	4	4
2	2	4	4
2	3	4	5
3	3	4	5
3	4	3	4
2	2	4	4
2	3	4	3
1	2	4	3
2	2	4	3
2	1	5	5
3	2	5	4
4	2	4	4
2	2	4	4
2	1	4	4

SAMPEL II			
WARNA	AROMA	TEKSTUR	RASA
2	3	5	4
1	2	4	5
1	1	5	4
2	1	4	4
2	2	4	3
1	4	1	4
2	2	1	4
3	2	1	4
2	3	1	5
3	2	5	4
3	2	4	5
1	3	5	4
1	2	5	4
2	1	4	4
2	3	3	3
2	2	3	4
1	2	4	4
1	2	3	3
1	2	4	4
1	1	4	4
5	2	3	3
5	3	3	4
2	2	3	5
2	1	3	3
1	3	3	4
3	1	3	4
2	2	4	2
1	3	4	3
2	2	3	5
2	1	3	3

SAMPEL III			
WARNA	AROMA	TEKSTUR	RASA
1	3	3	4
1	4	4	4
1	4	4	4
3	4	5	5
2	2	3	5
4	3	4	3
3	3	3	5
2	2	5	4
4	4	4	4
3	4	4	3
2	3	5	4
1	2	3	4
4	3	4	3
4	2	4	4
5	1	4	4
5	2	4	3
4	1	4	4
4	2	4	5
4	3	4	4
3	2	5	4
4	1	4	4
2	1	4	5
4	1	5	5
5	1	3	4
1	3	4	4
4	2	3	3
5	1	3	5
1	3	4	4
4	2	4	4
1	2	4	4

SAMPEL IV			
WARNA	AROMA	TEKSTUR	RASA
1	2	2	2
2	3	3	3
1	2	4	3
3	2	4	4
2	2	4	2
3	4	3	3
1	2	2	4
1	1	4	4
1	2	3	4
5	3	2	2
1	2	2	2
3	3	4	3
2	2	2	2
2	2	4	4
2	3	3	4
1	1	4	4
2	2	4	4
2	1	3	3
3	2	4	4
5	3	4	4
4	3	2	4
4	2	3	4
3	3	4	5
1	2	4	3
2	4	2	2
3	2	2	2
2	2	2	2
5	2	4	4
1	2	4	4
1	2	3	3

LAMPIRAN 2

HASIL UJI ORGANOLEPTIK HEDONIK

SAMPEL I				
WARNA	AROMA	TEKSTUR	RASA	KESELURUHAN
3	3	4	4	5
5	7	5	5	6
5	5	5	6	6
3	4	4	4	5
3	3	4	3	4
5	5	5	5	5
5	5	5	5	4
7	6	7	7	5
5	6	5	6	6
4	5	5	6	6
5	7	4	4	5
4	4	4	4	4
6	5	5	4	5
5	5	5	6	6
5	5	5	6	6
5	6	6	6	6
5	5	5	5	5
4	5	5	5	5
6	6	6	5	5
4	5	4	4	4
5	4	5	5	5
5	4	5	5	5
5	5	5	5	5
5	4	6	6	7
4	6	5	5	7
4	5	5	5	5
4	4	5	4	5
4	5	4	5	5
3	6	6	6	5
3	4	7	7	7

SAMPEL II				
WARNA	AROMA	TEKSTUR	RASA	KESELURUHAN
3	4	5	4	4
6	7	5	6	6
5	5	5	5	5
4	5	5	5	5
3	4	4	5	5
5	4	5	5	5
2	5	5	5	6
1	5	7	5	6
5	4	6	6	4
1	5	6	6	6
3	6	5	3	4
5	5	5	5	6
1	5	6	5	6
5	5	5	5	5
4	5	4	5	5
5	4	6	7	7
5	4	5	4	5
5	4	4	4	4
1	5	2	7	7
4	4	4	2	3
4	5	4	4	4
1	5	5	4	4
2	4	2	2	4
5	5	5	5	5
3	4	3	4	4
5	5	5	5	5
4	4	5	5	5
3	4	4	3	5
4	4	4	2	3
3	5	3	1	3

SAMPel III				
WARNA	AROMA	TEKSTUR	RASA	KESELURUHAN
3	5	4	5	4
5	7	5	5	5
5	5	5	7	5
2	6	2	6	6
1	4	4	7	5
4	6	4	7	6
4	4	5	7	5
7	5	6	6	6
4	5	5	5	5
5	5	6	5	5
7	4	6	6	5
5	4	5	7	5
6	4	5	5	5
5	5	5	5	5
5	4	5	5	6
5	5	5	4	5
5	4	6	4	7
5	6	5	7	5
6	5	6	6	5
2	5	6	2	7
4	5	7	5	5
5	5	5	6	7
4	4	5	5	4
5	5	6	5	6
4	4	5	5	6
4	4	6	5	5
4	4	5	5	5
6	5	5	6	6
4	4	5	5	6
5	4	5	5	7

SAMPEL IV				
WARNA	AROMA	TEKSTUR	RASA	KESELURUHAN
4	5	2	3	5
4	6	5	5	5
5	5	5	4	5
3	5	4	4	5
2	6	3	1	4
3	5	4	4	4
5	5	5	5	5
6	4	6	6	5
5	6	5	5	5
5	4	3	3	4
2	5	3	2	3
1	4	3	4	4
5	5	3	4	4
1	6	6	6	6
2	5	5	5	5
6	7	6	6	6
3	4	5	5	5
5	5	5	5	6
6	7	7	7	7
3	5	4	2	3
5	5	3	3	4
2	5	5	4	3
2	4	5	3	5
5	5	5	5	5
3	4	4	3	2
3	4	3	3	5
5	5	4	5	6
4	4	4	3	4
4	5	5	5	5
5	5	5	5	6

LAMPIRAN 3

HASIL UJI STATISTIK UJI MUTU HEDONIK TERHADAP RASA NUGGET KEONG SAWAH

Test of Homogeneity of Variances

RASA

Levene Statistic	df1	df2	Sig.
2.912	3	116	.037

ANOVA

RASA

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	11.892	3	3.964	6.578	.000
Within Groups	69.900	116	.603		
Total	81.792	119			

Descriptives

RASA

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
					T0	30		
T1	30	3.87	.730	.133	3.59	4.14	2	5
T2	30	4.10	.662	.121	3.85	4.35	3	5
T3	30	3.27	.907	.166	2.93	3.61	2	5
Total	120	3.79	.829	.076	3.64	3.94	2	5

Multiple Comparisons

Dependent Variable: RASA

Bonferroni

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.067	.200	1.000	-.47	.60
	T2	-.167	.200	1.000	-.70	.37
	T3	.667*	.200	.007	.13	1.20
T1	T0	-.067	.200	1.000	-.60	.47
	T2	-.233	.200	1.000	-.77	.30
	T3	.600*	.200	.020	.06	1.14
T2	T0	.167	.200	1.000	-.37	.70
	T1	.233	.200	1.000	-.30	.77
	T3	.833*	.200	.000	.30	1.37
T3	T0	-.667*	.200	.007	-1.20	-.13
	T1	-.600*	.200	.020	-1.14	-.06
	T2	-.833*	.200	.000	-1.37	-.30

*. The mean difference is significant at the 0.05 level.

Multiple Comparisons

Dependent Variable: RASA

Tukey HSD

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.067	.200	.987	-.46	.59
	T2	-.167	.200	.839	-.69	.36
	T3	.667*	.200	.006	.14	1.19
T1	T0	-.067	.200	.987	-.59	.46
	T2	-.233	.200	.651	-.76	.29
	T3	.600*	.200	.018	.08	1.12
T2	T0	.167	.200	.839	-.36	.69
	T1	.233	.200	.651	-.29	.76
	T3	.833*	.200	.000	.31	1.36
T3	T0	-.667*	.200	.006	-1.19	-.14
	T1	-.600*	.200	.018	-1.12	-.08
	T2	-.833*	.200	.000	-1.36	-.31

*. The mean difference is significant at the 0.05 level.

RASA

Tukey HSD^a

PERLAKUAN	N	Subset for alpha = 0.05	
		1	2
T3	30	3.27	
T1	30		3.87
T0	30		3.93
T2	30		4.10
Sig.		1.000	.651

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.

LAMPIRAN 4

HASIL UJI STATISTIK UJI MUTU HEDONIK TERHADAP WARNA NUGGET KEONG SAWAH

Test of Homogeneity of Variances

WARNA

Levene Statistic	df1	df2	Sig.
9.778	3	116	.000

ANOVA

WARNA

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	17.292	3	5.764	4.240	.007
Within Groups	157.700	116	1.359		
Total	174.992	119			

Descriptives

WARNA

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
T0	30	2.13	.681	.124	1.88	2.39	1	4
T1	30	1.97	1.066	.195	1.57	2.36	1	5
T2	30	2.97	1.474	.269	2.42	3.52	1	5
T3	30	2.30	1.291	.236	1.82	2.78	1	5
Total	120	2.34	1.213	.111	2.12	2.56	1	5

Multiple Comparisons

Dependent Variable: WARNA

Bonferroni

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.167	.301	1.000	-.64	.97
	T2	-.833*	.301	.039	-1.64	-.03
	T3	-.167	.301	1.000	-.97	.64
T1	T0	-.167	.301	1.000	-.97	.64
	T2	-1.000*	.301	.007	-1.81	-.19
	T3	-.333	.301	1.000	-1.14	.47
T2	T0	.833*	.301	.039	.03	1.64
	T1	1.000*	.301	.007	.19	1.81
	T3	.667	.301	.173	-.14	1.47
T3	T0	.167	.301	1.000	-.64	.97
	T1	.333	.301	1.000	-.47	1.14
	T2	-.667	.301	.173	-1.47	.14

*. The mean difference is significant at the 0.05 level.

Multiple Comparisons

Dependent Variable: WARNA

Tukey HSD

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.167	.301	.945	-.62	.95
	T2	-.833*	.301	.033	-1.62	-.05
	T3	-.167	.301	.945	-.95	.62
T1	T0	-.167	.301	.945	-.95	.62
	T2	-1.000*	.301	.006	-1.78	-.22
	T3	-.333	.301	.686	-1.12	.45
T2	T0	.833*	.301	.033	.05	1.62
	T1	1.000*	.301	.006	.22	1.78
	T3	.667	.301	.125	-.12	1.45
T3	T0	.167	.301	.945	-.62	.95
	T1	.333	.301	.686	-.45	1.12
	T2	-.667	.301	.125	-1.45	.12

*. The mean difference is significant at the 0.05 level.

WARNA

Tukey HSD^a

PERLAKUAN	N	Subset for alpha = 0.05	
		1	2
T1	30	1.97	
T0	30	2.13	
T3	30	2.30	2.30
T2	30		2.97
Sig.		.686	.125

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.

LAMPIRAN 5

HASIL UJI STATISTIK UJI MUTU HEDONIK TERHADAP AROMA NUGGET KEONG SAWAH

Test of Homogeneity of Variances

AROMA

Levene Statistic	df1	df2	Sig.
2.472	3	116	.065

ANOVA

AROMA

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	1.425	3	.475	.637	.593
Within Groups	86.567	116	.746		
Total	87.992	119			

Descriptives

AROMA

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
T0	30	2.27	.868	.159	1.94	2.59	1	4
T1	30	2.07	.785	.143	1.77	2.36	1	4
T2	30	2.37	1.033	.189	1.98	2.75	1	4
T3	30	2.27	.740	.135	1.99	2.54	1	4
Total	120	2.24	.860	.078	2.09	2.40	1	4

Multiple Comparisons

Dependent Variable: AROMA

Bonferroni

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.200	.223	1.000	-.40	.80
	T2	-.100	.223	1.000	-.70	.50
	T3	.000	.223	1.000	-.60	.60
T1	T0	-.200	.223	1.000	-.80	.40
	T2	-.300	.223	1.000	-.90	.30
	T3	-.200	.223	1.000	-.80	.40
T2	T0	.100	.223	1.000	-.50	.70
	T1	.300	.223	1.000	-.30	.90
	T3	.100	.223	1.000	-.50	.70
T3	T0	.000	.223	1.000	-.60	.60
	T1	.200	.223	1.000	-.40	.80
	T2	-.100	.223	1.000	-.70	.50

Multiple Comparisons

Dependent Variable: AROMA

Tukey HSD

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.200	.223	.807	-.38	.78
	T2	-.100	.223	.970	-.68	.48
	T3	.000	.223	1.000	-.58	.58
T1	T0	-.200	.223	.807	-.78	.38
	T2	-.300	.223	.536	-.88	.28
	T3	-.200	.223	.807	-.78	.38
T2	T0	.100	.223	.970	-.48	.68
	T1	.300	.223	.536	-.28	.88
	T3	.100	.223	.970	-.48	.68
T3	T0	.000	.223	1.000	-.58	.58
	T1	.200	.223	.807	-.38	.78
	T2	-.100	.223	.970	-.68	.48

AROMA

Tukey HSD^a

PERLAKUAN	N	Subset for alpha = 0.05
		1
T1	30	2.07
T0	30	2.27
T3	30	2.27
T2	30	2.37
Sig.		.536

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.

LAMPIRAN 6

HASIL UJI STATISTIK UJI MUTU HEDONIK TERHADAP TEKSTUR NUGGET KEONG SAWAH

Test of Homogeneity of Variances

TEKSTUR

Levene Statistic	df1	df2	Sig.
8.181	3	116	.000

ANOVA

TEKSTUR

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	14.892	3	4.964	6.601	.000
Within Groups	87.233	116	.752		
Total	102.125	119			

Descriptives

TEKSTUR

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
T0	30	4.00	.643	.117	3.76	4.24	2	5
T1	30	3.40	1.192	.218	2.95	3.85	1	5
T2	30	3.93	.640	.117	3.69	4.17	3	5
T3	30	3.17	.874	.160	2.84	3.49	2	4
Total	120	3.63	.926	.085	3.46	3.79	1	5

Multiple Comparisons

Dependent Variable: TEKSTUR

Bonferroni

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.600	.224	.051	.00	1.20
	T2	.067	.224	1.000	-.53	.67
	T3	.833*	.224	.002	.23	1.43
T1	T0	-.600	.224	.051	-1.20	.00
	T2	-.533	.224	.113	-1.13	.07
	T3	.233	.224	1.000	-.37	.83
T2	T0	-.067	.224	1.000	-.67	.53
	T1	.533	.224	.113	-.07	1.13
	T3	.767*	.224	.005	.17	1.37
T3	T0	-.833*	.224	.002	-1.43	-.23
	T1	-.233	.224	1.000	-.83	.37
	T2	-.767*	.224	.005	-1.37	-.17

*. The mean difference is significant at the 0.05 level.

Multiple Comparisons

Dependent Variable: TEKSTUR

Tukey HSD

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.600*	.224	.041	.02	1.18
	T2	.067	.224	.991	-.52	.65
	T3	.833*	.224	.002	.25	1.42
T1	T0	-.600*	.224	.041	-1.18	-.02
	T2	-.533	.224	.086	-1.12	.05
	T3	.233	.224	.725	-.35	.82
T2	T0	-.067	.224	.991	-.65	.52
	T1	.533	.224	.086	-.05	1.12
	T3	.767*	.224	.005	.18	1.35
T3	T0	-.833*	.224	.002	-1.42	-.25
	T1	-.233	.224	.725	-.82	.35
	T2	-.767*	.224	.005	-1.35	-.18

*. The mean difference is significant at the 0.05 level.

TEKSTUR

Tukey HSD^a

PERLAKUAN	N	Subset for alpha = 0.05		
		1	2	3
T3	30	3.17		
T1	30	3.40	3.40	
T2	30		3.93	3.93
T0	30			4.00
Sig.		.725	.086	.991

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.

LAMPIRAN 7

HASIL UJI STATISTIK UJI HEDONIK TERHADAP RASA NUGGET KEONG SAWAH

Test of Homogeneity of Variances

RASA

Levene Statistic	df1	df2	Sig.
1.937	3	116	.127

ANOVA

RASA

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	30.092	3	10.031	6.622	.000
Within Groups	175.700	116	1.515		
Total	205.792	119			

Descriptives

RASA

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
					T0	30		
T1	30	4.47	1.432	.261	3.93	5.00	1	7
T2	30	5.43	1.104	.202	5.02	5.85	2	7
T3	30	4.17	1.367	.250	3.66	4.68	1	7
Total	120	4.79	1.315	.120	4.55	5.03	1	7

Multiple Comparisons

Dependent Variable: RASA

Bonferroni

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.633	.318	.292	-.22	1.49
	T2	-.333	.318	1.000	-1.19	.52
	T3	.933*	.318	.024	.08	1.79
T1	T0	-.633	.318	.292	-1.49	.22
	T2	-.967*	.318	.017	-1.82	-.11
	T3	.300	.318	1.000	-.55	1.15
T2	T0	.333	.318	1.000	-.52	1.19
	T1	.967*	.318	.017	.11	1.82
	T3	1.267*	.318	.001	.41	2.12
T3	T0	-.933*	.318	.024	-1.79	-.08
	T1	-.300	.318	1.000	-1.15	.55
	T2	-1.267*	.318	.001	-2.12	-.41

*. The mean difference is significant at the 0.05 level.

Multiple Comparisons

Dependent Variable: RASA

Tukey HSD

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.633	.318	.197	-.19	1.46
	T2	-.333	.318	.721	-1.16	.49
	T3	.933*	.318	.021	.11	1.76
T1	T0	-.633	.318	.197	-1.46	.19
	T2	-.967*	.318	.015	-1.79	-.14
	T3	.300	.318	.781	-.53	1.13
T2	T0	.333	.318	.721	-.49	1.16
	T1	.967*	.318	.015	.14	1.79
	T3	1.267*	.318	.001	.44	2.09
T3	T0	-.933*	.318	.021	-1.76	-.11
	T1	-.300	.318	.781	-1.13	.53
	T2	-1.267*	.318	.001	-2.09	-.44

*. The mean difference is significant at the 0.05 level.

RASA

Tukey HSD^a

PERLAKUAN	N	Subset for alpha = 0.05		
		1	2	3
T3	30	4.17		
T1	30	4.47	4.47	
T0	30		5.10	5.10
T2	30			5.43
Sig.		.781	.197	.721

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.

LAMPIRAN 8

HASIL UJI STATISTIK UJI HEDONIK TERHADAP WARNA NUGGET KEONG SAWAH

Test of Homogeneity of Variances

WARNA

Levene Statistic	df1	df2	Sig.
3.308	3	116	.023

ANOVA

WARNA

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	22.492	3	7.497	4.120	.008
Within Groups	211.100	116	1.820		
Total	233.592	119			

Descriptives

WARNA

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
T0	30	4.53	.973	.178	4.17	4.90	3	7
T1	30	3.57	1.524	.278	3.00	4.14	1	6
T2	30	4.53	1.332	.243	4.04	5.03	1	7
T3	30	3.80	1.495	.273	3.24	4.36	1	6
Total	120	4.11	1.401	.128	3.86	4.36	1	7

Multiple Comparisons

Dependent Variable: WARNA

Bonferroni

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.967 [*]	.348	.039	.03	1.90
	T2	.000	.348	1.000	-.93	.93
	T3	.733	.348	.224	-.20	1.67
T1	T0	-.967 [*]	.348	.039	-1.90	-.03
	T2	-.967 [*]	.348	.039	-1.90	-.03
	T3	-.233	.348	1.000	-1.17	.70
T2	T0	.000	.348	1.000	-.93	.93
	T1	.967 [*]	.348	.039	.03	1.90
	T3	.733	.348	.224	-.20	1.67
T3	T0	-.733	.348	.224	-1.67	.20
	T1	.233	.348	1.000	-.70	1.17
	T2	-.733	.348	.224	-1.67	.20

*. The mean difference is significant at the 0.05 level.

Multiple Comparisons

Dependent Variable: WARNA

Tukey HSD

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.967 [*]	.348	.032	.06	1.87
	T2	.000	.348	1.000	-.91	.91
	T3	.733	.348	.157	-.17	1.64
T1	T0	-.967 [*]	.348	.032	-1.87	-.06
	T2	-.967 [*]	.348	.032	-1.87	-.06
	T3	-.233	.348	.908	-1.14	.67
T2	T0	.000	.348	1.000	-.91	.91
	T1	.967 [*]	.348	.032	.06	1.87
	T3	.733	.348	.157	-.17	1.64
T3	T0	-.733	.348	.157	-1.64	.17
	T1	.233	.348	.908	-.67	1.14
	T2	-.733	.348	.157	-1.64	.17

*. The mean difference is significant at the 0.05 level.

WARNA

Tukey HSD^a

PERLAKUAN	N	Subset for alpha = 0.05	
		1	2
T1	30	3.57	
T3	30	3.80	3.80
T0	30		4.53
T2	30		4.53
Sig.		.908	.157

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.

LAMPIRAN 9

HASIL UJI STATISTIK UJI HEDONIK TERHADAP AROMA NUGGET KEONG SAWAH

Test of Homogeneity of Variances

AROMA

Levene Statistic	df1	df2	Sig.
.746	3	116	.527

ANOVA

AROMA

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	2.200	3	.733	1.084	.359
Within Groups	78.467	116	.676		
Total	80.667	119			

Descriptives

AROMA

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
					T0	30		
T1	30	4.67	.711	.130	4.40	4.93	4	7
T2	30	4.73	.785	.143	4.44	5.03	4	7
T3	30	4.97	.765	.140	4.68	5.25	4	7
Total	120	4.83	.823	.075	4.68	4.98	3	7

Multiple Comparisons

Dependent Variable: AROMA

Bonferroni

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.300	.212	.963	-.27	.87
	T2	.233	.212	1.000	-.34	.80
	T3	.000	.212	1.000	-.57	.57
T1	T0	-.300	.212	.963	-.87	.27
	T2	-.067	.212	1.000	-.64	.50
	T3	-.300	.212	.963	-.87	.27
T2	T0	-.233	.212	1.000	-.80	.34
	T1	.067	.212	1.000	-.50	.64
	T3	-.233	.212	1.000	-.80	.34
T3	T0	.000	.212	1.000	-.57	.57
	T1	.300	.212	.963	-.27	.87
	T2	.233	.212	1.000	-.34	.80

Multiple Comparisons

Dependent Variable: AROMA

Tukey HSD

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.300	.212	.494	-.25	.85
	T2	.233	.212	.691	-.32	.79
	T3	.000	.212	1.000	-.55	.55
T1	T0	-.300	.212	.494	-.85	.25
	T2	-.067	.212	.989	-.62	.49
	T3	-.300	.212	.494	-.85	.25
T2	T0	-.233	.212	.691	-.79	.32
	T1	.067	.212	.989	-.49	.62
	T3	-.233	.212	.691	-.79	.32
T3	T0	.000	.212	1.000	-.55	.55
	T1	.300	.212	.494	-.25	.85
	T2	.233	.212	.691	-.32	.79

AROMA

Tukey HSD^a

PERLAKUAN	N	Subset for alpha = 0.05
		1
T1	30	4.67
T2	30	4.73
T0	30	4.97
T3	30	4.97
Sig.		.494

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.

LAMPIRAN 10

HASIL UJI STATISTIK UJI HEDONIK TERHADAP TEKSTUR NUGGET KEONG SAWAH

Test of Homogeneity of Variances

TEKSTUR

Levene Statistic	df1	df2	Sig.
3.334	3	116	.022

ANOVA

TEKSTUR

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	10.600	3	3.533	3.456	.019
Within Groups	118.600	116	1.022		
Total	129.200	119			

Descriptives

TEKSTUR

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
T0	30	5.03	.809	.148	4.73	5.34	4	7
T1	30	4.63	1.129	.206	4.21	5.05	2	7
T2	30	5.13	.900	.164	4.80	5.47	2	7
T3	30	4.40	1.163	.212	3.97	4.83	2	7
Total	120	4.80	1.042	.095	4.61	4.99	2	7

Multiple Comparisons

Dependent Variable: TEKSTUR

Bonferroni

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.400	.261	.769	-.30	1.10
	T2	-.100	.261	1.000	-.80	.60
	T3	.633	.261	.101	-.07	1.33
T1	T0	-.400	.261	.769	-1.10	.30
	T2	-.500	.261	.348	-1.20	.20
	T3	.233	.261	1.000	-.47	.93
T2	T0	.100	.261	1.000	-.60	.80
	T1	.500	.261	.348	-.20	1.20
	T3	.733*	.261	.035	.03	1.43
T3	T0	-.633	.261	.101	-1.33	.07
	T1	-.233	.261	1.000	-.93	.47
	T2	-.733*	.261	.035	-1.43	-.03

*. The mean difference is significant at the 0.05 level.

Multiple Comparisons

Dependent Variable: TEKSTUR

Tukey HSD

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.400	.261	.422	-.28	1.08
	T2	-.100	.261	.981	-.78	.58
	T3	.633	.261	.078	-.05	1.31
T1	T0	-.400	.261	.422	-1.08	.28
	T2	-.500	.261	.227	-1.18	.18
	T3	.233	.261	.808	-.45	.91
T2	T0	.100	.261	.981	-.58	.78
	T1	.500	.261	.227	-.18	1.18
	T3	.733*	.261	.029	.05	1.41
T3	T0	-.633	.261	.078	-1.31	.05
	T1	-.233	.261	.808	-.91	.45
	T2	-.733*	.261	.029	-1.41	-.05

*. The mean difference is significant at the 0.05 level.

TEKSTUR

Tukey HSD^a

PERLAKUAN	N	Subset for alpha = 0.05	
		1	2
T3	30	4.40	
T1	30	4.63	4.63
T0	30	5.03	5.03
T2	30		5.13
Sig.		.078	.227

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.

LAMPIRAN 11

HASIL UJI STATISTIK UJI TINGKAT KESUKAAN SECARA KESELURUHAN TERHADAP NUGGET KEONG SAWAH

Test of Homogeneity of Variances

KESELURUHAN

Levene Statistic	df1	df2	Sig.
.792	3	116	.501

ANOVA

KESELURUHAN

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	13.267	3	4.422	4.770	.004
Within Groups	107.533	116	.927		
Total	120.800	119			

Descriptives

KESELURUHAN

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
T0	30	5.30	.837	.153	4.99	5.61	4	7
T1	30	4.87	1.074	.196	4.47	5.27	3	7
T2	30	5.53	.819	.150	5.23	5.84	4	7
T3	30	4.70	1.088	.199	4.29	5.11	2	7
Total	120	5.10	1.008	.092	4.92	5.28	2	7

Multiple Comparisons

Dependent Variable: KESELURUHAN

Bonferroni

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.433	.249	.504	-.23	1.10
	T2	-.233	.249	1.000	-.90	.43
	T3	.600	.249	.104	-.07	1.27
T1	T0	-.433	.249	.504	-1.10	.23
	T2	-.667	.249	.050	-1.33	.00
	T3	.167	.249	1.000	-.50	.83
T2	T0	.233	.249	1.000	-.43	.90
	T1	.667	.249	.050	.00	1.33
	T3	.833*	.249	.006	.17	1.50
T3	T0	-.600	.249	.104	-1.27	.07
	T1	-.167	.249	1.000	-.83	.50
	T2	-.833*	.249	.006	-1.50	-.17

*. The mean difference is significant at the 0.05 level.

Multiple Comparisons

Dependent Variable: KESELURUHAN

Tukey HSD

(I) PERLAKUAN	(J) PERLAKUAN	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
T0	T1	.433	.249	.306	-.21	1.08
	T2	-.233	.249	.784	-.88	.41
	T3	.600	.249	.080	-.05	1.25
T1	T0	-.433	.249	.306	-1.08	.21
	T2	-.667*	.249	.041	-1.31	-.02
	T3	.167	.249	.908	-.48	.81
T2	T0	.233	.249	.784	-.41	.88
	T1	.667*	.249	.041	.02	1.31
	T3	.833*	.249	.006	.19	1.48
T3	T0	-.600	.249	.080	-1.25	.05
	T1	-.167	.249	.908	-.81	.48
	T2	-.833*	.249	.006	-1.48	-.19

*. The mean difference is significant at the 0.05 level.

KESELURUHAN

Tukey HSD^a

PERLAKUAN	N	Subset for alpha = 0.05	
		1	2
T3	30	4.70	
T1	30	4.87	
T0	30	5.30	5.30
T2	30		5.53
Sig.		.080	.784

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 30.000.