

DAFTAR LAMPIRAN

Statistics

	umur	pendidikan	pekerjaan	Asupan Natrium	Pola konsumsi Makanan asin	Pola konsumsi Bumbu penyedap	Aktifitas fisik	IMT
N Valid	889	889	889	889	889	889	889	889
Missing	0	0	0	0	0	0	0	0
Mean	48.65	.23	.06	107.72	.49	.93	.48	1.13
Median	48.00	.00	.00	15.00	.00	1.00	.00	1.00
Std. Deviation	2.932	.500	.231	398.877	.500	.257	.500	.738
Minimum	45	0	0	1	0	0	0	0
Maximum	54	2	1	3872	1	1	1	3

Statistics

hipertension

N Valid	889
Missing	0
Mean	.61
Median	1.00
Std. Deviation	.489
Minimum	0
Maximum	1


Hypertension

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tidak hipertensi	350	39.4	39.4	39.4
	hipertensi	539	60.6	60.6	100.0
	Total	889	100.0	100.0	

kategori umur

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	45-49th	515	57.9	57.9	57.9
	50-54	374	42.1	42.1	100.0
	Total	889	100.0	100.0	

umur responden




Pendidikan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	pendidikan rendah	716	80.5	80.5	80.5
	pendidikan sedang	141	15.9	15.9	96.4
	pendidikan tinggi	32	3.6	3.6	100.0
	Total	889	100.0	100.0	

Pekerjaan

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	bekerja	839	94.4	94.4	94.4
	tidak bekerja	50	5.6	5.6	100.0
	Total	889	100.0	100.0	

asupan natrium



Pola konsumsi makanan asin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	jarang	454	51.1	51.1	51.1
	sering	435	48.9	48.9	100.0
	Total	889	100.0	100.0	

pola konsumsi penyedap

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	jarang	63	7.1	7.1	7.1
	sering	826	92.9	92.9	100.0
	Total	889	100.0	100.0	

Aktifitas fisik

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	cukup	460	51.7	51.7	51.7
	kurang	429	48.3	48.3	100.0
	Total	889	100.0	100.0	

IMT

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	BB kurang	152	17.1	17.1	17.1
	normal	510	57.4	57.4	74.5
	BB lebih	187	21.0	21.0	95.5
	obesitas	40	4.5	4.5	100.0
	Total	889	100.0	100.0	

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
umur responden * hipertension	889	100.0%	0	.0%	889	100.0%
kategori asupan natrium * hipertension	889	100.0%	0	.0%	889	100.0%
kategori makanan asin * hipertension	889	100.0%	0	.0%	889	100.0%
kategori konsumsi penyedap * hipertension	889	100.0%	0	.0%	889	100.0%
kategori aktifitas * hipertension	889	100.0%	0	.0%	889	100.0%
kategori indeks massa tubuh * hipertension	889	100.0%	0	.0%	889	100.0%

Crosstab

			hipertension		Total
			tidak hipertensi	hipertensi	
kategori umur	45-49th	Count	218	297	515
		% within kategori umur	42.3%	57.7%	100.0%
	50-54	Count	132	242	374
		% within kategori umur	35.3%	64.7%	100.0%
Total		Count	350	539	889
		% within kategori umur	39.4%	60.6%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.493 ^a	1	.034		
Continuity Correction ^b	4.203	1	.040		
Likelihood Ratio	4.513	1	.034		
Fisher's Exact Test				.037	.020
Linear-by-Linear Association	4.488	1	.034		
N of Valid Cases ^b	889				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 147.24.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for kategori umur (45-49th / 50-54)	1.346	1.022	1.771
For cohort hipertension = tidak hipertensi	1.199	1.012	1.422
For cohort hipertension = hipertensi	.891	.802	.990
N of Valid Cases	889		

Crosstab

			hipertension		Total
			tidak hipertensi	hipertensi	
kategori asupan natrium	cukup	Count	345	534	879
		% within kategori asupan natrium	39.2%	60.8%	100.0%
	lebih	Count	5	5	10
		% within kategori asupan natrium	50.0%	50.0%	100.0%
Total		Count	350	539	889
		% within kategori asupan natrium	39.4%	60.6%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.479 ^a	1	.489		
Continuity Correction ^b	.134	1	.714		
Likelihood Ratio	.468	1	.494		
Fisher's Exact Test				.527	.351
Linear-by-Linear Association	.478	1	.489		
N of Valid Cases ^b	889				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 3.94.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for kategori asupan natrium (cukup / lebih)	.646	.186	2.248
For cohort hipertension = tidak hipertensi	.785	.420	1.467
For cohort hipertension = hipertensi	1.215	.652	2.263
N of Valid Cases	889		

Crosstab

			hipertension		Total
			tidak hipertensi	hipertensi	
kategori makanan asin	jarang	Count	175	279	454
		% within kategori makanan asin	38.5%	61.5%	100.0%
	sering	Count	175	260	435
		% within kategori makanan asin	40.2%	59.8%	100.0%
Total		Count	350	539	889
		% within kategori makanan asin	39.4%	60.6%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.264 ^a	1	.608		
Continuity Correction ^b	.198	1	.656		
Likelihood Ratio	.264	1	.608		
Fisher's Exact Test				.631	.328
Linear-by-Linear Association	.264	1	.608		
N of Valid Cases ^b	889				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 171.26.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for kategori makanan asin (jarang / sering)	.932	.712	1.220
For cohort hipertension = tidak hipertensi	.958	.814	1.128
For cohort hipertension = hipertensi	1.028	.925	1.143
N of Valid Cases	889		

Crosstab

		hipertension		Total
		tidak hipertensi	hipertensi	
kategori konsumsi penyedap jarang	Count	18	45	63
	% within kategori konsumsi penyedap	28.6%	71.4%	100.0%
sering	Count	332	494	826
	% within kategori konsumsi penyedap	40.2%	59.8%	100.0%
Total	Count	350	539	889
	% within kategori konsumsi penyedap	39.4%	60.6%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	3.312 ^a	1	.069		
Continuity Correction ^b	2.843	1	.092		
Likelihood Ratio	3.444	1	.063		
Fisher's Exact Test				.082	.044
Linear-by-Linear Association	3.309	1	.069		
N of Valid Cases ^b	889				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 24.80.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for kategori konsumsi penyedap (jarang / sering)	.595	.339	1.046
For cohort hipertension = tidak hipertensi	.711	.477	1.060
For cohort hipertension = hipertensi	1.194	1.012	1.410
N of Valid Cases	889		

Crosstab

			hipertension		Total
			tidak hipertensi	hipertensi	
kategori aktifitas	cukup	Count	174	286	460
		% within kategori aktifitas	37.8%	62.2%	100.0%
	kurang	Count	176	253	429
		% within kategori aktifitas	41.0%	59.0%	100.0%
Total		Count	350	539	889
		% within kategori aktifitas	39.4%	60.6%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.952 ^a	1	.329		
Continuity Correction ^b	.823	1	.364		
Likelihood Ratio	.952	1	.329		
Fisher's Exact Test				.337	.182
Linear-by-Linear Association	.951	1	.329		
N of Valid Cases ^b	889				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 168.90.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for kategori aktifitas (cukup / kurang)	.875	.668	1.145
For cohort hipertension = tidak hipertensi	.922	.783	1.085
For cohort hipertension = hipertensi	1.054	.948	1.173
N of Valid Cases	889		

Crosstab

			hipertension		Total
			tidak hipertensi	hipertensi	
kategori indeks massa tubuh	BB kurang	Count	86	66	152
		% within kategori indeks massa tubuh	56.6%	43.4%	100.0%
	normal	Count	203	307	510
		% within kategori indeks massa tubuh	39.8%	60.2%	100.0%
	BB lebih	Count	55	132	187
		% within kategori indeks massa tubuh	29.4%	70.6%	100.0%
	obesitas	Count	6	34	40
		% within kategori indeks massa tubuh	15.0%	85.0%	100.0%
Total		Count	350	539	889
		% within kategori indeks massa tubuh	39.4%	60.6%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	36.619 ^a	3	.000
Likelihood Ratio	37.811	3	.000
Linear-by-Linear Association	35.710	1	.000
N of Valid Cases	889		

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 15.75.

Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	kat_umur	4.493	1	.034
		kat_asupan	.479	1	.489
		kat_asin	.264	1	.608
		kat_penyedap	3.312	1	.069
		kat_aktifitas	.952	1	.329
		imtkurus	22.747	1	.000
		imtgemuk	19.947	1	.000
Overall Statistics			44.458	7	.000

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	1146.613 ^a	.050	.067

a. Estimation terminated at iteration number 4 because parameter estimates changed by less than .001.

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95.0% C.I. for EXP(B)	
							Lower	Upper
kat_umur	.328	.144	5.191	1	.023	1.389	1.047	1.842
kat_asupan	-.601	.660	.831	1	.362	.548	.150	1.997
kat_asin	-.041	.142	.082	1	.775	.960	.727	1.268
kat_penyedap	-.478	.295	2.619	1	.106	.620	.348	1.106
kat_aktifitas	-.237	.143	2.767	1	.096	.789	.596	1.043
imtkurus	-.706	.189	13.928	1	.000	.494	.341	.715
imtgemuk	.605	.176	11.776	1	.001	1.832	1.296	2.589
Constant	.867	.324	7.158	1	.007	2.380		

a. Variable(s) entered on step 1: kat_umur, kat_asupan, kat_asin, kat_penyedap, kat_aktifitas, imtkurus, imtgemuk.

Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	kat_umur	4.493	1	.034
		imtkurus	22.747	1	.000
		imtgemuk	19.947	1	.000
	Overall Statistics		38.917	3	.000

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	1152.557 ^a	.043	.059

a. Estimation terminated at iteration number 4 because parameter estimates changed by less than .001.

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95.0% C.I. for EXP(B)	
							Lower	Upper
kat_umur	.331	.143	5.349	1	.021	1.393	1.052	1.844
imtkurus	-.689	.188	13.457	1	.000	.502	.348	.726
imtgemuk	.598	.176	11.621	1	.001	1.819	1.290	2.566
Constant	.276	.108	6.564	1	.010	1.318		

a. Variable(s) entered on step 1: kat_umur, imtkurus, imtgemuk.