

Deskriptive data

Descriptive Statistics

	N	Mean	Std. Deviation
Sebelum1	7	68,71	7,868
Sesudah1	7	50,14	7,128
Selisih1	7	18,57	8,423
Sebelum2	7	70,29	10,095
Sesudah2	7	49,86	11,908
Selisih2	7	20,43	12,122
Valid N (listwise)	7		

Uji normalitas

Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Sebelum1	,150	7	,200 [*]	,987	7	,988
Sesudah1	,193	7	,200 [*]	,938	7	,618
Sebelum2	,176	7	,200 [*]	,941	7	,652
Sesudah2	,253	7	,198	,900	7	,330

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

Independent Samples Test

	Levene's Test for Equality of Variances		t-test for Equality of Means						
	F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
								Lower	Upper
Equal variances assumed	,946	0,350	-,325	12	,751	-1,571	4,837	-12,111	8,969
Sebelum1 um1 Equal variances not assumed			-,325	11,325	,751	-1,571	4,837	-12,182	9,039

Uji Homogenitas

Uji Hipotesis I

Paired Samples Test

	Paired Differences				t	df	Sig. (2-tailed)	
	Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
				Lower				Upper
Pair 1 Sebelum1 - Sesudah1	18,571	8,423	3,184	10,781	26,362	5,833	6	0,001

Uji Hipotesis II

Paired Samples Test

	Paired Differences					t	df	Sig. (2-tailed)
	Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference				
				Lower	Upper			
Pair 1 Sebelum2 - Sesudah2	20,429	12,122	4,582	9,217	31,640	4,459	6	0,004

Uji Hipotesis III

Independent Samples Test

	Levene's Test for Equality of Variances		t-test for Equality of Means						
	F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
								Lower	Upper
Selisih1	,932	0,353	Equal variances assumed	12	,745	-1,857	5,579	-14,014	10,299
			Equal variances not assumed	10,699	,746	-1,857	5,579	-14,180	10,465