

LAMPIRAN

Analisis Univariat

Statistics

	kategori BBLR	Jarak Hamil	Jenis Kelamin	Umur kehamilan	frekuensi periksa hamil	minum pil besi	pendidikan ibu	Paritas Ibu	komplikasi ada	status ekonomi
N Valid	353	353	353	353	353	353	353	353	353	353
Missing	0	0	0	0	0	0	0	0	0	0

Jenis Kelamin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Laki laki	185	52.4	52.4	52.4
	Perempuan	168	47.6	47.6	100.0
	Total	353	100.0	100.0	

Jarak Hamil

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	0 tahun	91	25.8	25.8	25.8
	kurang dari 2 tahun	27	7.6	7.6	33.4
	lebih dari sama dengan 2 tahun	235	66.6	66.6	100.0
	Total	353	100.0	100.0	

Minum pil besi

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	90-95 hari	92	26.1	26.1	26.1
	0-89 hari	261	73.9	73.9	100.0
	Total	353	100.0	100.0	

Komplikasi

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	tidak ada	321	90.9	90.9	90.9
	ada	32	9.1	9.1	100.0
	Total	353	100.0	100.0	

Umur Kehamilan

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid normal	342	96.9	96.9	96.9
prematur	11	3.1	3.1	100.0
Total	353	100.0	100.0	

Frekuensi periksa hamil

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid lebih sama dengan 4x	326	92.4	92.4	92.4
kurang dari 4x	27	7.6	7.6	100.0
Total	353	100.0	100.0	

Paritas Ibu

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid 1-5 anak	323	91.5	91.5	91.5
lebih dari 5	30	8.5	8.5	100.0
Total	353	100.0	100.0	

Pendidikan ibu

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Tinggi	127	36.0	36.0	36.0
Rendah	226	64.0	64.0	100.0
Total	353	100.0	100.0	

Status ekonomi

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Tinggi	193	54.7	54.7	54.7
Rendah	160	45.3	45.3	100.0
Total	353	100.0	100.0	

Analisis Bivariat

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
jarak Kehamilan * kategori BBLR	353	100.0%	0	.0%	353	100.0%
Umur Kehamilan * kategori BBLR	353	100.0%	0	.0%	353	100.0%
frekuensi periksa hamil * kategori BBLR	353	100.0%	0	.0%	353	100.0%
minum pil besi * kategori BBLR	353	100.0%	0	.0%	353	100.0%
pendidikan ibu * kategori BBLR	353	100.0%	0	.0%	353	100.0%
Paritas Ibu * kategori BBLR	353	100.0%	0	.0%	353	100.0%
komplikasi ada * kategori BBLR	353	100.0%	0	.0%	353	100.0%
jenis kelamin bayi * kategori BBLR	353	100.0%	0	.0%	353	100.0%
status ekonomi * kategori BBLR	353	100.0%	0	.0%	353	100.0%

Jenis kelamin bayi * kategori BBLR

Crosstab

			kategori BBLR		Total
			Tidak BBLR	BBLR	
jenis kelamin bayi	laki-laki	Count	171	14	185
		Expected Count	172.9	12.1	185.0
		% within jenis kelamin bayi	92.4%	7.6%	100.0%
		% within kategori BBLR	51.8%	60.9%	52.4%
	perempuan	Count	159	9	168
		Expected Count	157.1	10.9	168.0
		% within jenis kelamin bayi	94.6%	5.4%	100.0%
		% within kategori BBLR	48.2%	39.1%	47.6%
Total	Count	330	23	353	
	Expected Count	330.0	23.0	353.0	
	% within jenis kelamin bayi	93.5%	6.5%	100.0%	
	% within kategori BBLR	100.0%	100.0%	100.0%	

Chi-Square Tests^d

	Value	Df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	.706 ^a	1	.401	.518	.267	
Continuity Correction ^b	.390	1	.532			
Likelihood Ratio	.713	1	.398	.518	.267	
Fisher's Exact Test				.518	.267	
Linear-by-Linear Association	.704 ^c	1	.401	.518	.267	.122
N of Valid Cases	353					

- a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 10.95.
 b. Computed only for a 2x2 table
 c. The standardized statistic is -.839.
 d. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for jenis kelamin bayi (laki-laki / perempuan)	.691	.291	1.642
For cohort kategori BBLR = Tidak BBLR	.977	.925	1.032
For cohort kategori BBLR = BBLR	1.413	.628	3.178
N of Valid Cases	353		

Jarak Kehamilan * kategori BBLR

Crosstab

			kategori BBLR		Total
			Tidak BBLR	BBLR	
jarak Kehamilan	Tidak beresiko	Count	305	21	326
		Expected Count	304.8	21.2	326.0
		% within jarak Kehamilan	93.6%	6.4%	100.0%
		% within kategori BBLR	92.4%	91.3%	92.4%
	Beresiko	Count	25	2	27
		Expected Count	25.2	1.8	27.0
		% within jarak Kehamilan	92.6%	7.4%	100.0%
		% within kategori BBLR	7.6%	8.7%	7.6%
Total	Count	330	23	353	
	Expected Count	330.0	23.0	353.0	
	% within jarak Kehamilan	93.5%	6.5%	100.0%	
	% within kategori BBLR	100.0%	100.0%	100.0%	

Chi-Square Tests^d

	Value	Df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	.038 ^a	1	.845	1.000	.541	
Continuity Correction ^b	.000	1	1.000			
Likelihood Ratio	.037	1	.848	1.000	.541	
Fisher's Exact Test				.692	.541	
Linear-by-Linear Association	.038 ^c	1	.845	1.000	.541	.289
N of Valid Cases	353					

- a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 1.76.
 b. Computed only for a 2x2 table
 c. The standardized statistic is .195.
 d. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for jarak Kehamilan (Tidak beresiko / Beresiko)	1.162	.258	5.242
For cohort kategori BBLR = Tidak BBLR	1.010	.905	1.128
For cohort kategori BBLR = BBLR	.870	.215	3.513
N of Valid Cases	353		

Minum pil besi * kategori BBLR

Crosstab

			kategori BBLR		Total
			Tidak BBLR	BBLR	
minum pil besi	90-95 hari	Count	90	2	92
		Expected Count	86.0	6.0	92.0
		% within minum pil besi	97.8%	2.2%	100.0%
		% within kategori BBLR	27.3%	8.7%	26.1%
	0-89 hari	Count	240	21	261
		Expected Count	244.0	17.0	261.0
		% within minum pil besi	92.0%	8.0%	100.0%
		% within kategori BBLR	72.7%	91.3%	73.9%
Total		Count	330	23	353
		Expected Count	330.0	23.0	353.0
		% within minum pil besi	93.5%	6.5%	100.0%
		% within kategori BBLR	100.0%	100.0%	100.0%

Chi-Square Tests^d

	Value	Df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	3.851 ^a	1	.050	.082	.035	
Continuity Correction ^b	2.947	1	.086			
Likelihood Ratio	4.719	1	.030	.051	.035	
Fisher's Exact Test				.051	.035	
Linear-by-Linear Association	3.840 ^c	1	.050	.082	.035	.027
N of Valid Cases	353					

- a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5.99.
 b. Computed only for a 2x2 table
 c. The standardized statistic is 1.960.
 d. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for minum pil besi (90-95 hari / 0-89 hari)	3.938	.905	17.134
For cohort kategori BBLR = Tidak BBLR	1.064	1.015	1.115
For cohort kategori BBLR = BBLR	.270	.065	1.130
N of Valid Cases	353		

Komplikasi * kategori BBLR

Crosstab

			kategori BBLR		Total
			Tidak BBLR	BBLR	
komplikasi ada	tidak ada	Count	305	16	321
		Expected Count	300.1	20.9	321.0
		% within komplikasi ada	95.0%	5.0%	100.0%
		% within kategori BBLR	92.4%	69.6%	90.9%
ada	ada	Count	25	7	32
		Expected Count	29.9	2.1	32.0
		% within komplikasi ada	78.1%	21.9%	100.0%
		% within kategori BBLR	7.6%	30.4%	9.1%
Total		Count	330	23	353
		Expected Count	330.0	23.0	353.0
		% within komplikasi ada	93.5%	6.5%	100.0%
		% within kategori BBLR	100.0%	100.0%	100.0%

Chi-Square Tests^d

	Value	Df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	13.629 ^a	1	.000	.002	.002	
Continuity Correction ^b	10.997	1	.001			
Likelihood Ratio	9.320	1	.002	.002	.002	
Fisher's Exact Test				.002	.002	
Linear-by-Linear Association	13.591 ^c	1	.000	.002	.002	.002
N of Valid Cases	353					

- a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 2.08.
 b. Computed only for a 2x2 table
 c. The standardized statistic is 3.687.
 d. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for komplikasi ada (tidak ada / ada)	5.338	2.009	14.183
For cohort kategori BBLR = Tidak BBLR	1.216	1.011	1.463
For cohort kategori BBLR = BBLR	.228	.101	.512
N of Valid Cases	353		

Umur Kehamilan * kategori BBLR

Crosstab

			kategori BBLR		Total
			Tidak BBLR	BBLR	
Umur Kehamilan	normal	Count	323	19	342
		Expected Count	319.7	22.3	342.0
		% within Umur Kehamilan	94.4%	5.6%	100.0%
		% within kategori BBLR	97.9%	82.6%	96.9%
	prematurn	Count	7	4	11
		Expected Count	10.3	.7	11.0
		% within Umur Kehamilan	63.6%	36.4%	100.0%
		% within kategori BBLR	2.1%	17.4%	3.1%
Total		Count	330	23	353
		Expected Count	330.0	23.0	353.0
		% within Umur Kehamilan	93.5%	6.5%	100.0%
		% within kategori BBLR	100.0%	100.0%	100.0%

Chi-Square Tests^d

	Value	Df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	16.607 ^a	1	.000	.003	.003	
Continuity Correction ^b	11.934	1	.001			
Likelihood Ratio	8.914	1	.003	.003	.003	
Fisher's Exact Test				.003	.003	
Linear-by-Linear Association	16.560 ^c	1	.000	.003	.003	.003
N of Valid Cases	353					

- a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is .72.
 b. Computed only for a 2x2 table
 c. The standardized statistic is 4.069.
 d. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Umur Kehamilan (normal / prematur)	9.714	2.614	36.101
For cohort kategori BBLR = Tidak BBLR	1.484	.949	2.322
For cohort kategori BBLR = BBLR	.153	.062	.374
N of Valid Cases	353		

Frekuensi periksa hamil * kategori BBLR

Crosstab

		kategori BBLR		Total	
		Tidak BBLR	BBLR		
frekuensi periksa hamil	lebih sama dengan 4x	Count	303	23	326
		Expected Count	304.8	21.2	326.0
		% within frekuensi periksa hamil	92.9%	7.1%	100.0%
		% within kategori BBLR	91.8%	100.0%	92.4%
	kurang dari 4x	Count	27	0	27
		Expected Count	25.2	1.8	27.0
		% within frekuensi periksa hamil	100.0%	.0%	100.0%
		% within kategori BBLR	8.2%	.0%	7.6%
Total	Count	330	23	353	
	Expected Count	330.0	23.0	353.0	
	% within frekuensi periksa hamil	93.5%	6.5%	100.0%	
	% within kategori BBLR	100.0%	100.0%	100.0%	

Chi-Square Tests^d

	Value	Df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2.038 ^a	1	.153	.238	.151	
Continuity Correction ^b	1.044	1	.307			
Likelihood Ratio	3.790	1	.052	.174	.151	
Fisher's Exact Test				.238	.151	
Linear-by-Linear Association	2.032 ^c	1	.154	.238	.151	.151
N of Valid Cases	353					

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 1.76.

b. Computed only for a 2x2 table

c. The standardized statistic is -1.425.

d. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
For cohort kategori BBLR = Tidak BBLR	.929	.902	.958
N of Valid Cases	353		

Paritas Ibu * kategori BBLR

Crosstab

			kategori BBLR		Total
			Tidak BBLR	BBLR	
Paritas Ibu	1-5 anak	Count	303	20	323
		Expected Count	302.0	21.0	323.0
		% within Paritas Ibu	93.8%	6.2%	100.0%
		% within kategori BBLR	91.8%	87.0%	91.5%
	lebih dari 5	Count	27	3	30
		Expected Count	28.0	2.0	30.0
		% within Paritas Ibu	90.0%	10.0%	100.0%
		% within kategori BBLR	8.2%	13.0%	8.5%
Total	Count	330	23	353	
	Expected Count	330.0	23.0	353.0	
	% within Paritas Ibu	93.5%	6.5%	100.0%	
	% within kategori BBLR	100.0%	100.0%	100.0%	

Chi-Square Tests^d

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	.654 ^a	1	.419	.430	.309	
Continuity Correction ^b	.178	1	.673			
Likelihood Ratio	.576	1	.448	.707	.309	
Fisher's Exact Test				.430	.309	
Linear-by-Linear Association	.652 ^c	1	.420	.430	.309	.189
N of Valid Cases	353					

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 1.95.

b. Computed only for a 2x2 table

c. The standardized statistic is .807.

d. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Paritas Ibu (1-5 anak / lebih dari 5)	1.683	.470	6.029
For cohort kategori BBLR = Tidak BBLR	1.042	.922	1.178
For cohort kategori BBLR = BBLR	.619	.195	1.964
N of Valid Cases	353		

Pendidikan ibu * kategori BBLR**Crosstab**

			kategori BBLR		Total
			Tidak BBLR	BBLR	
pendidikan ibu	Tinggi	Count	122	5	127
		Expected Count	118.7	8.3	127.0
		% within pendidikan ibu	96.1%	3.9%	100.0%
		% within kategori BBLR	37.0%	21.7%	36.0%
	Rendah	Count	208	18	226
		Expected Count	211.3	14.7	226.0
		% within pendidikan ibu	92.0%	8.0%	100.0%
		% within kategori BBLR	63.0%	78.3%	64.0%
Total	Count	330	23	353	
	Expected Count	330.0	23.0	353.0	
	% within pendidikan ibu	93.5%	6.5%	100.0%	
	% within kategori BBLR	100.0%	100.0%	100.0%	

Chi-Square Tests^d

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	2.165 ^a	1	.141	.179	.104	
Continuity Correction ^b	1.555	1	.212			
Likelihood Ratio	2.332	1	.127	.179	.104	
Fisher's Exact Test				.179	.104	
Linear-by-Linear Association	2.159 ^c	1	.142	.179	.104	.063
N of Valid Cases	353					

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 8.27.

b. Computed only for a 2x2 table

c. The standardized statistic is 1.469.

d. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for pendidikan ibu (Tinggi / Rendah)	2.112	.765	5.831
For cohort kategori BBLR = Tidak BBLR	1.044	.991	1.100
For cohort kategori BBLR = BBLR	.494	.188	1.300
N of Valid Cases	353		

Status ekonomi * kategori BBLR**Crosstab**

			kategori BBLR		Total
			Tidak BBLR	BBLR	
status ekonomi	Tinggi	Count	182	11	193
		Expected Count	180.4	12.6	193.0
		% within status ekonomi	94.3%	5.7%	100.0%
		% within kategori BBLR	55.2%	47.8%	54.7%
	Rendah	Count	148	12	160
		Expected Count	149.6	10.4	160.0
		% within status ekonomi	92.5%	7.5%	100.0%
		% within kategori BBLR	44.8%	52.2%	45.3%
Total	Count	330	23	353	
	Expected Count	330.0	23.0	353.0	
	% within status ekonomi	93.5%	6.5%	100.0%	
	% within kategori BBLR	100.0%	100.0%	100.0%	

Chi-Square Tests^d

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)	Point Probability
Pearson Chi-Square	.466 ^a	1	.495	.522	.319	
Continuity Correction ^b	.217	1	.641			
Likelihood Ratio	.463	1	.496	.522	.319	
Fisher's Exact Test				.522	.319	
Linear-by-Linear Association	.464 ^c	1	.496	.522	.319	.135
N of Valid Cases	353					

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 10.42.

b. Computed only for a 2x2 table

c. The standardized statistic is .681.

d. For 2x2 crosstabulation, exact results are provided instead of Monte Carlo results.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for status ekonomi (Tinggi / Rendah)	1.342	.575	3.127
For cohort kategori BBLR = Tidak BBLR	1.019	.964	1.078
For cohort kategori BBLR = BBLR	.760	.345	1.676
N of Valid Cases	353		

T-Test

Notes

Output Created	05-Jan-1980 23:11:56		
Comments			
Input	Data	E:\Tahniyah Nurfajriani S Masba S.Gz\allahu akbar - Copy (2).sav	
	Active Dataset	DataSet1	
	File Label	Aggregated File	
	Filter	<none>	
	Weight	<none>	
	Split File	<none>	
	N of Rows in Working Data File	353	
Missing Value Handling	Definition of Missing Cases Used	User defined missing values are treated as missing. Statistics for each analysis are based on the cases with no missing or out-of-range data for any variable in the analysis.	
Syntax	T-TEST GROUPS=JKbayi(0 1) /MISSING=ANALYSIS /VARIABLES=EA05bayi /CRITERIA=CI(.9500).		
Resources	Processor Time	00:00:00.031	
	Elapsed Time	00:00:00.020	

Group Statistics

jenis kelamin bayi		N	Mean	Std. Deviation	Std. Error Mean
Berat badan [NAMA] ketika lahir	laki-laki	185	3207.87	536.904	39.474
	Perempuan	168	3122.92	480.826	37.096

Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Berat badan [NAMA] ketika lahir	Equal variances assumed	1.213	.271	1.560	351	.120	84.954	54.458	22.151	192.058
	Equal variances not assumed			1.568	350.935	.118	84.954	54.170	21.584	191.491

Analisis Multivariat

Logistic Regression

Tahap 1

Notes

Output Created		12-Jan-1980 21:42:41
Comments		
Input	Data	E:\Tahniyah Nurfajriani S. Masba S.Gz\FIX ALHAMDULILLAH FIX\allahu akbar.sav
	Active Dataset	DataSet1
	File Label	Aggregated File
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	353
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES katBBLR /METHOD=ENTER JarakLahir Umurlahir frekperiksa pilbesi pendidikanibu ParitasIbu komplikasiada JKbayi ekonomi /PRINT=CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.063
	Elapsed Time	00:00:00.037

Case Processing Summary

Unweighted Cases ^a		N	Percent
Selected Cases	Included in Analysis	353	100.0
	Missing Cases	0	.0
	Total	353	100.0
Unselected Cases		0	.0
Total		353	100.0

a. If weight is in effect, see classification table for the total number of cases.

Dependent Variable Encoding

Original Value	Internal Value
Tidak BBLR	0
BBLR	1

Block 1: Method = Enter**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	25.076	9	.003
	Block	25.076	9	.003
	Model	25.076	9	.003

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	145.017 ^a	.069	.179

a. Estimation terminated at iteration number 20 because maximum iterations has been reached. Final solution cannot be found.

Classification Table^a

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 1	kategori BBLR	Tidak BBLR	329	1	99.7
		BBLR	23	0	.0
Overall Percentage					93.2

a. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95.0% C.I. for EXP(B)		
							Lower	Upper	
Step 1 ^a	JarakLahir	.054	.828	.004	1	.948	1.056	.208	5.351
	Umurlahir	2.006	.713	7.920	1	.005	7.435	1.839	30.068
	frekperiksa	-18.634	7.444E3	.000	1	.998	.000	.000	.
	pilbesi	.867	.779	1.237	1	.266	2.379	.517	10.957
	pendidikanibu	.638	.559	1.299	1	.254	1.892	.632	5.663
	ParitasIbu	.402	.699	.330	1	.565	1.494	.380	5.878
	komplikasiada	1.463	.527	7.712	1	.005	4.318	1.538	12.124
	JKbayi	-.352	.473	.554	1	.457	.703	.278	1.777
	ekonomi	-.080	.481	.027	1	.869	.924	.360	2.371
	Constant	-4.013	.827	23.544	1	.000	.018		

a. Variable(s) entered on step 1: JarakLahir, Umurlahir, frekperiksa, pilbesi, pendidikanibu, ParitasIbu, komplikasiada, JKbayi, ekonomi.

Block 0: Beginning Block**Classification Table^{a,b}**

Observed		Predicted			
		kategori BBLR		Percentage Correct	
		Tidak BBLR	BBLR		
Step 0	kategori BBLR	Tidak BBLR	330	0	100.0
		BBLR	23	0	.0
Overall Percentage					93.5

a. Constant is included in the model.

b. The cut value is .500

Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-2.664	.216	152.547	1	.000	.070

Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	JarakLahir	.038	1	.845
		Umurlahir	16.607	1	.000
		frekperiksa	2.038	1	.153
		pilbesi	3.851	1	.050
		pendidikanibu	2.165	1	.141
		ParitasIbu	.654	1	.419
		komplikasiada	13.629	1	.000
		JKbayi	.706	1	.401
		ekonomi	.466	1	.495
Overall Statistics			32.924	9	.000

Logistic Regression

Tahap 2

Notes

Output Created		12-Jan-1980 21:43:43
Comments		
Input	Data	E:\Tahniyah Nurfajriani S. Masba S.Gz\FIX ALHAMDULILLAH FIX\allahu akbar.sav
	Active Dataset	DataSet1
	File Label	Aggregated File
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	353
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES katBBLR /METHOD=ENTER JarakLahir Umurlahir pilbesi pendidikanibu ParitasIbu komplikasiada JKbayi ekonomi /PRINT=CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.046
	Elapsed Time	00:00:00.035

Case Processing Summary

Unweighted Cases ^a		N	Percent
Selected Cases	Included in Analysis	353	100.0
	Missing Cases	0	.0
	Total	353	100.0
Unselected Cases		0	.0
Total		353	100.0

a. If weight is in effect, see classification table for the total number of cases.

Dependent Variable Encoding

Original Value	Internal Value
Tidak BBLR	0
BBLR	1

Block 1: Method = Enter**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	21.254	8	.007
	Block	21.254	8	.007
	Model	21.254	8	.007

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	148.839 ^a	.058	.153

a. Estimation terminated at iteration number 7 because parameter estimates changed by less than .001.

Classification Table^a

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 1	kategori BBLR	Tidak BBLR	329	1	99.7
		BBLR	23	0	.0
Overall Percentage					93.2

a. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95.0% C.I. for EXP(B)		
							Lower	Upper	
Step 1 ^a	JarakLahir	.126	.829	.023	1	.879	1.134	.224	5.754
	Umurlahir	2.115	.713	8.800	1	.003	8.288	2.049	33.520
	pilbesi	.834	.777	1.151	1	.283	2.302	.502	10.565
	pendidikanibu	.584	.559	1.090	1	.297	1.793	.599	5.363
	ParitasIbu	.274	.695	.156	1	.693	1.315	.337	5.131
	komplikasiada	1.470	.525	7.856	1	.005	4.350	1.556	12.159
	JKbayi	-.341	.469	.529	1	.467	.711	.283	1.784
	ekonomi	-.134	.482	.077	1	.781	.875	.340	2.248
	Constant	-4.016	.830	23.421	1	.000	.018		

a. Variable(s) entered on step 1: JarakLahir, Umurlahir, pilbesi, pendidikanibu, ParitasIbu, komplikasiada, JKbayi, ekonomi.

Block 0: Beginning Block**Classification Table^{a,b}**

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 0	kategori BBLR	Tidak BBLR	330	0	100.0
		BBLR	23	0	.0
Overall Percentage					93.5

a. Constant is included in the model.

b. The cut value is .500

Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-2.664	.216	152.547	1	.000	.070

Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	JarakLahir	.038	1	.845
		Umurlahir	16.607	1	.000
		pilbesi	3.851	1	.050
		pendidikanibu	2.165	1	.141
		ParitasIbu	.654	1	.419
		komplikasiada	13.629	1	.000
		JKbayi	.706	1	.401
		ekonomi	.466	1	.495
Overall Statistics			31.076	8	.000

Logistic Regression
Tahap 3

Notes

Output Created		12-Jan-1980 21:43:43
Comments		
Input	Data	E:\Tahniyah Nurfaejriani S. Masba S.Gz\FIX ALHAMDULILLAH FIX\allahu akbar.sav
	Active Dataset	DataSet1
	File Label	Aggregated File
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	353
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES katBBLR /METHOD=ENTER JarakLahir Umurlahir pilbesi pendidikanibu ParitasIbu komplikasiada JKbayi ekonomi /PRINT=CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.046
	Elapsed Time	00:00:00.035

Case Processing Summary

Unweighted Cases ^a		N	Percent
Selected Cases	Included in Analysis	353	100.0
	Missing Cases	0	.0
	Total	353	100.0
Unselected Cases		0	.0
Total		353	100.0

a. If weight is in effect, see classification table for the total number of cases.

Dependent Variable Encoding

Original Value	Internal Value
Tidak BBLR	0
BBLR	1

Block 1: Method = Enter**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	21.254	8	.007
	Block	21.254	8	.007
	Model	21.254	8	.007

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	148.839 ^a	.058	.153

a. Estimation terminated at iteration number 7 because parameter estimates changed by less than .001.

Classification Table^a

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 1	kategori BBLR	Tidak BBLR	329	1	99.7
		BBLR	23	0	.0
Overall Percentage					93.2

a. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95.0% C.I. for EXP(B)		
							Lower	Upper	
Step 1 ^a	JarakLahir	.126	.829	.023	1	.879	1.134	.224	5.754
	Umurlahir	2.115	.713	8.800	1	.003	8.288	2.049	33.520
	pilbesi	.834	.777	1.151	1	.283	2.302	.502	10.565
	pendidikanibu	.584	.559	1.090	1	.297	1.793	.599	5.363
	ParitasIbu	.274	.695	.156	1	.693	1.315	.337	5.131
	komplikasiada	1.470	.525	7.856	1	.005	4.350	1.556	12.159
	JKbayi	-.341	.469	.529	1	.467	.711	.283	1.784
	ekonomi	-.134	.482	.077	1	.781	.875	.340	2.248
	Constant	-4.016	.830	23.421	1	.000	.018		

a. Variable(s) entered on step 1: JarakLahir, Umurlahir, pilbesi, pendidikanibu, ParitasIbu, komplikasiada, JKbayi, ekonomi.

Block 0: Beginning Block**Classification Table^{a,b}**

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 0	kategori BBLR	Tidak BBLR	330	0	100.0
		BBLR	23	0	.0
Overall Percentage					93.5

a. Constant is included in the model.

b. The cut value is .500

Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-2.664	.216	152.547	1	.000	.070

Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	JarakLahir	.038	1	.845
		Umurlahir	16.607	1	.000
		pilbesi	3.851	1	.050
		pendidikanibu	2.165	1	.141
		ParitasIbu	.654	1	.419
		komplikasiada	13.629	1	.000
		JKbayi	.706	1	.401
		ekonomi	.466	1	.495
Overall Statistics			31.076	8	.000

Logistic Regression
Tahap 4

Notes

Output Created		12-Jan-1980 21:46:11
Comments		
Input	Data	E:\Tahniyah Nurfajriani S. Masba S.Gz\FIX ALHAMDULILLAH FIX\allahu akbar.sav
	Active Dataset	DataSet1
	File Label	Aggregated File
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	353
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES katBBLR /METHOD=ENTER Umurlahir pilbesi pendidikanibu komplikasiada JKbayi /PRINT=CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.062
	Elapsed Time	00:00:00.031

Case Processing Summary

Unweighted Cases ^a		N	Percent
Selected Cases	Included in Analysis	353	100.0
	Missing Cases	0	.0
	Total	353	100.0
Unselected Cases		0	.0
Total		353	100.0

a. If weight is in effect, see classification table for the total number of cases.

Dependent Variable Encoding

Original Value	Internal Value
Tidak BBLR	0
BBLR	1

Block 1: Method = Enter**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	21.037	5	.001
	Block	21.037	5	.001
	Model	21.037	5	.001

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	149.055 ^a	.058	.151

a. Estimation terminated at iteration number 7 because parameter estimates changed by less than .001.

Classification Table^a

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 1	kategori BBLR	Tidak BBLR	329	1	99.7
		BBLR	22	1	4.3
Overall Percentage					93.5

a. The cut value is .500

Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)	95.0% C.I. for EXP(B)	
								Lower	Upper
Step 1 ^a	Umurlahir	2.086	.697	8.947	1	.003	8.054	2.053	31.601
	pilbesi	.836	.777	1.160	1	.282	2.308	.504	10.571
	pendidikanibu	.564	.546	1.065	1	.302	1.758	.602	5.129
	komplikasiada	1.468	.521	7.945	1	.005	4.339	1.564	12.037
	JKbayi	-.364	.467	.607	1	.436	.695	.279	1.735
	Constant	-4.015	.821	23.946	1	.000	.018		

a. Variable(s) entered on step 1: Umurlahir, pilbesi, pendidikanibu, komplikasiada, JKbayi.

Block 0: Beginning Block**Classification Table^{a,b}**

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 0	kategori BBLR	Tidak BBLR	330	0	100.0
		BBLR	23	0	.0
	Overall Percentage				93.5

a. Constant is included in the model.

b. The cut value is .500

Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-2.664	.216	152.547	1	.000	.070

Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	Umurlahir	16.607	1	.000
		pilbesi	3.851	1	.050
		pendidikanibu	2.165	1	.141
		komplikasiada	13.629	1	.000
		JKbayi	.706	1	.401
	Overall Statistics		30.894	5	.000

Logistic Regression
Tahap 5

Notes

Output Created		12-Jan-1980 21:46:30
Comments		
Input	Data	E:\Tahniyah Nurfajriani S. Masba S.Gz\FIX ALHAMDULILLAH FIX\allahu akbar.sav
	Active Dataset	DataSet1
	File Label	Aggregated File
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	353
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES katBBLR /METHOD=ENTER Umurlahir pilbesi pendidikanibu komplikasiada /PRINT=CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.063
	Elapsed Time	00:00:00.043

Case Processing Summary

Unweighted Cases ^a		N	Percent
Selected Cases	Included in Analysis	353	100.0
	Missing Cases	0	.0
	Total	353	100.0
Unselected Cases		0	.0
Total		353	100.0

a. If weight is in effect, see classification table for the total number of cases.

Dependent Variable Encoding

Original Value	Internal Value
Tidak BBLR	0
BBLR	1

Block 1: Method = Enter**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	20.418	4	.000
	Block	20.418	4	.000
	Model	20.418	4	.000

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	149.675 ^a	.056	.147

a. Estimation terminated at iteration number 7 because parameter estimates changed by less than .001.

Classification Table^a

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 1	kategori BBLR	Tidak BBLR	329	1	99.7
		BBLR	22	1	4.3
Overall Percentage					93.5

a. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95.0% C.I. for EXP(B)	
							Lower	Upper
Step 1 ^a Umurlahir	2.061	.701	8.630	1	.003	7.852	1.986	31.052
pilbesi	.876	.774	1.282	1	.257	2.402	.527	10.946
pendidikanibu	.549	.546	1.012	1	.314	1.732	.594	5.052
komplikasiada	1.461	.523	7.803	1	.005	4.312	1.546	12.021
Constant	-4.188	.797	27.610	1	.000	.015		

a. Variable(s) entered on step 1: Umurlahir, pilbesi, pendidikanibu, komplikasiada.

Block 0: Beginning Block**Classification Table^{a,b}**

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 0	kategori BBLR	Tidak BBLR	330	0	100.0
		BBLR	23	0	.0
	Overall Percentage				93.5

a. Constant is included in the model.

b. The cut value is .500

Variables in the Equation

		B	S.E.	Wald	df	Sig.	Exp(B)
Step 0	Constant	-2.664	.216	152.547	1	.000	.070

Variables not in the Equation

			Score	df	Sig.
Step 0	Variables	Umurlahir	16.607	1	.000
		pilbesi	3.851	1	.050
		pendidikanibu	2.165	1	.141
		komplikasiada	13.629	1	.000
	Overall Statistics		30.537	4	.000

Logistic Regression
Tahap 6

Notes

Output Created		12-Jan-1980 21:46:57
Comments		
Input	Data	E:\Tahniyah Nurfajriani S. Masba S.Gz\FIX ALHAMDULILLAH FIX\allahu akbar.sav
	Active Dataset	DataSet1
	File Label	Aggregated File
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	353
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES katBBLR /METHOD=ENTER Umurlahir pilbesi komplikasiada /PRINT=CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.078
	Elapsed Time	00:00:00.046

Case Processing Summary

Unweighted Cases ^a		N	Percent
Selected Cases	Included in Analysis	353	100.0
	Missing Cases	0	.0
	Total	353	100.0
Unselected Cases		0	.0
Total		353	100.0

a. If weight is in effect, see classification table for the total number of cases.

Dependent Variable Encoding

Original Value	Internal Value
Tidak BBLR	0
BBLR	1

Block 1: Method = Enter**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	19.329	3	.000
	Block	19.329	3	.000
	Model	19.329	3	.000

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	150.763 ^a	.053	.139

a. Estimation terminated at iteration number 6 because parameter estimates changed by less than .001.

Classification Table^a

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 1	kategori BBLR	Tidak BBLR	329	1	99.7
		BBLR	22	1	4.3
Overall Percentage					93.5

a. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95.0% C.I. for EXP(B)	
							Lower	Upper
Step 1 ^a Umurlahir	2.054	.704	8.507	1	.004	7.800	1.962	31.015
pilbesi	.967	.768	1.583	1	.208	2.629	.583	11.856
komplikasiada	1.503	.523	8.263	1	.004	4.497	1.613	12.534
Constant	-3.875	.718	29.159	1	.000	.021		

a. Variable(s) entered on step 1: Umurlahir, pilbesi, komplikasiada.

Block 0: Beginning Block**Classification Table^{a,b}**

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 0	kategori BBLR	Tidak BBLR	330	0	100.0
		BBLR	23	0	.0
Overall Percentage					93.5

a. Constant is included in the model.

b. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 0 Constant	-2.664	.216	152.547	1	.000	.070

Variables not in the Equation

	Score	df	Sig.
Step 0 Variables Umurlahir	16.607	1	.000
Pilbesi	3.851	1	.050
komplikasiada	13.629	1	.000
Overall Statistics	29.762	3	.000

Logistic Regression

Tahap 7

Notes

Output Created		12-Jan-1980 21:47:19
Comments		
Input	Data	E:\Tahniyah Nurfajriani S. Masba S.Gz\FIX ALHAMDULILLAH FIX\allahu akbar.sav
	Active Dataset	DataSet1
	File Label	Aggregated File
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	353
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES katBBLR /METHOD=ENTER Umurlahir komplikasiada /PRINT=CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.047
	Elapsed Time	00:00:00.035

Case Processing Summary

Unweighted Cases ^a		N	Percent
Selected Cases	Included in Analysis	353	100.0
	Missing Cases	0	.0
	Total	353	100.0
Unselected Cases		0	.0
Total		353	100.0

a. If weight is in effect, see classification table for the total number of cases.

Dependent Variable Encoding

Original Value	Internal Value
Tidak BBLR	0
BBLR	1

Block 1: Method = Enter**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	17.367	2	.000
	Block	17.367	2	.000
	Model	17.367	2	.000

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	152.725 ^a	.048	.126

a. Estimation terminated at iteration number 6 because parameter estimates changed by less than .001.

Classification Table^a

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 1	kategori BBLR	Tidak BBLR	329	1	99.7
		BBLR	22	1	4.3
Overall Percentage					93.5

a. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95.0% C.I. for EXP(B)	
							Lower	Upper
Step 1 ^a Umurlahir	2.226	.703	10.022	1	.002	9.259	2.334	36.725
komplikasiada	1.643	.518	10.049	1	.002	5.170	1.872	14.277
Constant	-3.107	.276	126.609	1	.000	.045		

a. Variable(s) entered on step 1: Umurlahir, komplikasiada.

Block 0: Beginning Block**Classification Table^{a,b}**

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 0	kategori BBLR	Tidak BBLR	330	0	100.0
		BBLR	23	0	.0
Overall Percentage					93.5

a. Constant is included in the model.

b. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	
Step 0	Constant	-2.664	.216	152.547	1	.000	.070

Variables not in the Equation

	Score	df	Sig.		
Step 0	Variables	Umurlahir	16.607	1	.000
		komplikasiada	13.629	1	.000
Overall Statistics			28.615	2	.000

Logistic Regression

Tahap 8

Notes

Output Created		12-Jan-1980 21:47:42
Comments		
Input	Data	E:\Tahniah Nurfaejriani S. Masba S.Gz\FIX ALHAMDULILLAH FIX\allahu akbar.sav
	Active Dataset	DataSet1
	File Label	Aggregated File
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	353
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES katBBLR /METHOD=ENTER Umurlahir /PRINT=CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.062

Notes

Output Created		12-Jan-1980 21:47:42
Comments		
Input	Data	E:\Tahniyah Nurfajriani S. Masba S.Gz\FIX ALHAMDULILLAH FIX\allahu akbar.sav
	Active Dataset	DataSet1
	File Label	Aggregated File
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	353
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing
Syntax		LOGISTIC REGRESSION VARIABLES katBBLR /METHOD=ENTER Umurlahir /PRINT=CI(95) /CRITERIA=PIN(0.05) POUT(0.10) ITERATE(20) CUT(0.5).
Resources	Processor Time	00:00:00.062
	Elapsed Time	00:00:00.034

Case Processing Summary

Unweighted Cases ^a		N	Percent
Selected Cases	Included in Analysis	353	100.0
	Missing Cases	0	.0
	Total	353	100.0
Unselected Cases		0	.0
Total		353	100.0

a. If weight is in effect, see classification table for the total number of cases.

Dependent Variable Encoding

Original Value	Internal Value
Tidak BBLR	0
BBLR	1

Block 1: Method = Enter**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	8.914	1	.003
	Block	8.914	1	.003
	Model	8.914	1	.003

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	161.179 ^a	.025	.065

a. Estimation terminated at iteration number 5 because parameter estimates changed by less than .001.

Classification Table^a

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 1	kategori BBLR	Tidak BBLR	330	0	100.0
		BBLR	23	0	.0
Overall Percentage					93.5

a. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95.0% C.I. for EXP(B)		
							Lower	Upper	
Step 1 ^a	Umurlahir	2.274	.670	11.523	1	.001	9.714	2.614	36.101
	Constant	-2.833	.236	144.042	1	.000	.059		

a. Variable(s) entered on step 1: Umurlahir.

Block 0: Beginning Block**Classification Table^{a,b}**

Observed			Predicted		
			kategori BBLR		Percentage Correct
			Tidak BBLR	BBLR	
Step 0	kategori BBLR	Tidak BBLR	330	0	100.0
		BBLR	23	0	.0
Overall Percentage					93.5

a. Constant is included in the model.

b. The cut value is .500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 0 Constant	-2.664	.216	152.547	1	.000	.070

Variables not in the Equation

	Score	df	Sig.
Step 0 Variables Umurlahir	16.607	1	.000
Overall Statistics	16.607	1	.000