


Frequencies

Statistics

TOTAL_ISPA		
N	Valid	72
	Missing	0
Mean		3.46
Median		4.00
Mode		4
Std. Deviation		.963
Variance		.928
Minimum		1
Maximum		5

TOTAL_ISPA				
	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1	6	8.3	8.3
	2	5	6.9	15.3
	3	12	16.7	31.9
	4	48	66.7	98.6
	5	1	1.4	100.0
	Total	72	100.0	100.0



Frequencies


Statistics

ISPA_BALITA

N	Valid	72
	Missing	0

ISPA_BALITA

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak ISPA	23	31.9	31.9	31.9
	ISPA	49	68.1	68.1	100.0
	Total	72	100.0	100.0	



Frequencies

Statistics

TOTAL_PERILAKU


N	Valid	72
	Missing	0
Mean		4.97
Median		5.00
Mode		3 ^a
Std. Deviation		2.129
Variance		4.534
Minimum		1
Maximum		9

a. Multiple modes exist. The smallest value is shown

TOTAL_PERILAKU

	Frequency	Percent	Valid Percent	Cumulative Percent
1	2	2.8	2.8	2.8
2	9	12.5	12.5	15.3
3	12	16.7	16.7	31.9
Valid 4	6	8.3	8.3	40.3
5	12	16.7	16.7	56.9
6	11	15.3	15.3	72.2
7	10	13.9	13.9	86.1

Histogram




8	8	11.1	11.1	97.2
9	2	2.8	2.8	100.0
Total	72	100.0	100.0	

Frequencies

Statistics		
perilaku_merokok_keluarga		
N	Valid	72
	Missing	0

perilaku_merokok_keluarga				
	Frequency	Percent	Valid Percent	Cumulative Percent
ringan	29	40.3	40.3	40.3
Valid berat	43	59.7	59.7	100.0
Total	72	100.0	100.0	



Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
perilaku_merokok_keluarga * ISPA_BALITA	72	100.0%	0	0.0%	72	100.0%

perilaku_merokok_keluarga * ISPA_BALITA Crosstabulation

			ISPA_BALITA		Total
			Tidak ISPA	ISPA	
perilaku_merokok_keluarga	ringan	Count	5	24	29
		% within perilaku_merokok_keluarga	17.2%	82.8%	100.0%
	berat	Count	18	25	43
		% within perilaku_merokok_keluarga	41.9%	58.1%	100.0%
	Total	Count	23	49	72
		% within perilaku_merokok_keluarga	31.9%	68.1%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.829 ^a	1	.028		
Continuity Correction ^b	3.763	1	.052		
Likelihood Ratio	5.081	1	.024		
Fisher's Exact Test				.039	.024
Linear-by-Linear Association	4.762	1	.029		
N of Valid Cases	72				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 9.26.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for perilaku_merokok_keluarga (ringan / berat)	.289	.093	.903
For cohort ISPA_BALITA = Tidak ISPA	.412	.172	.985
For cohort ISPA_BALITA = ISPA	1.423	1.051	1.928
N of Valid Cases	72		

