

HASIL OUTPUT SPSS

Pendidikan Responden

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Pendidikan * Kematian Neonatal	90	100.0%	0	.0%	90	100.0%

Pendidikan * Kematian Neonatal Crosstabulation

			Kematian Neonatal		
			Neonatal Mati	Neonatal Hidup	Total
Pendidikan SD	Count	1	0	1	
	% within Kematian Neonatal	2.2%	.0%	1.1%	
SMP	Count	5	2	7	
	% within Kematian Neonatal	11.1%	4.4%	7.8%	
SMU	Count	20	10	30	
	% within Kematian Neonatal	44.4%	22.2%	33.3%	
D3	Count	3	11	14	
	% within Kematian Neonatal	6.7%	24.4%	15.6%	
S1	Count	16	22	38	
	% within Kematian Neonatal	35.6%	48.9%	42.2%	
Total	Count	45	45	90	
	% within Kematian Neonatal	100.0%	100.0%	100.0%	

Pekerjaan Responden

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Pekerjaan * Kematian Neonatal	90	100.0%	0	.0%	90	100.0%

Pekerjaan * Kematian Neonatal Crosstabulation

			Kematian Neonatal		Total
			Neonatal Mati	Neonatal Hidup	
Pekerjaan	Swasta/ BUMN	Count % within Kematian Neonatal	15 33.3%	20 44.4%	35 38.9%
	PNS	Count % within Kematian Neonatal	5 11.1%	5 11.1%	10 11.1%
	Wiraswasta	Count % within Kematian Neonatal	4 8.9%	5 11.1%	9 10.0%
	Ibu Rumah Tangga	Count % within Kematian Neonatal	21 46.7%	15 33.3%	36 40.0%
Total		Count % within Kematian Neonatal	45 100.0%	45 100.0%	90 100.0%

Jenis Kelamin Bayi**Crosstabs****Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Jenis Kelamin * Kematian Neonatal	90	100.0%	0	.0%	90	100.0%

Jenis Kelamin * Kematian Neonatal Crosstabulation

			Kematian Neonatal		
			Neonatal Mati	Neonatal Hidup	Total
Jenis Kelamin	Laki-laki	Count	27	23	50
		% within Kematian Neonatal	60.0%	51.1%	55.6%
	Perempuan	Count	18	22	40
		% within Kematian Neonatal	40.0%	48.9%	44.4%
	Total	Count	45	45	90
		% within Kematian Neonatal	100.0%	100.0%	100.0%

Variabel Usia Ibu**Crosstabs****Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Kelompok Usia Ibu * Kematian Neonatal	90	100.0%	0	.0%	90	100.0%

Kelompok Usia Ibu * Kematian Neonatal Crosstabulation

		Kematian Neonatal		Total
		Neonatal Mati	Neonatal Hidup	
Kelompok Usia Ibu	Beresiko (>35 Tahun atau <20 Tahun)	Count 11 24.4%	Count 7 15.6%	Count 18 20.0%
	Tidak Beresiko (20-35 Tahun)	Count 34 75.6%	Count 38 84.4%	Count 72 80.0%
Total		Count 45 100.0%	Count 45 100.0%	Count 90 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.111 ^a	1	.292		
Continuity Correction ^b	.625	1	.429		
Likelihood Ratio	1.119	1	.290		
Fisher's Exact Test				.430	.215
Linear-by-Linear Association	1.099	1	.295		
N of Valid Cases	90				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 9.00.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Kelompok Usia Ibu (Beresiko (>35 Tahun atau <20 Tahun) / Tidak Beresiko (20-35 Tahun))	1.756	.612	5.042
For cohort Kematian Neonatal = Neonatal Mati	1.294	.832	2.014
For cohort Kematian Neonatal = Neonatal Hidup	.737	.397	1.368
N of Valid Cases	90		

Variabel Paritas

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Kelompok Paritas * Kematian Neonatal	90	100.0%	0	.0%	90	100.0%

Kelompok Paritas * Kematian Neonatal Crosstabulation

			Kematian Neonatal		Total
			Neonatal Mati	Total	
Kelompok Paritas	Berisiko (0 dan >3)	Count % within Kematian Neonatal	20 44.4%	39 43.3%	39 43.3%
	Tidak Berisiko (1-3)	Count % within Kematian Neonatal	25 55.6%	51 56.7%	51 56.7%
Total		Count % within Kematian Neonatal	45 100.0%	90 100.0%	90 100.0%

Kelompok Paritas * Kematian Neonatal Crosstabulation

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.045 ^a	1	.832		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.045	1	.832		
Fisher's Exact Test				1.000	.500
Linear-by-Linear Association	.045	1	.832		
N of Valid Cases	90				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 19.50.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Kelompok Paritas (Berisiko (0 dan >3) / Tidak Berisiko (1-3))	1.095	.475	2.521
For cohort Kematian Neonatal = Neonatal Mati	1.046	.691	1.584
For cohort Kematian Neonatal = Neonatal Hidup	.956	.628	1.454
N of Valid Cases	90		

Variabel Jarak Kelahiran

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Jarak Kelahiran * Kematian Neonatal	90	100.0%	0	.0%	90	100.0%

Jarak Kelahiran * Kematian Neonatal Crosstabulation

			Kematian Neonatal		Total
			Neonatal Mati	Neonatal Hidup	
Jarak Kelahiran	Berisiko (<2 Thn atau >4 Thn)	Count % within Kematian Neonatal	36 80.0%	34 75.6%	70 77.8%
	Tidak Berisiko (2-4 Thn)	Count % within Kematian Neonatal	9 20.0%	11 24.4%	20 22.2%
Total		Count % within Kematian Neonatal	45 100.0%	45 100.0%	90 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.257 ^a	1	.612		
Continuity Correction ^b	.064	1	.800		
Likelihood Ratio	.257	1	.612		
Fisher's Exact Test				.800	.400
Linear-by-Linear Association	.254	1	.614		
N of Valid Cases	90				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 10.00.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Jarak Kelahiran (Berisiko (<2 Thn atau >4 Thn) / Tidak Berisiko (2-4 Thn))	1.294	.477	3.510
For cohort Kematian Neonatal = Neonatal Mati	1.143	.669	1.952
For cohort Kematian Neonatal = Neonatal Hidup	.883	.555	1.404
N of Valid Cases	90		

Variabel Riwayat Kesehatan Ibu

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Riwayat Kesehatan Ibu * Kematian Neonatal	90	100.0%	0	.0%	90	100.0%

Riwayat Kesehatan Ibu * Kematian Neonatal
Crosstabulation

			Kematian Neonatal		Total
			Neonatal Mati	Neonatal Hidup	
Riwayat Kesehatan Ibu	Berisiko (Sakit)	Count % within Kematian Neonatal	43 95.6%	32 71.1%	75 83.3%
	Tidak Berisiko (Sehat)	Count % within Kematian Neonatal	2 4.4%	13 28.9%	15 16.7%
Total		Count % within Kematian Neonatal	45 100.0%	45 100.0%	90 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	9.680 ^a	1	.002		
Continuity Correction ^b	8.000	1	.005		
Likelihood Ratio	10.633	1	.001		
Fisher's Exact Test				.003	.002
Linear-by-Linear Association	9.572	1	.002		
N of Valid Cases	90				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 7.50.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Riwayat Kesehatan Ibu (Berisiko (Sakit) / Tidak Berisiko (Sehat))	8.734	1.840	41.459
For cohort Kematian Neonatal = Neonatal Mati	4.300	1.166	15.855
For cohort Kematian Neonatal = Neonatal Hidup	.492	.354	.684
N of Valid Cases	90		

Variabel Masa Gestasi

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Masa Gestasi * Kematian Neonatal	90	100.0%	0	.0%	90	100.0%

Masa Gestasi * Kematian Neonatal Crosstabulation

			Kematian Neonatal		Total
			Neonatal Mati	Neonatal Hidup	
Masa Gestasi	Berisiko (<37 atau >42 minggu)	Count % within Kematian Neonatal	36 80.0%	5 11.1%	41 45.6%
	Tidak Berisiko (37-42 minggu)	Count % within Kematian Neonatal	9 20.0%	40 88.9%	49 54.4%
Total		Count % within Kematian Neonatal	45 100.0%	45 100.0%	90 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	43.051 ^a	1	.000		
Continuity Correction ^b	40.319	1	.000		
Likelihood Ratio	47.623	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	42.573	1	.000		
N of Valid Cases	90				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 20.50.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Masa Gestasi (Berisiko (<37 atau >42 minggu) / Tidak Berisiko (37-42 minggu))	32.000	9.810	104.381
For cohort Kematian Neonatal = Neonatal Mati	4.780	2.620	8.721
For cohort Kematian Neonatal = Neonatal Hidup	.149	.065	.343
N of Valid Cases	90		

Variabel Berat Lahir Bayi

Crosstabs

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Berat Lahir * Kematian Neonatal	90	100.0%	0	.0%	90	100.0%

Berat Lahir * Kematian Neonatal Crosstabulation

			Kematian Neonatal		Total
			Neonatal Mati	Neonatal Hidup	
Berat Lahir	Berisiko (BB <2500 gram)	Count % within Kematian Neonatal	34 75.6%	7 15.6%	41 45.6%
	Tidak Berisiko (BB >2500 gram)	Count % within Kematian Neonatal	11 24.4%	38 84.4%	49 54.4%
	Total	Count % within Kematian Neonatal	45 100.0%	45 100.0%	90 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2- sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	32.658 ^a	1	.000		
Continuity Correction ^b	30.284	1	.000		
Likelihood Ratio	35.101	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	32.295	1	.000		
N of Valid Cases	90				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 20.50.

b. Computed only for a 2x2 table

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for Berat Lahir (Berisiko (BB <2500 gram) / Tidak Berisiko (BB >2500 gram))	16.779	5.845	48.168
For cohort Kematian Neonatal = Neonatal Mati	3.694	2.156	6.330
For cohort Kematian Neonatal = Neonatal Hidup	.220	.110	.439
N of Valid Cases	90		