

Case Processing Summary

| | Cases | | | | | |
|---|-------|---------|---------|---------|-------|---------|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| Kategori Asupan Energi * Status Gizi | 66 | 100.0% | 0 | .0% | 66 | 100.0% |

| | | | | | | |
|--|----|--------|---|-----|----|--------|
| Kategori Asupan Protein * Status Gizi | 66 | 100.0% | 0 | .0% | 66 | 100.0% |
| Kategori Asupan Lemak * Status Gizi | 66 | 100.0% | 0 | .0% | 66 | 100.0% |
| Kategori Asupan KH * Status Gizi | 66 | 100.0% | 0 | .0% | 66 | 100.0% |

Kategori Asupan Energi * Status Gizi Crosstabulation

| Count | | Status Gizi | | |
|------------------------|------------|-------------|--------------|-------|
| | | Normal | Tidak normal | Total |
| Kategori Asupan Energi | Baik | 25 | 8 | 33 |
| | Tidak baik | 31 | 2 | 33 |
| Total | | 56 | 10 | 66 |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|--------------------|----|--------------------------|-------------------------|-------------------------|
| Pearson Chi-Square | 4.243 ^a | 1 | .039 | | |
| Continuity Correction ^b | 2.946 | 1 | .086 | | |
| Likelihood Ratio | 4.499 | 1 | .034 | | |
| Fisher's Exact Test | | | | .082 | .041 |
| Linear-by-Linear Association | 4.179 | 1 | .041 | | |
| N of Valid Cases ^b | 66 | | | | |

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5,00.

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|--------------------|----|--------------------------|-------------------------|-------------------------|
| Pearson Chi-Square | 4.243 ^a | 1 | .039 | | |
| Continuity Correction ^b | 2.946 | 1 | .086 | | |
| Likelihood Ratio | 4.499 | 1 | .034 | | |
| Fisher's Exact Test | | | | .082 | .041 |
| Linear-by-Linear Association | 4.179 | 1 | .041 | | |
| N of Valid Cases ^b | 66 | | | | |

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5,00.

b. Computed only for a 2x2 table

Kategori Asupan Protein * Status Gizi

Crosstab

| Count | | Status Gizi | | |
|--------|----|-----------------------------------|--------------|-------|
| | | Normal | Tidak normal | Total |
| | | Kategori Asupan Protein ≥ 80 | 54 | 8 |
| < 80 | 2 | 2 | 4 | |
| Total | 56 | 10 | 66 | |

hi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|--------------------|----|--------------------------|-------------------------|-------------------------|
| Pearson Chi-Square | 4.022 ^a | 1 | .045 | | |
| Continuity Correction ^b | 1.654 | 1 | .198 | | |
| Likelihood Ratio | 2.915 | 1 | .088 | | |
| Fisher's Exact Test | | | | .016 | .016 |
| Linear-by-Linear Association | 3.961 | 1 | .047 | | |
| N of Valid Cases ^b | 66 | | | | |

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,61.

hi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|--------------------|----|--------------------------|-------------------------|-------------------------|
| Pearson Chi-Square | 4.022 ^a | 1 | .045 | | |
| Continuity Correction ^b | 1.654 | 1 | .198 | | |
| Likelihood Ratio | 2.915 | 1 | .088 | | |
| Fisher's Exact Test | | | | .016 | .016 |
| Linear-by-Linear Association | 3.961 | 1 | .047 | | |
| N of Valid Cases ^b | 66 | | | | |

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,61.

b. Computed only for a 2x2 table

Kategori Asupan Lemak * Status Gizi

Crosstab

| Count | | Status Gizi | | |
|--------|----|---------------------------------|--------------|-------|
| | | Normal | Tidak normal | Total |
| | | Kategori Asupan Lemak ≥ 80 | 55 | 8 |
| < 80 | 1 | 2 | 3 | |
| Total | 56 | 10 | 66 | |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|--------------------|----|--------------------------|-------------------------|-------------------------|
| Pearson Chi-Square | 6.488 ^a | 1 | .011 | | |
| Continuity Correction ^b | 2.969 | 1 | .085 | | |
| Likelihood Ratio | 4.367 | 1 | .037 | | |
| Fisher's Exact Test | | | | .038 | .038 |
| Linear-by-Linear Association | 6.389 | 1 | .011 | | |
| N of Valid Cases ^b | 66 | | | | |

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,45.

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|--------------------|----|--------------------------|-------------------------|-------------------------|
| Pearson Chi-Square | 6.488 ^a | 1 | .011 | | |
| Continuity Correction ^b | 2.969 | 1 | .085 | | |
| Likelihood Ratio | 4.367 | 1 | .037 | | |
| Fisher's Exact Test | | | | .038 | .038 |
| Linear-by-Linear Association | 6.389 | 1 | .011 | | |
| N of Valid Cases ^b | 66 | | | | |

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,45.

b. Computed only for a 2x2 table

Kategori Asupan KH * Status Gizi

Crosstab

| Count | | Status Gizi | | |
|-------|------------|--------------------|--------------|-------|
| | | Normal | Tidak normal | Total |
| | | Kategori Asupan KH | Baik | 56 |
| | Tidak Baik | 0 | 2 | 2 |
| Total | | 56 | 10 | 66 |

Chi-Square Tests

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|------------------------------------|---------------------|----|--------------------------|-------------------------|-------------------------|
| Pearson Chi-Square | 11.550 ^a | 1 | .010 | | |
| Continuity Correction ^b | 5.746 | 1 | .017 | | |
| Likelihood Ratio | 7.917 | 1 | .005 | | |
| Fisher's Exact Test | | | | .021 | .021 |
| Linear-by-Linear Association | 11.375 | 1 | .001 | | |
| N of Valid Cases ^b | 66 | | | | |

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is ,30.

b. Computed only for a 2x2 table

Jadwal menu makan santri

| No | Hari | Waktu Makan | | |
|----|--------|--------------------------------|--|---|
| | | Pagi | Siang | Malam |
| 1 | Minggu | Nasi ,Semur tahu cokelat | Nasi, Tumis taube + Bakwan jagung | Nasi, Tumis sawi putih + Tahu coklat goreng |
| 2 | Senin | Nasi, Sayur tempe | Nasi, Tumis sawi putih + Semur tahu cokelat | Nasi, Sayur sop + ayam santan |
| 3 | Selasa | Nasi, Bihun goreng | Nasi , Pecel sayur + Tahu cokelat bumbu kacang | Nasi, Sayur asem + orek tempe |
| 4 | Rabu | Nasi uduk + kerupuk udang | Nasi, Tumis labu siam + orek tempe cabe ijo | Nasi , Bening bayam+ Tongkol balado |
| 5 | Kamis | Nasi, Mie goreng | Nasi, Sayur asem + Teri kacang balado | Nasi, Sayur sop + Telur balado |
| 6 | Jumat | Nasi, Kentang balado | Nasi, Terong balado + orek tempe | Nasi, Tumis labu siam + telur dadar |
| 7 | Sabtu | Nasi goreng + kerupuk udang | Nasi , Tumis kangkung + tongol balado | Nasi, Sayur kol toge + tempe orek |