

Kuesioner

Hubungan tingkat Kepadatan Lalat dengan Kejadian Diare pada Balita yang Bermukim disekitar Tempat Pengolahan Hasil Perikanan Tradisional (PHPT), 2016

Nama Pewawancara :

Tanggal wawancara :

Responden

Nomor Responden

Nama :

Alamat :

Karakteristik Balita

1. Nama Balita :
2. Jenis Kelamin :
3. Umur Balita : Bulan

A. Pemberian Asi Eksklusif

- a. Pemberian Asi Eksklusif : 1. Iya
2. Tidak
- b. Jika Iya, Sampai Usia Berapa Bulan : < 6 bulan
≥ 6 bulan
- c. Kapan Ibu mulai memberikan Makanan tambahan : 1. < 6 bulan
2. ≥ 6 bulan

B. Kejadian Diare

1. Apakah 2 minggu terakhir anak ibu mengalami sakit berak-berak?
 - a. Tidak
 - b. Iya

[illegible]

Frequencies

Statistics

Kepadatan Lalat1

N	Valid	75
	Missing	0
Mean		,77
Std. Error of Mean		,049
Median		1,00
Mode		1
Std. Deviation		,421
Minimum		0
Maximum		1
Sum		58
	25	1,00
Percentiles	50	1,00
	75	1,00

Kepadatan Lalat1

	Frequency	Percent	Valid Percent	Cumulative Percent
0 - 5	17	22,7	22,7	22,7
Valid 6 - >21	58	77,3	77,3	100,0
Total	75	100,0	100,0	

Statistics

Diare1

N	Valid	75
	Missing	0
Mean		,73
Std. Error of Mean		,051
Median		1,00
Mode		1
Std. Deviation		,445
Minimum		0
Maximum		1
Sum		55
	25	,00
Percentiles	50	1,00
	75	1,00

Diare1

	Frequency	Percent	Valid Percent	Cumulative Percent
Diare	55	73,3	73,3	73,3
Valid Tidak Diare	20	26,7	26,7	100,0
Total	75	100,0	100,0	

Crosstabs**Case Processing Summary**

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
KL1 * Diare1	75	100,0%	0	0,0%	75	100,0%

Kepadatan Lalat1 * Diare1 Crosstabulation

			Diare1		Total
			Diare	Tidak Diare	
KL1	Tidak Padat	Count	4	13	17
		Expected Count	12,5	4,5	17,0
		% within KL1	23,5%	76,5%	100,0%
	Padat	Count	51	7	58
		Expected Count	42,5	15,5	58,0
		% within KL1	87,9%	12,1%	100,0%
Total	Count		55	20	75
	Expected Count		55,0	20,0	75,0
	% within KL1		73,3%	26,7%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	27,883 ^a	1	,000	,000	,000
Continuity Correction ^b	24,687	1	,000		
Likelihood Ratio	25,715	1	,000		
Fisher's Exact Test					
Linear-by-Linear Association	27,511	1	,000		
N of Valid Cases	75				

a. 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,53.

b. Computed only for a 2x2 table

Symmetric Measures

	Value	Approx. Sig.
Phi	,610	,000
Nominal by Cramer's V	,610	,000
Nominal Contingency Coefficient	,521	,000
N of Valid Cases	75	

a. Not assuming the null hypothesis.

b. Using the asymptotic standard error assuming the null hypothesis

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for KL1 (Tidak Padat / Padat)	23,679	6,010	93,288
For cohort Diare1 = Tidak Diare	6,336	3,014	13,320
For cohort Diare1 = Diare	,268	,113	,634
N of Valid Cases	75		

Frequencies

Statistics

Umur1

N	Valid	75
	Missing	0
Mean		,51
Median		1,00
Mode		1
Std. Deviation		,503
Minimum		0
Maximum		1
Sum		38
	25	,00
Percentiles	50	1,00
	75	1,00

Umur1

	Frequency	Percent	Valid Percent	Cumulative Percent
<36	37	49,3	49,3	49,3
Valid >37	38	50,7	50,7	100,0
Total	75	100,0	100,0	

Frequencies

Statistics

JenisKelamin1

N	Valid	75
	Missing	0
Mean		,47
Median		,00
Mode		0
Std. Deviation		,502
Minimum		0
Maximum		1
Sum		35
	25	,00
Percentiles	50	,00
	75	1,00

JenisKelamin1

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	LakiLaki	40	53,3	53,3	53,3
	Perempuan	35	46,7	46,7	100,0
	Total	75	100,0	100,0	



Gambar : 2.7

Tempat Pembuangan Akhir Sampah



Gambar : 2.8

Sisa-Sisa Kulit Ikan



Gamabar : 2.9

Fly grill



Gambar : 2.10

Lagi Mewawancarai Responden



Gambar : 2.11

Jeroan Sisa-Sisa Ikan



Gambar : 2.12

Jeroan Sisa-Sisa Ikan